

Title	Sequential Point Estimation of the Powers of an Exponential Scale Parameter (Asymptotic Statistical Theory)
Author(s)	Ali, Muktar; Isogai, Eiichi
Citation	数理解析研究所講究録 (2003), 1308: 75-83
Issue Date	2003-02
URL	<a href="http://hdl.handle.net/2433/42849">http://hdl.handle.net/2433/42849</a>
Right	
Type	Departmental Bulletin Paper
Textversion	publisher

## Sequential Point Estimation of the Powers of an Exponential Scale Parameter

Muktar ALI (アリ ムクター)

Graduate School of Science and Technology, Niigata University  
Niigata 950-2181, Japan

Eiichi ISOGAI (磯貝 英一)

Department of Mathematics, Niigata University  
Niigata 950-2181, Japan

### 1. Introduction

Let  $X_1, X_2, X_3, \dots$  be independent and identically distributed (i.i.d.) random variables with the probability density function (p.d.f.)

$$f_{\mu, \sigma}(x) = \frac{1}{\sigma} \exp\left(-\frac{x - \mu}{\sigma}\right) I_{(x \geq \mu)},$$

where both  $\mu \in (-\infty, \infty)$  and  $\sigma \in (0, \infty)$  are unknown and  $I(A)$  denotes the indicator function of the set  $A$ . For any given  $r \neq 0$  we want to estimate the power of the scale parameter  $\sigma^r$ . Let  $\delta_n = \delta_n(X_1, \dots, X_n)$  be an estimator of  $\sigma^r$  based on a random sample  $X_1, \dots, X_n$  of size  $n$ . Then as a loss function we use the squared error loss defined by  $L_n = (\delta_n - \sigma^r)^2$ . The risk associated with the estimator  $\delta_n$  is given by  $R_n = R_n(\delta_n) = E(L_n)$ . Let  $w > 0$  be a preassigned error bound for the risk. We want to find the smallest sample size  $n = n_0$  which satisfies that  $R_n \leq w$ . In bounded risk point estimation problems the error bound  $w$  is assumed to be sufficiently small, so that we suppose that the sample size  $n$  is sufficiently large. For  $n \geq 2$  set

$$T_n = \min\{X_1, \dots, X_n\} \quad \text{and} \quad \sigma_n = (n-1)^{-1} \sum_{i=1}^n (X_i - T_n).$$

We can show that  $c_n \sigma_n^r$  with  $c_n = (n-1)^r \Gamma(n-1) / \Gamma(r+n-1)$  is the uniformly minimum variance unbiased estimator of  $\sigma^r$  provided  $n > \max\{1, 1-r\}$ . We use, however,  $\sigma_n^r$  as an estimator of  $\sigma^r$ , because  $\sigma_n^r$  is approximately equal to  $c_n \sigma_n^r$  for large  $n$  and the calculation of  $\sigma_n^r$  is easier than that of  $c_n \sigma_n^r$ . Our goal is to find an asymptotically smallest sample size  $n_0$  satisfying that  $R_n = E(\sigma_n^r - \sigma^r)^2 \leq w$ .

Estimation of  $\sigma$  and  $\sigma^2$  are of great importance. For  $r = 1$ , namely, for the estimation of the standard deviation by  $\sigma_n$ , Isogai, Saito and Uno (1999) dealt with this bounded risk point estimation problem. Minimum risk point estimation problems for  $r = 1$  were considered by Mukhopadhyay and Ekwo (1987), Ghosh and Mukhopadhyay (1989) and Isogai and Uno (1994). Starr and Woodroffe (1972) treated the same problem for  $r = 2$ . Uno and Isogai (2002) proposed a fully sequential procedure for the estimation of  $\sigma^r$  with normal scale parameter  $\sigma$ . For a review one may refer to Mukhopadhyay (1988), Ghosh and Sen (1991) and Ghosh, Mukhopadhyay and Sen (1997). Sometimes, it is of interest to measure mean  $\lambda = \mu + \sigma$  in  $\sigma$ -units and hence to estimate  $\mu\sigma^{-1}$ . For a normal distribution with mean  $\mu$  and variance  $\sigma^2$  both unknown, Sriram (1990) considered the sequential point estimation problem for  $\mu\sigma^{-1}$  by using an estimator of  $\sigma^r$  with  $r = -1$ .

We shall now compute the risk  $R_n = E(\sigma_n^r - \sigma^r)^2$  to find  $n_0$ . We can show that for  $n > \max\{2, 1-2r\}$

$$R_n < \infty \quad \text{and} \quad R_n = r^2 \sigma^{2r} n^{-1} + O(n^{-2}) \quad \text{as} \quad n \rightarrow \infty.$$

Ignoring the order term above, we can find the asymptotically smallest sample size  $n_0$  satisfying that  $R_n \leq w$ . Suppose

$$r^2 \sigma^{2r} n^{-1} \leq w, \quad \text{or equivalently,} \quad n \geq \frac{r^2 \sigma^{2r}}{w} = n^* \quad (\text{say}). \quad (1.1)$$

For simplicity  $n^*$  is assumed to be an integer. Then  $n_0 = n^*$  is the asymptotically best fixed sample size if  $\sigma$  is known. Unfortunately, the asymptotically best fixed sample size procedure  $n_0$  cannot be used since  $\sigma$  is unknown. Further there is no fixed sample size procedure satisfying our condition. Thus we need to find a sequential sampling rule.

In Section 2 we shall propose a fully sequential procedure for this estimation problem and give two theorems concerning the second order approximation to its average sample

size and risk associated with our procedure. We shall also consider a class of sequential estimators derived on the basis of the idea of bias-correction and compare them from the point of view of risk. In Section 3 we shall provide brief simulation results.

## 2. Results

In this section we shall propose a fully sequential procedure  $N$  motivated by the form of  $n^*$  in (1.1) and give two theorems concerning the second order approximation to its average sample size  $E(N)$  and risk  $R_N = E(\sigma_N^r - \sigma^r)^2$  as  $w \rightarrow 0$ . We shall also consider a class of sequential estimators, including the ordinary estimator  $\sigma_N^r$ , based on the idea of bias-correction. The comparison will be made from the point of view of risk. It will turn out that we can find an appropriate sequential estimator to reduce the risk associated with the ordinary one.

In this paper we propose the stopping rule defined by

$$N = N_w(r) = \inf \left\{ n \geq m : n \geq \frac{r^2 \sigma_n^{2r}}{w} l_n \right\}, \quad (2.1)$$

where  $m$  is a starting sample size satisfying that  $m > \max\{2, 1 - 2r\}$  and  $l_x$  is a given positive function of  $x$  on  $(0, \infty)$  such that

$$l_x = 1 + \frac{l_0}{x} + o\left(\frac{1}{x}\right) \quad \text{as } x \rightarrow \infty \quad \text{with a constant } l_0.$$

We can show that  $P(N_w(r) < \infty) = 1$  for all  $w > 0$  and  $r \neq 0$ . Once the sampling stops at the  $N$ th stage, we estimate  $\sigma^r$  by  $\sigma_N^r$ . Then the risk associated with  $\sigma_N^r$  is given by  $R_N = E(\sigma_N^r - \sigma^r)^2$ . The following two theorems are concerned with the second order approximation to the average sample size and risk.

**Theorem 2.1.** If  $m > m_1(r)$ , then as  $w \rightarrow 0$

$$E(N) = n^* + \rho + l_0 - r(2r + 1) + o(1),$$

where

$$m_1(r) = \begin{cases} 1 + 6r & \text{if } r > 0 \\ 1 - 2r & \text{if } r < 0 \end{cases}$$

and  $\rho$  is a certain constant with  $0 \leq \rho \leq \frac{1}{2} + 2r^2$ .

**Theorem 2.2.** If  $m > m_2(r)$ , then as  $w \rightarrow 0$

$$n^* \left( \frac{R_N}{w} - 1 \right) = 11r^2 + 8r + 1 + \frac{3}{4}(r-1)^2 - \rho - l_0 + o(1),$$

where

$$m_2(r) = \begin{cases} \max \{1 + 10r, 7 + 8r\} & \text{if } r > 0 \\ 7 - 14r & \text{if } r < 0. \end{cases}$$

**Remark 2.1.** (i) If we take an arbitrary constant  $l_0$  such that

$$l_0 > 11r^2 + 8r + 1 + \frac{3}{4}(r-1)^2 - \rho, \quad (2.2)$$

then from Theorem 2.2 we have that  $R_N < w$  for sufficiently small  $w > 0$ . Thus our condition on the risk is satisfied for sufficiently small  $w$ .

(ii) Theorem 2.1 of Isogai, Saito and Uno (1999) with  $a = 0$  and  $b = 1$  is the same as Theorems 2.1 and 2.2 with  $r = 1$  except for the condition on the starting sample size. The difference of this condition is caused by the fact that this paper deals with all powers. Further the methods of the proofs are different.

We shall here evaluate the bias of  $\sigma_N^r$ .

**Proposition 2.1.** If  $m > m_3(r)$ , then as  $w \rightarrow 0$

$$E(\sigma_N^r) - \sigma^r = -\frac{1}{2} \text{sign}(r)(3r+1)(n^*)^{-1/2} w^{1/2} + o(w),$$

where

$$\text{sign}(r) = \begin{cases} 1 & \text{if } r > 0 \\ -1 & \text{if } r < 0 \end{cases} \quad \text{and} \quad m_3(r) = \begin{cases} \max\{1 + 6r, 3 + 3r\} & \text{if } r > 0 \\ 3 - 5r & \text{if } r < 0. \end{cases}$$

Taking Proposition 2.1 into account, we consider a class of sequential estimators  $\{\sigma_N^r(k), k \in (-\infty, \infty)\}$  for  $\sigma^r$  defined by

$$\sigma_N^r(k) = \sigma_N^r + k N^{-1/2} w^{1/2}.$$

Then we get the following proposition concerning the bias of  $\sigma_N^r(k)$ .

**Proposition 2.2.** If  $m > m_3(r)$ , then as  $w \rightarrow 0$

$$E(\sigma_N^r(k)) = \sigma^r + \{k - \frac{1}{2}\text{sign}(r)(3r + 1)\}(n^*)^{-1/2}w^{1/2} + o(w).$$

For  $k = \frac{1}{2}\text{sign}(r)(3r + 1)$ ,  $\sigma_N^r(k)$  is a second-order asymptotically unbiased estimator.

We shall now compare the risk of  $\sigma_N^r(k)$  with that of  $\sigma_N^r$ . Let  $R_N(k) = E(\sigma_N^r(k) - \sigma^r)^2$ .

**Theorem 2.3.** If  $m > m_2(r)$ , then as  $w \rightarrow 0$

$$\frac{n^*}{w}(R_N(k) - R_N) = k^2 - \text{sign}(r)(5r + 1)k + o(1).$$

**Remark 2.2** Let  $k = \frac{1}{2}\text{sign}(r)(5r + 1)$  for  $r \neq 0$ . Then we have

$$R_N(k) < R_N \quad \text{for sufficiently small } w > 0 \quad \text{if } r \neq -\frac{1}{5}.$$

Thus bias-correction is asymptotically effective to reduce the risk for all  $r \neq 0$  with using  $\sigma_N^r + \frac{1}{2}\text{sign}(r)(5r + 1)N^{-1/2}w^{1/2}$  which is not a second-order asymptotically unbiased estimator.

### 3. Simulation Results

We shall give brief simulation results which are based on 100,000 repetitions. We choose the constant  $l_0$  satisfying the inequality in (2.2) in Tables 1–3. In Tables 4 and 5 we choose  $l_0$  such that the average sample size  $E(N)$  approximately equals the optimal one  $n^*$ . Since we do not know any approximate value of  $\rho$  between 0 and  $\frac{1}{2} + 2r^2$ , we use here  $\rho = 0$  or  $\rho = \frac{1}{2} + 2r^2$  as  $\rho$ . From these simulation results we might need to improve the stopping rule  $N$  in (2.1).

**Table 1.**  $\rho = 0$ ,  $l_0 > 11r^2 + 8r + 1 + \frac{3}{4}(r - 1)^2 - \rho$ 

$n^* = 100$	$r = -1$	$r = 1$	$r = 2$
$\mu = 0, \sigma = 1$	$w = 0.01$	$w = 0.01$	$w = 0.04$
	$m = 22$	$m = 16$	$m = 24$
$l_n = 1 + l_0/n$	$l_0 = 8$	$l_0 = 21$	$l_0 = 62$
	$k = 2$	$k = 3$	$k = 5.5$
$E(N)$	108.630260	116.346200	137.330710
$E(\sigma_N^r)$	0.991965	0.983436	0.951511
$E(\sigma_N^r(k))$	1.011400	1.011567	1.048849
$R_N/w$	0.963153	0.984520	0.949165
$R_N(k)/w$	0.934459	0.921439	0.800041
$n^*(R_N(k) - R_N)/w$	-2.869400	-6.308015	-14.912360

**Table 2.**  $r = -1$ ,  $\rho = \frac{1}{2} + 2r^2 = 2.5$ ,  $l_0 > 11r^2 + 8r + 1 + \frac{3}{4}(r - 1)^2 - \rho = 4.5$ 

$n^* = 40$	$\sigma = 0.5$	$\sigma = 1$	$\sigma = 2$
$\mu = 0, l_0 = 5$	$w = 0.1$	$w = 0.025$	$w = 0.00625$
$m = 22, k = 2$	$\sigma^r = 2$	$\sigma^r = 1$	$\sigma^r = 0.5$
$E(N)$	45.752540	45.700810	45.738710
$E(\sigma_N^r)$	1.965109	0.982012	0.491141
$E(\sigma_N^r(k))$	2.061243	1.030103	0.515186
$R_N/w$	0.918271	0.914489	0.925572
$R_N(k)/w$	0.865381	0.859792	0.870948
$n^*(R_N(k) - R_N)/w$	-2.115595	-2.187878	-2.184982

**Table 3.**  $r = 1$ ,  $\rho = \frac{1}{2} + 2r^2 = 2.5$ ,  $l_0 > 11r^2 + 8r + 1 + \frac{3}{4}(r - 1)^2 - \rho = 17.5$ 

$n^* = 40$	$\sigma = 0.5$	$\sigma = 1$	$\sigma = 2$
$\mu = 0$ , $l_0 = 18$	$w = 0.00625$	$w = 0.025$	$w = 0.1$
$m = 16$ , $k = 3$	$\sigma^r = 0.5$	$\sigma^r = 1$	$\sigma^r = 2$
$E(N)$	52.049260	52.088870	52.200480
$E(\sigma_N^r)$	0.481075	0.962627	1.928253
$E(\sigma_N^r(k))$	0.514786	1.030025	2.062843
$R_N/w$	1.004848	1.002715	0.984277
$R_N(k)/w$	0.870552	0.871007	0.862504
$n^*(R_N(k) - R_N)/w$	-5.371849	-5.268332	-4.870901

**Table 4.**  $r = -1$ ,  $\rho = \frac{1}{2} + 2r^2 = 2.5$ ,  $l_0 \leq r(2r + 1) - \rho = -1.5$ 

$n^* = 40$	$\sigma = 0.5$	$\sigma = 1$	$\sigma = 2$
$\mu = 0$ , $l_0 = -2$	$w = 0.1$	$w = 0.025$	$w = 0.00625$
$m = 22$ , $k = 2$	$\sigma^r = 2$	$\sigma^r = 1$	$\sigma^r = 0.5$
$E(N)$	39.236610	39.303210	39.232770
$E(\sigma_N^r)$	1.951794	0.976822	0.487953
$E(\sigma_N^r(k))$	2.056594	1.029155	0.514149
$R_N/w$	1.034014	1.018080	1.026389
$R_N(k)/w$	0.942045	0.931146	0.935053
$n^*(R_N(k) - R_N)/w$	-3.678757	-3.477360	-3.653437



**Table 5.**  $r = 1$ ,  $\rho = \frac{1}{2} + 2r^2 = 2.5$ ,  $l_0 \leq r(2r + 1) - \rho = 0.5$

$n^* = 40$	$\sigma = 0.5$	$\sigma = 1$	$\sigma = 2$
$\mu = 0, l_0 = 0$	$w = 0.00625$	$w = 0.025$	$w = 0.1$
$m = 16, k = 3$	$\sigma^r = 0.5$	$\sigma^r = 1$	$\sigma^r = 2$
$E(N)$	37.647620	37.596930	37.616820
$E(\sigma_N^r)$	0.468557	0.936478	1.873344
$E(\sigma_N^r(k))$	0.509244	1.017909	2.036156
$R_N/w$	1.470935	1.470930	1.473031
$R_N(k)/w$	1.107491	1.103551	1.106660
$n^*(R_N(k) - R_N)/w$	-14.537765	-14.695145	-14.654831

## References

- Ghosh, M. and Mukhopadhyay, N. (1989). Sequential estimation of the percentiles of exponential and normal distributions. *South African Statist. J.*, **23**, 251–268.
- Ghosh, M., Mukhopadhyay, N. and Sen, P.K. (1997). *Sequential Estimation*. John Wiley & Sons, Inc., New York.
- Ghosh, B.K. and Sen, P.K. (1991). *Handbook of Sequential Analysis*, Marcel Dekker, Inc., New York.
- Isogai, E., Saito, K. and Uno, C. (1999). Sequential bounded risk point estimation of parameters of a negative exponential distribution. *Sequential Anal.*, **18**, 217–232.
- Isogai, E. and Uno, C. (1994). Sequential estimation of a parameter of an exponential distribution. *Ann. Inst. Statist. Math.*, **46**, 77–82.
- Mukhopadhyay, N. (1988). Sequential estimation problems for negative exponential populations. *Commun. Statist., Theory and Methods (Reviews Section)*, **17**, 2471–2506.
- Mukhopadhyay, N. and Ekwo, M.E. (1987). A note on minimum risk point estimation of the shape parameter of a Pareto distribution. *Calcutta Statist. Assoc. Bull.*, **36**, 69–78.

Sriram, T.N. (1990). Sequential estimation of ratio of normal parameters. *Statist. Plann. Inference*, **26**, 305–324.

Starr, N. and Woodroffe, M. (1972). Further remarks on sequential estimation: The exponential case. *Ann. Math. Statist.*, **43**, 1147–1154.

Uno, C. and Isogai, E. (2002). Sequential point estimation of the powers of a normal scale parameter. *Metrika*, **55**, 215–232.