

Title	Temperature dependent sex determination and hatching performance of green turtle ( <i>Chelonia mydas</i> ) at Chendor Rookery on the east coast of Peninsular Malaysia
Author(s)	ABDULLAH, SYED; ISMAIL, MAZLAN
Citation	Proceedings of the International Symposium on SEASTAR2000 and Bio-logging Science (The 5th SEASTAR2000 Workshop) (2004): 11-15
Issue Date	2004
URL	<a href="http://hdl.handle.net/2433/44096">http://hdl.handle.net/2433/44096</a>
Right	
Type	Conference Paper
Textversion	publisher

# Temperature dependent sex determination and hatching performance of green turtles (*Chelonia mydas*) at Chendor Rookery on the east coast of Peninsular Malaysia.

SYED ABDULLAH AND MAZLAN ISMAIL

*Marine Fishery Resources Development and Management Department (MFRDMD, SEAFDEC),*

*Fisheries Department of Malaysia, Kuala Terengganu, Terengganu, Malaysia.*

*e-mail: sabdullah@mfrdmd.org.my*

## ABSTRACT

The green turtle (*Chelonia mydas*) is a major species of sea turtle in Malaysia. This study was conducted in 1997 and 1998 at a rookery of green turtles in Pahang. Hatching success as well as sex ratio between male and female was mainly affected by incubation temperatures. The objectives of this study are to determine the optimum temperature that could produce the best hatching performance and to obtain the suitable incubation temperature for producing the female hatching as well male of green turtle. The results showed that the temperatures between 25°C to 33.0°C are suitable for egg incubation. The temperatures between 27.2°C to 29.2°C produced more male turtle hatchlings at the Chendor rookeries, Pahang. More than 80% male hatchlings were produced at the temperatures between 27.2°C to 28.2°C. Whereas female hatchlings are dominant at the temperature above 29.5°C. Temperature between 32.0°C to 33.0°C produce 100% female hatchlings. Thus determination on incubation temperature is vital to formulate the successful hatchery management of sea turtle in Malaysia as well as in the Southeast Asian region.

## INTRODUCTION

There are four species of sea turtles (*Dermochelys coriacea*, *Chelonia mydas*, *Eretmochelys imbricata* and *Lepidochelys olivacea*) nesting in Peninsular Malaysia. Out of four species green turtle is the major species in Malaysia as well as in the Southeast Asian region. It is well-known that the temperatures are a major factor on the success of turtle egg incubation. The temperature would affect the sex determination of turtle hatchlings as well as hatching performance. Therefore it is essential to obtain the information on the suitable temperatures of turtle hatching. The success of the clutch of eggs depends upon interaction of a number of factors, such as salinity, humidity, temperatures, gas flow, rainfall, tidal inundation, erosion and predation (Hendrickson, 1958; Ragotzie, 1959; Bustard and Greenham, 1968; Ackerman, 1974; Kreamer and Bell, 1980; Parmenter, 1980). The temperature also influences the sexual differential of *Caretta caretta* (Yntema and Mrosovsky, 1979, 1980; Reed, 1980; Limpus et al., 1983) *Chelonia mydas* (Miller and Limpus, 1981; Morreale et al., 1982; Limpus et al., 1983). Two main objectives of this study; firstly to determine suitable incubation temperatures which could produce the best hatching performance and secondly to determine the temperatures that produce the male as

well as female green turtles hatchlings at Chendor rookery, Pahang. The results could promote successful hatchery management.

## MATERIALS AND METHODS

The study was conducted in 1997 and 1998 at Chendor rookery, Pahang, Malaysia during two nesting seasons. The first season is from May to June, and the second season is from August to September, respectively. The sample eggs were collected from three different mothers in each sampling period. The eggs were kept in a Coleman container and cooled with ice-cubes and transported to a laboratory at MFRDMD, Kuala Terengganu for incubation. At the laboratory each clutch were divided randomly into groups of 10 eggs. Each group was set in a plastic container holding heat sterilized sand that had been collected from where the turtle normally nest. The containers were placed in constant temperature incubators such that the eggs from the clutch were set at different temperatures between 24°C to 34°C. The temperature of each incubator was monitored at least twice daily and minor adjustment was made as required to maintain each at its specified temperatures. Histology process was undertaken to determine the sexing of turtle hatchlings. Criteria for sexing the

hatchlings followed the methods described in Mrosovsky et al. (1984) and Dutton et al. (1985).

### Results and Discussion

Table 1 shows that the hatching rate in first season at Chendor, Pahang obtained higher than 90% at the temperatures between 27°C to 32°C. Whereas 100 % hatching rate occurred at the temperatures 27°C and 30°C.

Table 2 shows that the hatching rate in the second season obtained higher than 90%, at the temperatures between 25°C and 33°C. This result is quite similar with the sample in first season. The temperature at 26°C, 28°C, 30°C and 31°C has produced an optimum

hatchling rate (100%). Handling and transferring technique will also affect the eggs hatching. These experiments indicate that handling and transferring as well as incubation temperature were the major factor should be considered in hatchery management.

Figure 3 indicates that temperature values between 27.2 °C to 29.0°C produce more male turtle hatchlings at the Chendor rookeries, Pahang. Whereas female hatchlings are dominant at the temperature above 29.5 °C. Temperature values between 32.0 °C to 33.0°C produce 100 % female hatchlings.

Table 1: Hatchling performance in first season (May – June, 1997) of green turtles of Chendor (CH), Pahang.

Temperature (°C)	CH1(Mother - 1)		CH2 (Mother - 2)		CH3 (Mother - 3)		% Average
	eggs incubated	% Hatching	eggs incubated	% Hatching	eggs incubated	% Hatching	
24°C	10	0	10	0	10	100	33.33
25°C	10	0	10	80	10	100	60.00
26°C	10	0	10	90	10	100	63.33
27°C	10	100	10	100	10	100	100.00
28°C	10	100	10	90	10	100	96.67
29°C	10	90	10	100	10	90	93.33
30°C	10	100	10	100	10	100	100.00
31°C	10	80	10	100	10	100	93.33
32°C	10	90	10	90	10	90	90.00
33°C	10	100	10	20	10	80	36.67
34°C	10	70	10	20	0	80	56.67


Table 2: Hatching performance in second season of green turtles in laboratory of Chendor (CH), Pahang

Temperature (°C)	CH4(Mother - 4)		CH5 (Mother - 5)		CH6 (Mother - 6)		% Average
	eggs incubated	% Hatching	eggs incubated	% Hatching	eggs incubated	% Hatching	
24°C	10	0	10	0	10	0	0.00
25°C	10	90	10	90	10	100	93.33
26°C	10	100	10	100	10	100	100.00
27°C	10	90	10	80	10	100	90.00
28°C	10	100	10	100	10	100	100.00
29°C	10	100	10	90	10	90	93.33
30°C	10	100	10	100	10	100	100.00
31°C	10	100	10	100	10	100	100.00
32°C	10	100	10	100	10	90	96.67
33°C	10	90	10	100	10	80	90.00
34°C	10	0	10	100	0	80	60.00

Table 3: Sex identification using histological protocols of green turtle's hatchlings at Chendor, at the various temperatures in 1998.

Temperature °C	CH-1			CH-2			CH-3			Total			%		
	M	F	I	M	F	I	M	F	I	M	F	I	M	F	I
27.2°C	0	0	0	9	0	1	7	0	3	16	0	4	80	0	20
27.7°C	0	0	0	9	0	0	7	0	3	16	0	3	84	0	16
28.2°C	0	0	0	10	0	0	6	3	0	16	3	0	84	16	0
28.7°C	2	3	0	6	4	0	6	4	0	14	11	0	56	44	0
29.0°C	2	3	0	10	0	0	6	4	0	18	7	0	72	28	0
29.2°C	1	4	0	7	3	0	5	5	0	13	12	0	52	48	0
29.5°C	0	0	0	5	3	0	2	8	0	7	11	0	39	61	0
29.7°C	0	0	0	0	0	0	3	7	0	3	7	0	30	70	0
30°C	0	6	0	4	6	0	3	4	0	7	16	0	30	70	0
31°C	0	6	0	1	9	0	0	10	0	1	25	0	4	96	0
32°C	0	7	0	0	10	0	0	10	0	0	20	0	0	100	0
33°C	0	5	0	0	10	0	0	6	0	0	21	0	0	100	0

Figure 3: Percentage of sex determination at various temperatures in Chendor, Pahang



M = male; F = female; I = Indeterminate

The above graph indicates that temperature values between 27.2 °C to 29.0°C produce more male turtle

## DISCUSSION

The result on incubation of the first season (May - June) showed that the average hatching percentage obtained higher than 90% at the temperatures 27°C to 32°C (Table 1). Whereas in the second season (August - September) the average hatching percentage obtained higher than 90% at the temperature 25°C to 33°C (Table 2). This occurrence might be due to handling technique of turtle eggs during the transportation from the rookery to laboratory. For instance the turtle eggs should be incubated immediately in an incubator after arriving at the laboratory. Whereby proper handling would maintain the quality of turtle eggs for hatching. Miller and Limpus (1981) reported that temperature influences the duration of incubation and the rate of the development. Low temperatures increase the duration of incubation and slow the rate of development, whereas higher temperatures decrease the duration of incubation and increase the rate of development. In conservation efforts, improper handling of eggs during movement to hatcheries may increase mortality (Limpus et al., 1979; Blank and Sawyer, 1981).

The temperatures between 31°C to 33°C produce 96% to 100% of female hatchlings (Table 3). Thus the particular temperatures could be considered for female hatchlings at hatchery incubation. Percentage of male hatchlings obtained 80% to 84% at the temperatures ranging from 27.2 °C to 28.2°C. Whereas at the temperature 27.2 °C and 27.7°C, 20 % and 16 % respectively are indeterminate due to poor technique on preparation of specimen slides. This result can predict that the temperature between 27.2 °C and 27.7°C would produce the higher percentage of male green turtle hatchlings. Under natal beach conditions, the eggs incubate at temperatures between 24°C to 33°C (Cardwell, 1959; Bustard, 1972; Ewert, 1979; Limpus et al., 1983). *Chelonia mydas* the incubation period lasted 94 days at 23°C -24°C (Ackerman and Prange, 1972) and 47 - 49 days at 32°C (Bustard and Greenham, 1968). Within the limit of embryonic tolerance, cooler temperature produce male hatchlings at 26°C and 28°C for *Caretta caretta* and 26°C for *Chelonia mydas*.

## Conclusion

The average hatching performance obtained higher than 80 % at the temperature between 25°C to 33°C. The average temperatures between 27.2°C to 29.2°C produce more male green turtle hatchlings. Whereas more than 80% male hatchlings are produced at the temperature 27.2°C to 28.2°C. Female hatchlings are dominant at the temperature value above 29.5°C. Temperatures between 31.0°C to 33.0°C produce

more than 96% female hatchlings. Temperatures which are below 25°C and above 33°C are not suitable for eggs incubation of green turtle at Chendor rookery.

## ACKNOWLEDGEMENTS

Great thanks to the Director General of Fisheries Malaysia, for given the permission to conduct the study. Many thanks to Chief of MFRDMD and Deputy Chief of MFRDMD for their supporting and encouraging on implementation of this study. Last but not least the authors wish to express the appreciation to the staffs of MFRDMD Kuala Terengganu and Fisheries Department of Pahang for their assistances in conducting the study.

## REFERENCES

- Ackerman, R. A. 1977.** The respiratory gas exchange of sea turtle nests (*Chelonia, Caretta*). *Respir. Physiol.* **31**, 19 - 38.
- Ackerman, R. A. and Prange, H. D. 1972.** Oxygen diffusion across a sea turtle (*Chelonia, mydas*) egg shell. *Comp. Biochem. Physiol.* **43A**, 905-909.
- Blanck, C. E. and Sawyer, R. H. 1981.** Hatchery practices in relation to early embryology of the longgerhead sea turtle, *Caretta caretta* (Linne). *J. exp. mar. Biol. Ecol.* **49**, 163-177.
- Bustard, H. R. (1972).** *Sea turtles: Natural History and Conservation*. Collins, Sydney.
- Bustard, H. R. and Greenham, P. 1968.** Physical and chemical factors affecting hatching in the green sea turtle, *Chelonia mydas* (L). *Ecology* **49**, 269 - 276.
- Caldwell, D. K. 1959.** The longgerhead turtle of Cape Romain, South Carolina. *Bull. Florida State Mus.* **4**, 317-348.
- Dutton, P.H., Whitmore, C.P. & Mrosovsky, N. 1985.** Masculinization of leatherback turtle *Dermochelys coriacea* hantchlings from eggs incubated in styrofoam boxes. *Biol. Conserve.*, **31**, 249-64.
- Ewert, M. A. 1979.** The embryo and its egg: development and natural history. In *Turtle: Perspectives and Research*(M. Harless and H. Morlock, eds). J.Wiley and Sons, New York, pp. 333-413.
- Hendrickson, J. R. 1980.** The ecological strategies of sea turtle. *Amer. Zool.* **20**, 597-608. Kraemer, J. E. and bell, R. (1980). Rain-induced mortality of eggs and hatchlings of loggerhead sea turtle (*Caretta caretta*) on the Georgia coast. *Herpetologica* **36**, 72-77.

- Limpus, C. J. Baker, V. and Miller, J. D. 1979.** Movement induced mortality of longgerhead eggs. *Herpetologica* 35, 335-338.
- Limpus, C. J., Reed, P., and Miller, J. D. 1983c.** Islands and turtle. The influence of choice of nesting beach on sex ratio. In *Proceedings: Inaugural Great Barrier Reef Conference* (J.T. Baker, R. M. Carter, P.W. sammarco, and K. P. Stark, eds.) James Cook University Press, Townsville, Aust., pp. 397-402.
- Miller, J. D. and Limpus, C.J. 1981.** Incubation period and sexual differentiation in the green turtle *Chelonia mydas* L. In *Proceedings of the Melbourne Herpetological Symposium* (C.B. Banks and A. Martin, eds.) The Royal Melbourne Zoological Gardens, Melbourne, pp. 66-73.
- Morreale, S. J., Ruiz, G. J. , Spotila, J. R. and Standora, E. a. 1982.** Tempurature dependent sex determination: current practices threaten conservation of sea turtle. *Science, N. Y.* 216, 1245-1247.
- Mrosovsky, N. Dutton, P.H. & Whitmore, C.P. 1984.** Sex ratio of two species of sea turtle nesting in Surinam. *Can. J. Zool.*, 62, 227-
- Parmenter, C. J. 1980.** Environmental factors in turtle farming. In *Management of Turtle Resources: Research Monograph I*. James Cook University of North Queensland. Australia, 23-31.
- Ragotzkie, U. 1976.** Mortality of longgerhead turtle eggs from excessive rainfall. *Ecology* 40, 303-305.
- Reed, P. C. 1980.** "The Sex Ratio of Hatchling Longgerhead Turtle – the Progeny of Two Nesting Adult Female". Honours thesis, James Cook University, Queensland, Australia.
- Yntema, C. and Mrosovsky, N. 1979.** Incubation temperature and sex ratio in hatchling longgerhead turtle: a preliminary report. *Mar. Turtle Newslet.* 11, 9-10.
- Yntema, C. and Mrosovsky, N. 1980. Sexual differentiation in hatchling longgerheads (*Caretta caretta*) incubated at different controlled temperatures. *Herpetologica* 36, 33-36.