

Title	On The N-Fractional Calculus of Some Products of Some Power Functions(Sakaguchi Functions in Univalent Function Theory and Its Applications)
Author(s)	Nishimoto, Katsuyuki
Citation	数理解析研究所講究録 (2006), 1470: 11-17
Issue Date	2006-02
URL	http://hdl.handle.net/2433/48102
Right	
Type	Departmental Bulletin Paper
Textversion	publisher

On The N-Fractional Calculus of Some Products of Some Power Functions

Katsuyuki Nishimoto

Abstract

In this article, N-fractional calculus of products of power functions

$$((z-c)^\alpha \cdot (z-c)^\beta)_\nu, ((z-c)^\beta \cdot (z-c)^\alpha)_\nu, \text{ and } ((z-c)^{\alpha+\beta})_\nu$$

are discussed again.

§ 0. Introduction (Definition of Fractional Calculus)

(I) Definition. (by K. Nishimoto) ([1] Vol. 1)

Let $D = \{D_-, D_+\}$, $C = \{C_-, C_+\}$,

C_- be a curve along the cut joining two points z and $-\infty + i\text{Im}(z)$,

C_+ be a curve along the cut joining two points z and $\infty + i\text{Im}(z)$,

D_- be a domain surrounded by C_- , D_+ be a domain surrounded by C_+ .

(Here D contains the points over the curve C).

Moreover, let $f = f(z)$ be a regular function in $D(z \in D)$,

$$f_\nu(z) = (f)_\nu = {}_c(f)_\nu = \frac{\Gamma(\nu+1)}{2\pi i} \int_C \frac{f(\xi)}{(\xi-z)^{\nu+1}} d\xi \quad (\nu \notin \mathbf{Z}), \quad (1)$$

$$(f)_{-m} = \lim_{\nu \rightarrow -m} (f)_\nu \quad (m \in \mathbf{Z}^+), \quad (2)$$

where $-\pi \leq \arg(\xi-z) \leq \pi$ for C_- , $0 \leq \arg(\xi-z) \leq 2\pi$ for C_+ ,

$\xi \neq z$, $z \in C$, $\nu \in \mathbf{R}$, Γ ; Gamma function,

then $(f)_\nu$ is the fractional differintegration of arbitrary order ν (derivatives of order ν for $\nu > 0$, and integrals of order $-\nu$ for $\nu < 0$), with respect to z , of the function f , if $|(f)_\nu| < \infty$.

(II) On the fractional calculus operator N^ν [3]

Theorem A. Let fractional calculus operator (Nishimoto's Operator) N^ν be

$$N^\nu = \left(\frac{\Gamma(\nu+1)}{2\pi i} \int_C \frac{d\xi}{(\xi-z)^{\nu+1}} \right) \quad (\nu \notin \mathbf{Z}), \quad [\text{Refer to (1)}] \quad (3)$$

with

$$N^{-m} = \lim_{\nu \rightarrow -m} N^\nu \quad (m \in \mathbf{Z}^+), \quad (4)$$

and define the binary operation \circ as

$$N^\beta \circ N^\alpha f = N^\beta N^\alpha f = N^\beta(N^\alpha f) \quad (\alpha, \beta \in \mathbf{R}), \quad (5)$$

then the set

$$\{N^\nu\} = \{N^\nu \mid \nu \in \mathbf{R}\} \quad (6)$$

is an Abelian product group (having continuous index ν) which has the inverse transform operator $(N^\nu)^{-1} = N^{-\nu}$ to the fractional calculus operator N^ν , for the function f such that $f \in F = \{f; 0 \neq |f_\nu| < \infty, \nu \in \mathbf{R}\}$, where $f = f(z)$ and $z \in \mathbf{C}$. (vis. $-\infty < \nu < \infty$).

(For our convenience, we call $N^\beta \circ N^\alpha$ as product of N^β and N^α .)

Theorem B. " F.O.G. $\{N^\nu\}$ " is an " Action product group which has continuous index ν " for the set of F . (F.O.G. ; Fractional calculus operator group)

Theorem C. Let

$$S := \{\pm N^\nu\} \cup \{0\} = \{N^\nu\} \cup \{-N^\nu\} \cup \{0\} \quad (\nu \in \mathbf{R}). \quad (7)$$

Then the set S is a commutative ring for the function $f \in F$, when the identity

$$N^\alpha + N^\beta = N^\gamma \quad (N^\alpha, N^\beta, N^\gamma \in S) \quad (8)$$

holds. [5]

(III) Lemma. We have [1]

$$(i) \quad ((z-c)^\beta)_\alpha = e^{-i\pi\alpha} \frac{\Gamma(\alpha-\beta)}{\Gamma(-\beta)} (z-c)^{\beta-\alpha} \quad \left(\left| \frac{\Gamma(\alpha-\beta)}{\Gamma(-\beta)} \right| < \infty \right),$$

$$(ii) \quad (\log(z-c))_\alpha = -e^{-i\pi\alpha} \Gamma(\alpha) (z-c)^{-\alpha} \quad (|\Gamma(\alpha)| < \infty),$$

$$(iii) \quad ((z-c)^{-\alpha})_{-\alpha} = -e^{i\pi\alpha} \frac{1}{\Gamma(\alpha)} \log(z-c) \quad (|\Gamma(\alpha)| < \infty),$$

where $z-c \neq 0$ in (i), and $z-c \neq 0, 1$ in (ii) and (iii). (Γ ; Gamma function),

$$(iv) \quad (u \cdot v)_\alpha := \sum_{k=0}^{\infty} \frac{\Gamma(\alpha+1)}{k! \Gamma(\alpha+1-k)} u_{\alpha-k} v_k \quad \left(\begin{array}{l} u = u(z), \\ v = v(z) \end{array} \right).$$

§ 1. N-fractional calculus of products of some power functions

In the following $\alpha, \beta, \gamma \in \mathbf{R}$, for our convenience.

Theorem 1. *Let*

$$P = P(\alpha, \beta, \gamma) := \frac{\sin \pi \alpha \cdot \sin \pi(\gamma - \alpha - \beta)}{\sin \pi(\alpha + \beta) \cdot \sin \pi(\gamma - \alpha)} \quad (|P(\alpha, \beta, \gamma)| = M < \infty) \quad (1)$$

and

$$Q = Q(\alpha, \beta, \gamma) := P(\beta, \alpha, \gamma) \quad (|P(\beta, \alpha, \gamma)| = M < \infty) \quad (2)$$

When $\alpha, \beta, \gamma \notin \mathbf{Z}_0^+$, we have ;

$$(i) \quad ((z-c)^\alpha \cdot (z-c)^\beta)_\gamma = e^{-i\pi\gamma} P(\alpha, \beta, \gamma) \frac{\Gamma(\gamma - \alpha - \beta)}{\Gamma(-\alpha - \beta)} (z-c)^{\alpha+\beta-\gamma}, \quad (3)$$

$$(\operatorname{Re}(\alpha + \beta + 1) > 0, (1 + \alpha - \gamma) \notin \mathbf{Z}_0^-),$$

$$(ii) \quad ((z-c)^\beta \cdot (z-c)^\alpha)_\gamma = e^{-i\pi\gamma} Q(\alpha, \beta, \gamma) \frac{\Gamma(\gamma - \alpha - \beta)}{\Gamma(-\alpha - \beta)} (z-c)^{\alpha+\beta-\gamma}, \quad (4)$$

$$(\operatorname{Re}(\alpha + \beta + 1) > 0, (1 + \beta - \gamma) \notin \mathbf{Z}_0^-)$$

$$(iii) \quad ((z-c)^{\alpha+\beta})_\gamma = e^{-i\pi\gamma} \frac{\Gamma(\gamma - \alpha - \beta)}{\Gamma(-\alpha - \beta)} (z-c)^{\alpha+\beta-\gamma}, \quad (5)$$

where

$$z-c \neq 0, \quad \left| \frac{\Gamma(\gamma - \alpha - \beta)}{\Gamma(-\alpha - \beta)} \right| < \infty.$$

Proof of (i). We have

$$((z-c)^\alpha \cdot (z-c)^\beta)_\gamma = \sum_{k=0}^{\infty} \frac{\Gamma(\gamma+1)}{k! \Gamma(\gamma+1-k)} ((z-c)^\alpha)_{\gamma-k} ((z-c)^\beta)_k \quad (6)$$

by Lemma (iv).

Next we have

$$((z-c)^\alpha)_{\gamma-k} = e^{-i\pi(\gamma-k)} \frac{\Gamma(\gamma-k-\alpha)}{\Gamma(-\alpha)} (z-c)^{\alpha-\gamma+k} \quad \left(\left| \frac{\Gamma(\gamma-k-\alpha)}{\Gamma(-\alpha)} \right| < \infty \right) \quad (7)$$

and

$$((z-c)^\beta)_k = e^{-i\pi k} \frac{\Gamma(k-\beta)}{\Gamma(-\beta)} (z-c)^{\beta-k} \quad (8)$$

by Lemma (i), respectively.

Substitute (7) and (8) into (6), we have then

$$((z-c)^\alpha \cdot (z-c)^\beta)_\gamma = e^{-i\pi\gamma} \sum_{k=0}^{\infty} \frac{\Gamma(\gamma+1)\Gamma(\gamma-\alpha-k)\Gamma(k-\beta)}{k!\Gamma(\gamma+1-k)\Gamma(-\alpha)\Gamma(-\beta)} (z-c)^{\alpha+\beta-\gamma} \quad (9)$$

$$= e^{-i\pi\gamma} \frac{\Gamma(\gamma-\alpha)}{\Gamma(-\alpha)} (z-c)^{\alpha+\beta-\gamma} \sum_{k=0}^{\infty} \frac{[-\beta]_k [-\gamma]_k}{k! [1+\alpha-\gamma]_k} \quad (10)$$

using the relationship

$$\Gamma(\lambda+1-k) = (-1)^{-k} \frac{\Gamma(\lambda+1)\Gamma(-\lambda)}{\Gamma(k-\lambda)}, \quad (11)$$

where

$$[\lambda]_k = \lambda(\lambda+1)\cdots(\lambda+k-1) = \Gamma(\lambda+k)/\Gamma(\lambda), \text{ with } [\lambda]_0 = 1. \\ (\text{notation of Pochhammer}).$$

Therefore, applying the following relationships

$$\sum_{k=0}^{\infty} \frac{[a]_k [b]_k}{k! [c]_k} = {}_2F_1(a, b; c; 1) = \frac{\Gamma(c)\Gamma(c-a-b)}{\Gamma(c-a)\Gamma(c-b)} \left(\begin{array}{l} \operatorname{Re}(c-a-b) > 0, \\ c \notin \mathbf{Z}_0^- \end{array} \right) \quad (12)$$

and

$$\Gamma(\lambda)\Gamma(1-\lambda) = \frac{\pi}{\sin \pi\lambda} \quad (\lambda \notin \mathbf{Z}) \quad (13)$$

to (10), we obtain

$$((z-c)^\alpha \cdot (z-c)^\beta)_\gamma = e^{-i\pi\gamma} \frac{\Gamma(\gamma-\alpha)}{\Gamma(-\alpha)} (z-c)^{\alpha+\beta-\gamma} {}_2F_1(-\beta, -\gamma; 1+\alpha-\gamma; 1) \quad (14)$$

$$= e^{-i\pi\gamma} \frac{\Gamma(\gamma-\alpha)\Gamma(1+\alpha+\beta)\Gamma(1+\alpha-\gamma)}{\Gamma(-\alpha)\Gamma(1+\alpha)\Gamma(1+\alpha+\beta-\gamma)} (z-c)^{\alpha+\beta-\gamma} \quad (15) \\ \left(\begin{array}{l} \operatorname{Re}(\alpha+\beta+1) > 0, \\ (1+\alpha-\gamma) \notin \mathbf{Z}_0^- \end{array} \right)$$

$$= e^{-i\pi\gamma} \frac{\sin \pi\alpha \cdot \sin \pi(\gamma-\alpha-\beta)}{\sin \pi(\alpha+\beta) \cdot \sin \pi(\gamma-\alpha)} \cdot \frac{\Gamma(\gamma-\alpha-\beta)}{\Gamma(-\alpha-\beta)} (z-c)^{\alpha+\beta-\gamma} \quad (16)$$

We have then (3) from (16) using the notation (1), under the conditions.

Proof of (ii). In the same way as the proof of (i) we obtain (4) using (2) instead of (1), under the conditions. (Simply, changing α and β in (3) we obtain (4), using (2).)

Proof of (iii). We have

$$\begin{aligned} ((z-c)^{\alpha+\beta})_y &= e^{-i\pi y} \frac{\Gamma(\gamma-\alpha-\beta)}{\Gamma(-\alpha-\beta)} (z-c)^{\alpha+\beta-\gamma} & (17) \\ &\left(\left| \frac{\Gamma(\gamma-\alpha-\beta)}{\Gamma(-\alpha-\beta)} \right| < \infty \right) \end{aligned}$$

directly by Lemma (i).

Corollary 1.

When $\alpha, \beta, \gamma \notin \mathbf{Z}_0^+$, we have ;

$$(i) \quad (z^\alpha \cdot z^\beta)_y = e^{-i\pi y} P(\alpha, \beta, \gamma) \frac{\Gamma(\gamma-\alpha-\beta)}{\Gamma(-\alpha-\beta)} z^{\alpha+\beta-\gamma}, \quad (18)$$

$$(\operatorname{Re}(\alpha+\beta+1) > 0, (1+\alpha-\gamma) \notin \mathbf{Z}_0^-),$$

$$(ii) \quad (z^\beta \cdot z^\alpha)_y = e^{-i\pi y} Q(\alpha, \beta, \gamma) \frac{\Gamma(\gamma-\alpha-\beta)}{\Gamma(-\alpha-\beta)} z^{\alpha+\beta-\gamma}, \quad (19)$$

$$(\operatorname{Re}(\alpha+\beta+1) > 0, (1+\beta-\gamma) \notin \mathbf{Z}_0^-)$$

$$(iii) \quad (z^{\alpha+\beta})_y = e^{-i\pi y} \frac{\Gamma(\gamma-\alpha-\beta)}{\Gamma(-\alpha-\beta)} z^{\alpha+\beta-\gamma}, \quad (20)$$

where

$$\left| \frac{\Gamma(\gamma-\alpha-\beta)}{\Gamma(-\alpha-\beta)} \right| < \infty,$$

and P and Q are the ones shown by (1) and (2) respectively.

Proof. Set $c=0$ in Theorem 1.

Theorem 2. When $\alpha, \beta, \gamma \notin \mathbf{Z}_0^+$, we have ;

$$(i) \quad ((z-c)^\alpha \cdot (z-c)^\beta)_y = P(\alpha, \beta, \gamma) ((z-c)^{\alpha+\beta})_y, \quad (21)$$

$$(\operatorname{Re}(\alpha+\beta+1) > 0, (1+\alpha-\gamma) \notin \mathbf{Z}_0^-),$$

$$(ii) \quad ((z-c)^\beta \cdot (z-c)^\alpha)_y = Q(\alpha, \beta, \gamma) ((z-c)^{\alpha+\beta})_y, \quad (22)$$

$$(\operatorname{Re}(\alpha+\beta+1) > 0, (1+\beta-\gamma) \notin \mathbf{Z}_0^-)$$

$$(iii) \quad \frac{1}{P} ((z-c)^\alpha \cdot (z-c)^\beta)_y = \frac{1}{Q} ((z-c)^\beta \cdot (z-c)^\alpha)_y = ((z-c)^{\alpha+\beta})_y, \quad (23)$$

$$(\operatorname{Re}(\alpha+\beta+1) > 0, (1+\alpha-\gamma) \notin \mathbf{Z}_0^-, (1+\beta-\gamma) \notin \mathbf{Z}_0^-, PQ \neq 0)$$

where

$$z - c \neq 0, \quad \left| \frac{\Gamma(\gamma - \alpha - \beta)}{\Gamma(-\alpha - \beta)} \right| < \infty,$$

and P and Q are the ones shown by (1) and (2) respectively.

Proof. It is clear by Theorem 1.

Corollary 2. When $\alpha, \beta, \gamma \notin \mathbf{Z}_0^+$, we have ;

$$(i) \quad (z^\alpha \cdot z^\beta)_\gamma = P(\alpha, \beta, \gamma) (z^{\alpha+\beta})_\gamma, \quad (24)$$

$$(\operatorname{Re}(\alpha + \beta + 1) > 0, (1 + \alpha - \gamma) \notin \mathbf{Z}_0^-),$$

$$(ii) \quad (z^\beta \cdot z^\alpha)_\gamma = Q(\alpha, \beta, \gamma) (z^{\alpha+\beta})_\gamma, \quad (25)$$

$$(\operatorname{Re}(\alpha + \beta + 1) > 0, (1 + \beta - \gamma) \notin \mathbf{Z}_0^-)$$

$$(iii) \quad \frac{1}{P} (z^\alpha \cdot z^\beta)_\gamma = \frac{1}{Q} (z^\beta \cdot z^\alpha)_\gamma = (z^{\alpha+\beta})_\gamma, \quad (26)$$

$$(\operatorname{Re}(\alpha + \beta + 1) > 0, (1 + \alpha - \gamma) \notin \mathbf{Z}_0^-, (1 + \beta - \gamma) \notin \mathbf{Z}_0^-, PQ \neq 0)$$

where

$$\left| \frac{\Gamma(\gamma - \alpha - \beta)}{\Gamma(-\alpha - \beta)} \right| < \infty,$$

and P and Q are the ones shown by (1) and (2) respectively.

Proof. Set $c = 0$ in Theorem 2.

Corollary 3.

(i) When $\alpha, \beta, \gamma \notin \mathbf{Z}_0^+$ and

$$P(\alpha, \beta, \gamma) = Q(\alpha, \beta, \gamma) = 1, \quad (27)$$

we have

$$((z-c)^\alpha \cdot (z-c)^\beta)_\gamma = ((z-c)^\beta \cdot (z-c)^\alpha)_\gamma = ((z-c)^{\alpha+\beta})_\gamma. \quad (28)$$

$$(\operatorname{Re}(\alpha + \beta + 1) > 0, (1 + \alpha - \gamma) \notin \mathbf{Z}_0^-, (1 + \beta - \gamma) \notin \mathbf{Z}_0^-)$$

(ii) When $\gamma = m \in \mathbf{Z}_0^+$ we have

$$((z-c)^\alpha \cdot (z-c)^\beta)_m = ((z-c)^\beta \cdot (z-c)^\alpha)_m = ((z-c)^{\alpha+\beta})_m. \quad (29)$$

Proof of (i). We have (28) from Theorem 2 (23), clearly.

Proof of (ii). When $\gamma = m \in \mathbf{Z}_0^+$ we have (29) from Theorem 2 (23), since

$$P(\alpha, \beta, m) = Q(\alpha, \beta, m) = 1. \quad (30)$$

References

- [1] K. Nishimoto ; Fractional Calculus, Vol. 1 (1984), Vol. 2 (1987), Vol. 3 (1989), Vol. 4 (1991), Vol. 5, (1996), Descartes Press, Koriyama, Japan.
- [2] K. Nishimoto ; An Essence of Nishimoto's Fractional Calculus (Calculus of the 21st Century); Integrals and Differentiations of Arbitrary Order (1991), Descartes Press, Koriyama, Japan.
- [3] K. Nishimoto ; On Nishimoto's fractional calculus operator N^ν (On an action group), J. Frac. Calc. Vol. 4, Nov. (1993), 1 - 11.
- [4] K. Nishimoto ; Unification of the integrals and derivatives (A serendipity in fractional calculus), J. Frac. Calc. Vol. 6, Nov. (1994), 1 - 14.
- [5] K. Nishimoto ; Ring and Field Produced from The Set of N- Fractional Calculus Operator, J. Frac Calc. Vol. 24, Nov. (2003), 29 - 36.
- [6] K. Nishimoto, Ding- Kuo Chyan, Shy- Der Lin and Shih- Tong Tu; On some infinite sums derived by N- fractional calculus, J. Frac. Calc. Vol.20 (2001), 91 - 97.
- [7] Pin Yu Wang, Tsu- Chen Wu and Shih- Tong Tu; Some Infinite Sums via N- fractional calculus, J. Frac. Calc. Vol.21, May (2002), 71 -77.
- [8] Shy- Der Lin, Shih- Tong Tu , Tsai- Ming Hsieh and H.M. Srivastava ; Some Finite and Infinite Sums Associated with the Digamma and Related Functions, J. Frac.Calc.Vol.22, Nov. (2002), 103- 114.
- [9] K. Nishimoto ; N- Fractional Calculus of the Power and Logarithmic Functions and Some Identities (Continue), J. Frac. Calc. Vol.22, Nov. (2002), 59 - 65.
- [10] K.B. Oldham and J. Spanier ; The Fractional Calculus (1974), Academic Press.
- [11] A.C. McBride ; Fractional Calculus and Integral Transforms of Generalized Functions, Research Notes, Vol. 31, (1979), Pitman.
- [12] S. G. Samko, A.A. Kilbas and O.I. Marichev ; Fractional Integrals and Derivatives, and Some Their Applications (1987), Nauka, USSR.
- [13] K.S. Miller and B. Ross ; An Introduction to The Fractional Calculus and Fractional Differential Equations, John Wiley & Sons, (1993).
- [14] V. Kiryakova ; Generalized fractional calculus and applications, Pitman Research Notes, No. 301, (1994), Longman.
- [15] Igor Podlubny ; Fractional Differential Equations (1999), Academic Press.
- [16] R. Hilfer (Ed.) ; Applications of Fractional Calculus in Physics, (2000), World Scientific, Singapore, New Jersey, London, Hong Kong.

Katsuyuki Nishimoto
 Institute of Applied Mathematics
 Descartes Press Co.
 2 - 13 - 10 Kaguike, Koriyama
 963 - 8833 Japan