

## **Thermal history of the NE Japan frontal arc since the Late Miocene inferred from vitrinite reflectance**

Atsushi Yamaji

Institute of Geology and Paleontology, Tohoku University, Aobayama, Sendai 980, Japan

### **ABSTRACT**

Paleogeothermal gradients are estimated from the vitrinite reflectance-depth ( $Ro$ - $z$ ) relations of Middle to Late Miocene sediments in the Tanakura and Hirono areas, Northeast Japan. Both areas have been situated in the frontal arc since 15 Ma and have been free from magmatism. Thus, the sediments would have been undergone a limited amount of local, thermal disturbances, and are suitable to study regional-scale paleogeothermal conditions.

Late Miocene to Early Pliocene uplift resulted in an angular unconformity between the Miocene and Pliocene rocks and caused cooling of the Miocene sediments that would have retarded and effectively stopped coalification. The  $Ro$ - $z$  relations of the sediments therefore reveal the paleogeothermal gradient between deposition and unroofing. The estimated paleogeothermal gradient is  $64 \pm 11^\circ \text{C km}^{-1}$  at Tanakura and  $45 \pm 10^\circ \text{C km}^{-1}$  at Hirono. The present gradients are respectively  $30^\circ$  and  $18^\circ \text{C km}^{-1}$ , indicating that subsurface temperature was higher in the Late Miocene than the present under the frontal arc.

Cooling from the Pliocene through Quaternary appears to have been of regional scale across the arc as it is also suggested by the geochemical signature of volcanic rocks erupted at the volcanic front [Ban et al., 1992]. Westward retreat of the volcanic front [Ohguchi et al., 1989] is also in accord with the regional cooling.

### **INTRODUCTION**

Geothermics is an important clue to island arc dynamics and has been investigated mainly on the basis of heat flow measurements [e.g., Uyeda, 1972]. However, such geophysical measurements show only a snapshot: island arcs have their own history. The NE Japan arc, as an example, experienced a sequence of tectonic events in the Late Cenozoic such as the spreading and subsequent subduction-initiation in the back arc [Jolivet and Tamaki, 1992; Nakamura, 1983; Kobayashi, 1983]. Geothermal regime must have varied together with the tectonics. The secular change of arc magmatism suggests such variations under the arc. Volcanic front has migrated back and forth across the NE Japan arc during the Cenozoic [Ohguchi et al., 1989; Ohki et al., 1993]. Sato et al. [in preparation] document the variation of production rate of volcanic materials in the arc since the mid Tertiary. They found that the rate has changed up to two orders of magnitude. These observations suggest that the

estimation of geothermal history provides an important clue to island arc dynamics.

The analysis of the thermal history of NE Japan has been undertaken recently by the petrology of volcanic rocks [Ban et al., 1992; Yoshida 1992] and by the studies of organic maturation [Suzuki, 1989]. Ban et al. and Yoshida calculate the silica-normalized abundance of LIL elements in Cenozoic volcanics and show tentative results of the spatio-temporal variations of the depth of an isotherm that determines the element abundances. Suzuki measured vitrinite reflectance, the most widely used parameter of organic diagenesis, of coaly particles in Neogene sediments and suggested that the gross pattern of the distribution of paleogeothermal gradient in the Middle Miocene did not differ from that of the present. However, a significant part of his data was obtained in the volcanic arc of NE Japan in the exploration of metallic ores that




Figure 1. Map showing the location of the Hirono and Tanakura areas and the position of volcanic front at and after 15 Ma [Ohguchi et al., 1989]. (a): volcanic front at 16–15 Ma, (b): in the Late Miocene (8–5 Ma), and (c): 15–8 Ma and 0–5 Ma.

are often formed at thermally abnormal loci. The maximum temperature to which coal was exposed exerts a strong control on vitrinite reflectance [Hood et al., 1975; Barker and Pawlewicz, 1986]. Therefore Suzuki's estimation is possibly biased by local anomalies.

In this paper an attempt is made to estimate ancient geothermal gradients in the NE Japan arc from vitrinite reflectance. The Tanakura and Hirono areas are selected around the Abukuma mountains that have been situated in the frontal arc since 15 Ma (Fig. 1). The Miocene sequence in the studied areas was uplifted in Early Pliocene that resulted in an angular unconformity. Due to the associated unroofing and cooling of the Miocene strata, coalification would have been stopped so that an analysis of them enables us to estimate geothermal gradient in the Miocene.

## METHOD

The monotonic increase of the reflectance of vitrinite, a constituent of coals, through diagenesis has permitted quantitative reconstructions of the burial and temperature

histories of stratigraphic sequences. There are a number of methods proposed to correlate vitrinite reflectance  $R_o$  with the degree of thermal stress undergone by strata containing the vitrinite. The present study uses Middleton's formulation [Middleton, 1982a, b; Middleton and Schmidt, 1982]:

$$R_o^a = B \int \exp[bT(t)] dt \quad (1)$$

because this is simple but approximate to recent sophisticated formulations [Morrow and Issler, 1993]. In this equation,  $t$  is time since deposition and  $T(t)$  is the temperature at which the coal was exposed. Middleton determines the coefficients,

$$a = 5.5, b = 0.069^\circ \text{C}^{-1} \text{ and } B = 2.8 \times 10^{-5} \text{ Myr}^{-1}. \quad (2)$$

$T$  is approximated by the equation

$$T = T_o + gz \quad (3)$$

where  $z$  is the burial depth,  $T_o$  and  $g$  are respectively the surface temperature and geothermal gradient. From equations (1) and (3) we have

$$\log R_o^a = \frac{1}{a} \log \left[ \int_0^t \exp(bgz) dt \right] + \frac{bT_o + \log B}{a}.$$

If the geothermal gradient and depth do not significantly change, this reduces to

$$\log R_o = (b/a)gz + \text{const.} \quad (4)$$

Since coalification is much more sensitive to temperature than to time,  $R_o$  is determined mostly by the maximum temperature which the coal experienced [Hood et al., 1975]. Accordingly the reflectance records the maximum burial temperature [Yamaji, 1986]. The linear correlation of  $\log R_o$  with depth  $z$  represented by equation (4) is commonly observed in sedimentary basins [Dow, 1977].

Cooling due to uplift and erosion of overlying sediments effectively stops coalification. In this case, the reflectance gradient

$$\Gamma = (\log R_o) / z$$

is related to the thermal gradient before the uplift through the equation

$$g = (a/b)\Gamma. \quad (5)$$

This is the upper limit for the bed temperature between the deposition and uplift because of the irreversibility of coalification.

## GEOLOGICAL BACKGROUND OF STUDIED AREAS AND RESULTS

### Tanakura area

Coal particles (> #200 mesh) were collected from cuttings of a bore-hole that penetrates the Akasaka and Kubota formations in the Tanakura area. The bore-hole is at proximity to a syncline so that dips of strata are very gentle. The formations consist of Middle to early Late Miocene paralic sediments that cover Mesozoic metamorphic rocks (Fig. 2). The formations yield calcareous nanofossils from CN5a to CN8




Figure 2. Geologic map of the Tanakura area [Otsuki, 1975]. Circle: bore-hole site.

zones [Amano and Takahashi, 1986] which are dated at 14–8 Ma [Oda, 1986].

The formations were uplifted at the Miocene-Pliocene boundary and the non-marine Nikogi formation unconformably covers the Miocene sequence [Otsuki, 1975]. The formation is correlated with the Gauss normal epoch (~3 Ma) [Tohoku Agricultural Administration, 1986].




Figure 4. Geologic map of the Hirono area [Taketani et al., 1986].





Figure 3. Vitrinite reflectance versus depth for the Tanakura (a) and Hirono (b) samples.

Figure 3a shows the linear regression between  $\log R_o$  and depth with a reflectance gradient of  $\Gamma = 0.81 \pm 0.14 \text{ km}^{-1}$  where accuracy is represented by one sigma. Using the coefficients (2) and the formula (5), the  $\Gamma$  is converted to the thermal gradient at  $64 \pm 11^\circ \text{ Ckm}^{-1}$ . The present geothermal gradient is  $30^\circ \text{ Ckm}^{-1}$  [Tohoku Agricultural Administration, 1986], significantly lower than the paleogeothermal gradient.

#### Hirono area

The Lower Miocene Yunagaya group is overlain by the Pliocene Hirono Formation with a sharp angular unconformity in this area [Sugai et al., 1957]. The lowermost part of the group consists of a thin volcanic breccia that yields K-Ar and fission track ages of 20.9 and 23.4 Ma respectively [Kimura, 1988]. The breccia is immediately overlain by shallow marine, non-volcanic sediments. Planktonic fossils indicate that the marine



sediments were deposited from ~18 to 16 Ma [Koizumi, 1986;

Figure 5. Maximum paleogeothermal gradients in the Tanakura (a) and Hirono (b) areas estimated from vitrinite reflectance-depth relations. Indicated is the period in which the maximum gradient must have occurred and it is not meant to imply this gradient acted over the whole period. Closed circles represent the present geothermal gradient.

Yanagisawa et al., 1989]. The Yunagaya group is covered by the Hirono formation that yields a variety of fossil planktons which are correlated to 3.5–4 Ma [Taketani et al., 1986]. The group was uplifted at the end of Miocene or Early Pliocene [Yanagisawa et al., 1989]. Thrusting on the Futaba fault vertically tilted the group in the Hirono area (Fig. 4). The thickness of the stratigraphic section that was truncated by the faulting is not well constrained, but the equation (4) is linear with respect to depth  $z$  so that the missing section does not affect the estimated paleogeothermal gradient—only affects the constant in the equation. Therefore, we use the stratigraphic thickness as  $z$  in the equation. Measurement of vitrinite reflectance of the Yunagaya group therefore allows an estimation of the maximum geothermal gradient in the Miocene.

Figure 3b shows the measured reflectance versus depth that is approximated by stratigraphic thickness. The linear regression of the data gives a slope  $\Gamma = 0.57 \pm 0.13 \text{ km}^{-1}$  which is converted to a temperature gradient of  $45 \pm 10^\circ \text{ Ckm}^{-1}$ . Nakamura and Wakita [1982] show the present geothermal gradient is  $18^\circ \text{ Ckm}^{-1}$ . The paleogeothermal gradient before the Pliocene was therefore higher than at present.

## DISCUSSION

The maximum paleogeothermal gradient for Tanakura occurred in the Late Miocene to early Pliocene, and would

represent a regional geothermal state, as the area has been free from volcanism since early Middle Miocene (~15 Ma).

For the Hirono, it is less clear than for Tanakura when the temperature reached its maximum because of the long period between deposition and uplift. Although there were volcanic activities in the NE Japan fore-arc at 21–25 and 15–16 Ma, small volcanos were produced tens of kilometers away from the studied areas. Thus regional rather than local paleogeothermal gradients are estimated. Figure 5 shows the inferred thermal history. In the Tanakura and Hirono cases, the paleogeothermal gradient in the Miocene was greater than at present, though the estimated gradients represent the upper bound in the period between the maximum burial and uplift. Subsurface temperature appears to have fallen through the Pliocene to Quaternary in these areas.

Cooling appears to have occurred on a regional scale across the arc as secular changes in arc-volcanism exhibit parallel trends. The silica-normalized  $\text{K}_2\text{O}$  content of volcanics at the present volcanic front shows a gradual increase through the Plio-Pleistocene time, suggesting a cooling of their source region [Ban et al., 1992]. In the same period, the volcanic front retreated westward [Ohguchi et al., 1989]. Volcanic activity, as estimated from production rate, has also waned [Sato et al., in preparation]. The regional cooling may be the surface expression of a secular change in the underlying mantle wedge.

Many factors control variation in temperature of the wedge. Davies and Stevenson [1992], who take temporal changes into account in numerical modelling, stressed the roles of subduction dip angles and the mechanical coupling between the slab and asthenosphere. However, little is known about the temporal variation of the coupling under the NE Japan. Volcanic front lies above the subducting slab with a depth of about 110 km that is thought to be determined by pressure-dependent dehydration of the slab [Tatsumi, 1986]. The shallowing of the slab is suggested by the observation of the westward migration of the front [Ohguchi et al., 1989] and by the inference of insignificant migration of the Japan trench after Middle Miocene [Lallemand et al., 1992]. Therefore, the observed cooling from the Pliocene may represent the decrease in the subduction dip angle.

*Acknowledgments.* I am grateful to Drs. N. Kitamura, H. Nakagawa, K. Otsuki, T. Yoshida and H. Sato for discussions and encouragement. The Tanakura samples were provided from Mr. J. Miyakita of the Tohoku Agricultural Administration. Thanks are also due to the anonymous reviewers for the improvement of the manuscript.

## REFERENCES

- Amano, K. and H. Takahashi, 1986. Tertiary sequence in the Tanakura area (in Japanese). in: Kanto District, edited by M. Omori et al., Kyoritsu Publ., Tokyo, 132–134.

- Ban, M., Y. Oba, K. Ishikawa and T. Takaoka, 1992. K-Ar dating of Mutsu-Hiuchidake, Osoreyama, Nanashigure, and Aoso volcanoes of the Aoso-Osore volcanic zone—The formation of the present volcanic zonation of the Northeast Japan arc (in Japanese with English abstract). *Jour. Min. Petrol. Econ. Geol.*, 87, 39–49.
- Barker, C.E. and M.J. Pawlewicz, 1986. The orrelation of vitrinite reflectance with maximum temperature in humic organic matter. in: *Paleogeothermics—evaluation of geothermal conditions in the geologic past*, edited by G. Buntebarth and L. Stegena, Springer-Verlag, New York, 79–94.
- Davies, J.H. and D.J. Stevenson, 1992. Physical model of source region of subduction zone volcanism. *J. Geophys. Res.*, 97, 2037–2070.
- Dow, W.G., 1977. Kerogen studies and geological interpretation. *J. Geochem. Explor.*, 7, 97–99.
- Hood, A., C.C.M. Gutjahr and R.I. Heacock, 1975. Organic metamorphism and the generation of petroleum. *Bull. Am. Assoc. Petrol. Geol.*, 59, 986–996.
- Jolivet, L. and Tamaki, K., 1992. Neogene kinematics in the Japan Sea region and volcanic activity of the Northeast Japan arc. *Proc. ODP Sci. Results*, 127/128, College Station, 1311–1331.
- Kimura, K., 1988. Basement tectonics in Japanese oil fields (in Japanese), *Ann. Rep. JPC Petrol. Explor. Center*, S63, 14–17.
- Kobayashi, Y., 1983. On the initiation of subduction of plates (in Japanese). *Earth Monthly*, 5, 510–514.
- Koizumi, I., 1986. Diatom biochronology of the Neogene in the Joban coal-field, Northeast Japan—Yunagaya, Shirado, and Takaku Groups (in Japanese with English abstract). in *Essays in Geology*, edited by H. Nakagawa et al., Toko Publ., Sendai, 175–191.
- Lallemand, S.E., Schnurle, P. and Manoussis, S., 1992. Reconstruction of subduction zone paleogeometries and quantification of upper plate material losses by tectonic erosion. *J. Geophys. Res.*, 97, 217–239.
- Middleton, M.F., 1982a. The subsidence and thermal history of the Bass Basin, Southeastern Australia. *Tectonophysics*, 87, 383–397.
- Middleton, M.F., 1982b. Tectonic history from vitrinite reflectance. *Geophys. J. roy. Astron. Soc.*, 68, 121–132.
- Middleton, M.F. and P.W. Schmidt, 1982. Paleogeothermics of the Sydney Basin. *J. Geophys. Res.*, 87, 5351–5359.
- Morrow, D.W. and D.R. Issler, 1993. Calculation of vitrinite reflectance from thermal histories: a comparison of some methods. *Am. Assoc. Petrol. Geol. Bull.*, 77, 610–624.
- Nakamura, K., 1983. Possible nascent trench along the eastern Japan sea as the convergent boundary between Eurasian and North American plates (in Japanese with English abstract). *Bull. Earthq. Inst., Univ. Tokyo*, 58, 711–722.
- Nakamura, Y. and H. Wakita, 1982. Terrestrial heat flow around the aseismic front of the Japanese island arc. *Tectonophysics*, 81, T25 ñ T36.
- Oda, M., 1986. Some aspects and problems concerned with microfossil biochronology for the Neogene in Central and Northeast Honshu, Japan (in Japanese with English abstract). in: *Essays in Geology*, edited by H. Nakagawa et al., Toko Publ., Sendai, 297–312.
- Ohguchi, T., T. Yoshida and K. Okami, 1989. Historical change of the Neogene and Quaternary volcanic field in the Northeast Honshu arc, Japan (in Japanese with English abstract). *Mem. Geol. Soc. Jpn.*, 32, 431–455.
- Ohki, J., N. Watanabe, K. Shuto and T. Itaya, 1993. Shifting of the volcanic fronts during Early to Late Miocene in the Northeast Japan arc. *Island Arc*, 2, 87–93.
- Otsuki, K., 1975. Geology of the Tanakura Shear Zone and adjacent area (in Japanese with English abstract). *Contrib. Inst. Geol. Paleontol. Tohoku Univ.*, 76, 1–71.
- Sato, H., T. Yoshida and A. Yamaji, Secular variation of volcanic activity in the NE Japan arc since the mid Tertiary. *J. Geophys. Res.*, in preparation.
- Sugai, K. et al., 1957, *Geological Maps of the Coal Fileds of Japan*, 1, Explanatory Text of the Joban Coal Field (in Japanese with English abstract). *Geol. Surv. Jpn*, 143pp.
- Suzuki, S., 1989. The Cenozoic geothermal history of Northeast Japan (in Japanese with English abstract). *Mem. Geol. Soc. Jpn.*, 32, 321–337.
- Taketani, Y. et al., 1986. Microfossils from the Futaba Group (in Japanese). *Fukushima Prefectural Mus. Rep.*, 12, 1–53.
- Tatsumi, Y., 1986. Formation of the volcanic front in subduction zones. *Geophys. Res. Lett.*, 13, 717–720.
- Tohoku Agricultural Administration, 1986. Hydrological report for the Tanakura Area (in Japanese). Tohoku Agricultural Administration, Sendai, 150pp.
- Uyeda, S., 1972. Heat flow. in: *The Crust and Upper Mantle of the Japanese Area*, Part 1, Geophysics, edited by S. Miyamura and S. Uyeda, *Earthquake Res. Inst., Tokyo*, 96–105.
- Yamaji, A., 1986. Analysis of vitrinite reflectance-burial depth relations in dynamic geological settings by the direct integration method. *J. Jpn. Assoc. petrol. Technol.*, 51, 197–204.
- Yanagisawa, Y. et al., 1989. Tertiary biostratigraphy and subsurface geology of the Futaba district, Joban Coalfield, northeast Japan (in Japanese with English abstract). *Bull. Geol. Soc. Jpn.*, 40, 405–467.
- Yoshida, T., 1992. Geological and geochemical context of Northeast Honshu arc volcanism, 70 Ma to Quaternary. Abstract presented at the 29th IGC, Kyoto, vol 2, pp. 554.