TITLE:

All-or-none switching of transcriptional activity on single DNA molecules caused by a discrete conformational transition

AUTHOR(S):

Yamada, A; Kubo, K; Nakai, T; Yoshikawa, K; Tsumoto, K

CITATION:

Yamada, A ...[et al]. All-or-none switching of transcriptional activity on single DNA molecules caused by a discrete conformational transition. APPLIED PHYSICS LETTERS 2005, 86(22): 223901.

ISSUE DATE:

2005-05-30

URL:

<http://hdl.handle.net/2433/49903>

RIGHT:

Copyright 2005 American Institute of Physics. This article may be downloaded for personal use only. Any other use requires prior permission of the author and the American Institute of Physics.

All-or-none switching of transcriptional activity on single DNA molecules caused by a discrete conformational transition

Ayako Yamada, Koji Kubo, Tonau Nakai, and Kenichi Yoshikawa^{a)}

Department of Physics, Graduate School of Science, Kyoto University & CREST, Kyoto 606-8502, Japan

Kanta Tsumoto

Department of Chemistry for Materials, Faculty of Engineering, Mie University, Tsu 514-8507, Japan

(Received 16 August 2004; accepted 18 April 2005; published online 25 May 2005)

Recently, it has been confirmed that long duplex DNA molecules with a size larger than several tens of kilo-base pairs (kbp), exhibit a discrete conformational transition from an unfolded coil state to a folded compact state upon the addition of various kinds of chemical species that usually induce DNA condensation. In this study, we performed a single-molecule observation on a large DNA, Lambda ZAP II DNA (~41 kbp), in a solution containing RNA polymerase and substrates along with spermine, a tetravalent cation, at different concentrations, by use of fluorescence staining of both DNA and RNA. We found that transcription, or RNA production, is completely inhibited in the compact globule state, but is actively performed in the unfolded coil state. Such an all-or-none effect on transcriptional activity induced by the discrete conformational transition of single DNA molecules is discussed in relation to the mechanism of the regulation of large-scale genetic activity. © 2005 American Institute of Physics. [DOI: 10.1063/1.1937990]

An important goal in biological science is to clarify the mechanism of the self-regulation of transcriptional activity. Currently, many regulatory factors have been discovered, indicating that individual factors interact with certain regions by the recognition of specific sequences along long genomic DNA. It is considered that life is maintained through the use of very complicated networks composed of many key-lock relationships.^{1,2} However, it is still unclear how, in living cells, it is possible to achieve a delicate control of numerous genes with such complicated networks. In a different field of research, cell biologists have been actively studying the relationship between morphology and function of chromatin, both in prokaryote³ and eukaryote,⁴ and have shown that a marked morphological change is accompanied by a significant functional change such as differentiation, cell cycle, and malignancy. Such large structural changes in chromatin have often been discussed in relation to the epigenetic modification of DNA and binding proteins like histone, in terms of methylation, acetylation, etc.^{5,6} On the other hand, for the past decade it has been established that giant DNA molecules larger than several tens of kilo-base pairs exhibit a large discrete conformational transition between unfolded coil and folded compact states⁷ upon the addition of various DNA condensing agents including biochemical species.^{8,9} Thus, it is strongly suspected that this unique property of the folding transition of giant DNA should somehow be closely related to its biological function/activity. However, there have been some apparently conflicting studies on the biological activity of condensed DNA,^{10–17} suggesting that the manner of “condensation” of DNA has not yet been adequately clarified biophysically. For example, it has been reported that “condensed DNA” shows rapid cyclization¹⁰ and also higher transcriptional activity,¹¹ compared to the uncondensed DNA. In contrast, the inhibition of transcription¹² and depression of the activity of endonuclease¹³ have also been reported. In the present letter we report our results in the direct observation

of the on/off change of transcriptional activity induced by the folding/unfolding transition of a giant DNA in the presence of spermine, a natural tetracation.

We used Lambda ZAP II DNA (Stratagene, CA), which includes a single T7 promoter region, as a template for transcription by T7 RNA polymerase (T7 RNAP; Invitrogen, MD). The transcriptional reactions were performed under conditions similar to those described in Ref. 12, with 100 μ L of well-mixed reaction solution containing 5 mM dithiothreitol (provided with the polymerase kit), 0.5 mM each of ATP, CTP, and GTP, 10 μ M UTP (Roche Diagnostics, Switzerland), 0.5 μ M ChromaTide BODIPY TMR-14-UTP (BODIPY; Molecular Probes, Inc., OR), 0.3 μ M 4, 6-diamidino-2-phenylindole (DAPI; Wako Pure Chemicals Industries, Ltd., Japan), and 200 (or 700) μ M spermine tetrachloride (Nacalai Tesque, Japan) in T7 buffer (polymerase kit). After 80 (or 10, respectively) ng/mL Lambda ZAP II DNA and 50 U of T7 RNAP were added under gentle stirring, the solution was incubated at 37 °C for 30 min. To avoid aggregation of DNA molecules, the concentration of DNA was adjusted to be very low: around 10 ng/mL. Although RNAP activity on a single DNA molecule has been observed in a variety of adroit ways in previous pioneering studies,^{18–21} here we used a very simple and concise procedure to visualize both the DNA conformation and its transcripts: after the transcriptional reaction, we elongated DNA molecules on the glass surface by gentle shear stress.²² Both the Lambda ZAP II DNA and RNA molecules were visualized with a Carl Zeiss fluorescence microscope (Axiovert 200 TV) equipped with a 100 $\times$ oil-immersed lens and a high-sensitivity Hamamatsu SIT TV camera. DAPI bound to DNA and BODIPY incorporated into RNA by polymerization was excited at wavelengths of 365 and 535 nm, respectively. We also used an atomic force microscope (AFM) equipped with a fluorescence microscope (AFM; NBV100, FM; IX70, Olympus Optical Co. Ltd., Japan) to observe the same DNA and RNA molecules. AFM was performed in Tapping ModeTM in aqueous solution.

^{a)}Electronic mail: yoshikaw@scphys.kyoto-u.ac.jp

FIG. 1. (Color) Fluorescence microscopic images of single Lambda ZAP II DNA at (a) [spermine]=200 μ M and (b) [spermine]=700 μ M. Upper pictures: blue and red colors correspond to the fluorescence from labeled DNA and RNA (transcripts), respectively. Lower pictures: quasi-three-dimensional representations of the spatial distribution of fluorescence intensity on DNA.

Figure 1 shows typical fluorescence microscopic images of Lambda ZAP II DNA: (a) in an unfolded elongated coil state (200 μ M spermine) and (b) in a folded compact state (700 μ M spermine). In the lower pictures in Fig. 1, the intensity distribution of fluorescence emitted from DAPI is given as a quasi-three-dimensional representation. In Fig. 1, the elongated state of DNA (a) is obtained as described earlier. The red spot in the upper left picture in Fig. 1 indicates the active production of RNA, where fluorescence from BODIPY is observed. In contrast, no red fluorescence is emitted from the folded compact DNA, as shown in the upper right picture in Fig. 1.

Figure 2 shows experimental data on the dependence of transcriptional activity on the spermine concentration through conventional test-tube measurement, i.e., the activity is measured as the ensemble average of many DNA molecules. The experimental condition in this measurement is essentially the same as in the single-DNA observation. The results in Fig. 2 indicate that the change in transcriptional activity around [spermine]=400 μ M is rather steep, although it is not clear, based solely on the observation of the ensemble of DNAs, whether the change in transcriptional

FIG. 2. Transcriptional activity vs concentration of spermine (4+). The vertical broken line indicates the region of the conformational transition as evaluated from the fluorescence microscopic observation of bulk DNA solution. (Modified from the original figure in Ref. 12.)

FIG. 3. (Color) Fluorescence microscopic image of Lambda ZAP II DNA at the coexistence condition between unfolded and folded state with 350 μ M spermine. RNA transcripts exist only on the unfolded DNA (indicated by the arrowhead) and are absent on the folded DNA. Blue and red colors correspond to the fluorescence from labeled DNA and RNA, respectively.

activity is on/off type or cooperative and continuous type. The single-chain observation clarifies that this activity is completely inhibited for all of the compact DNA molecules.

In order to confirm the on/off nature of the transcriptional reaction, we have performed the measurement of the transcriptional activity at the coexistence condition (350 μ M spermine) between unfolded coil and folded compact states. Figure 3 exemplifies the result of such observation, where RNA transcripts exist on the unfolded DNA (indicated by the arrowhead), and the absence of transcripts on the folded DNA. It is to be noted that in this observation the unfolded DNA is not fully elongated by shear stress.

To measure the transcription along a DNA with higher resolution, we performed AFM/FM observation. Figure 4 shows (a) the AFM image of an unfolded DNA (200 μ M spermine) together with (b) the fluorescence microscopic image of RNA labeled with BODIPY. As the AFM image on the transcripts together with the DNA chain was obtained under wet conditions inside the aqueous solution containing substrates necessary for the RNA synthesis, the resolution is not so good compared to the usual AFM image of DNA in a dry state. Due to the resolution limit with optical observation, the fluorescence observation of BODIPY gives only a blurred image around the region that is white in the AFM image. Based on a comparison of the AFM and fluorescence images, in Fig. 4(c) we show a schematic representation of our observation. Although we could not recognize individual RNAP/RNA molecules, the white part in the AFM image is attributable to RNAP molecules that stay near the promoter in transcription.

FIG. 4. (Color) A single DNA molecule transcribed into RNA, as observed simultaneously by (a) AFM and (b) FM. (c) Schematic representation.

TABLE I. Transcriptional activity at different spermine concentrations obtained from single-DNA measurement.

[Spermine]	Number of active DNA	Total number of DNA ^a	Fraction of active DNA
200 μ M	47	79	0.60
700 μ M	0	103	0.00

^aUnfragmented DNA only.

Table I summarizes the results regarding transcriptional activity with the single-chain observation. At a spermine concentration of 700 μ M, the activity of transcription is completely inhibited and all of the DNA molecules assumes a folded compact state. In contrast, at 200 μ M, where DNA molecules show an unfolded coil conformation, most of the DNAs exhibit the transcriptional activity. It is now clear that individual DNA shows on/off switching in transcriptional activity that is accompanied by a folding transition.

The above result indicates that the transcriptional activity of an individual chain is strongly related to its conformation, i.e., to a drastic change in its density. The density ratio between the folded compact globule and unfolded coil states $\rho_{\text{globule}}/\rho_{\text{coil}}$ equals the inverse ratio of the cube of the respective radii $R_{\text{coil}}^3/R_{\text{globule}}^3$. The coil radius is given as $R_{\text{coil}} \sim \lambda N^{3/5}$ under good solvent conditions for a chain,²³ where λ is the persistence length (about 150 bp for DNA at room temperature) and N is defined as the number of independent segments of length λ per contour length L (40820 bp for Lambda ZAP II DNA). On the other hand, the globule size R_{globule} is proportional to $(\lambda s N)^{1/3}$, where s is the cross-sectional area of the chain ($3 \text{ nm}^2 \approx 27.5 \text{ bp}^2$). The density ratio $\rho_{\text{globule}}/\rho_{\text{coil}}$ can thus be estimated as $s^{-1} \lambda^2 N^{4/5}$. For the giant DNA adopted in this experiment, this ratio is $\sim 10^5$. It has been confirmed from the measurement of the Brownian motion of an individual giant DNA molecule that the relative ratio of the hydrodynamic radii is on the order of 10^4 – 10^5 on giant DNA.²⁴ The high density present in a globular conformation should prevent the binding of enzymes (here, RNA polymerase) to DNA, which could lead to a total inhibition of transcriptional activity. Table I indicates that the relative activity decreased from 0.60 at 200 μ M spermine to 0.00 at 700 μ M spermine. The abrupt change in density, with a discrete conformational transition, is the direct cause of the on/off switching of transcriptional activity. The present study shows that DNA transcription undergoes an on/off change that is accompanied by the conformational transition of DNA, which is induced by a small change in the solution; i.e., the spermine concentration in the present study. It has been reported that a small change in concentration such as of a 1:1 salt, RNA and ATP, can cause a large discrete transition in the conformation of giant DNA.²⁴ Such a conformational transition in giant DNA can cover a large genetic region, possibly including tens or hundreds of genes. Thus, it is strongly suspected that such a conformational transition corresponds to the opening/closing of a folder that contains several genes.

In the present study, we have shown that transcription is completely inhibited upon compaction. In contrast to this inhibitory effect, it has been reported that transcriptional activity was promoted by the condensation on small circular plasmid DNA molecules.¹¹ Circular DNA with superhelicity tends to undergo a continuous-type folding transition,²⁵ in

contrast to the discrete conformational transition of linear giant DNA.⁷ circular DNA exhibits a stepwise collapse, which causes local stress along double-stranded DNA chains.

It is expected that the generation of such local stress in circular DNAs induced by the collapsing transition may be associated with the enhancement of enzymatic activity, through weakening of the double-strand structure. A similar discussion may be applicable to the genetic activity in a loop attached to a scaffold in giant genomic DNA in chromatin,⁴ although other factors such as topoisomerases and structural maintenance of chromosomes family proteins, etc., are also considerable. Since the folding transition is a characteristic event generated on a large scale along a giant DNA, it may play a role in the large-scale regulation of multiple genes,²⁴ complementing key-lock interactions in individual genes.

The authors are grateful to Dr. Naoko Makita for providing helpful advice on the experimental methods, Mr. Hiroyuki Kitahata for help on analyzing video images, and Dr. Tatsuo Akitaya and Dr. Damien Baigl for their kind comments. This work is supported by the Grants in Aid for the 21st Century COE “Center for Diversity and Universality in Physics” and for Scientific Research in Priority Areas “Chemistry of Biological Processes Created by Water and Biomolecules” from the Ministry of Education, Culture, Sports, Science and Technology of Japan.

¹P. B. Warren and P. R. Wolde, Phys. Rev. Lett. **92**, 128101 (2004).

²J. Paulsson, Nature (London) **427**, 415 (2004).

³D. Frenkiel-Krispin, S. Levin-Zaidman, E. Shimoni, S. G. Wolf, E. J. Wachtel, T. Arad, S. E. Finkel, R. Kolter, and A. Minsky, EMBO J. **20**, 1184 (2001).

⁴A. Wolffe, *Chromatin*, 3rd ed. (Academic Press, San Diego, 1998).

⁵J. Zhang, F. Xu, T. Hashimshony, I. Keshet, and H. Cedar, Nature (London) **420**, 198 (2002).

⁶M. A. Karymov, M. Tomschik, S. H. Leuba, P. Caiafa, and J. Zlatanova, FASEB J. **15**, 2631 (2001).

⁷K. Yoshikawa, M. Takahashi, V. V. Vasilevskaya, and A. R. Khokhlov, Phys. Rev. Lett. **76**, 3029 (1996).

⁸V. A. Bloomfield, Curr. Opin. Struct. Biol. **6**, 334 (1996).

⁹E. Raspaud, M. Olvera de la Cruz, J.-L. Sikorav, and F. Livolant, Biophys. J. **74**, 381 (1998).

¹⁰D. Jary and J.-L. Sikorav, Biochemistry **38**, 3223 (1999).

¹¹I. Baeza, P. Gariglio, L. M. Rangel, P. Chavez, L. Cervantes, C. Arguello, C. Wong, and C. Montañez, Biochemistry **26**, 6387 (1987).

¹²K. Tsumoto, F. Luckel, and K. Yoshikawa, Biophys. Chem. **106**, 23 (2003).

¹³H. Oana, K. Tsumoto, Y. Yoshikawa, and K. Yoshikawa, FEBS Lett. **530**, 143 (2002).

¹⁴M. A. Krasnow and N. R. Cozzarelli, J. Biol. Chem. **257**, 2687 (1982).

¹⁵S. B. Zimmerman and A. P. Minton, Annu. Rev. Biophys. Biomol. Struct. **22**, 27 (1993).

¹⁶K. Sekine, T. Hase, and N. Sato, J. Biol. Chem. **277**, 24399 (2002).

¹⁷N. Sasaki, H. Kuroiwa, C. Nishitani, H. Takano, T. Higashiyama, T. Kobayashi, Y. Shirai, A. Sakai, S. Kawano, K. Murakami-Murofushi, and T. Kuroiwa, Mol. Biol. Cell **14**, 4758 (2003).

¹⁸Z. Gueroui, C. Place, E. Freyssingas, and B. Berge, Proc. Natl. Acad. Sci. USA **99**, 6005 (2002).

¹⁹Y. Harada, O. Ohara, A. Takatsuki, H. Itoh, N. Shimamoto, and K. Kinoshita Jr., Nature (London) **409**, 113 (2001).

²⁰H. Kabata, O. Kurosawa, I. Arai, M. Washizu, S. A. Margaron, R. E. Glass, and N. Shimamoto, Science **262**, 1561 (1993).

²¹C. Rivetti, S. Codeluppi, G. Dieci, and C. Bustamante, J. Mol. Biol. **326**, 1413 (2003).

²²Y. Matsuzawa and K. Yoshikawa, Nucleosides Nucleotides **13**, 1415 (1994).

²³A. Y. Grosberg and A. R. Khokhlov, *Statistical Physics of Macromolecules* (American Institute of Physics, Melville, NY, 1994).

²⁴K. Yoshikawa, J. Biol. Phys. **28**, 701 (2002).

²⁵Y. S. Velichko, K. Yoshikawa, and A. R. Khokhlov, J. Chem. Phys. **111**, 9424 (1999).