TITLE:

Electronic excitation and ionization spectra of cyclopentadiene: Revisit by the symmetry-adapted cluster-configuration interaction method

AUTHOR(S):

Wan, J; Ehara, M; Hada, M; Nakatsuji, H

CITATION:

Wan, J ...[et al]. Electronic excitation and ionization spectra of cyclopentadiene: Revisit by the symmetry-adapted cluster-configuration interaction method. JOURNAL OF CHEMICAL PHYSICS 2000, 113(13): 5245-5252

ISSUE DATE:

2000-10-01

URL:

<http://hdl.handle.net/2433/50179>

RIGHT:

Copyright 2000 American Institute of Physics. This article may be downloaded for personal use only. Any other use requires prior permission of the author and the American Institute of Physics.

Electronic excitation and ionization spectra of cyclopentadiene: Revisit by the symmetry-adapted cluster–configuration interaction method

Jian Wan, Masahiro Ehara, Masahiko Hada, and Hiroshi Nakatsuji^{a)}

Department of Synthetic Chemistry and Biological Chemistry, Graduate School of Engineering,
Kyoto University, Sakyo-ku, Kyoto 606-8501, Japan

(Received 26 May 2000; accepted 7 July 2000)

Electronic excitation and ionization spectra of cyclopentadiene (CP) were reinvestigated by the symmetry-adapted cluster (SAC) and SAC–configuration interaction (SAC-CI) method with an extended basis set and a wide active orbital space. To give a satisfactory interpretation of the general profile of the observed excitation spectrum, 40 low-lying excited singlet and triplet states (with excitation energies of up to 9.5 eV) were computed. The calculated excitation energies were greatly improved compared to those reported previously. All of the peaks in the experimental spectrum were reassigned theoretically with small deviations. The natures of the low-lying valence and Rydberg-excited states were discussed in detail, and the results were also compared with those of some other recent theoretical studies. The ionization energies calculated by the SAC-CI general-R method agree well with the experimental peaks in the photoelectron spectrum. A number of two-electron shake-up states were calculated below 23 eV. © 2000 American Institute of Physics. [S0021-9606(00)30437-8]

I. INTRODUCTION

Electronic excited states of cyclopentadiene (CP) have been extensively investigated by various spectroscopic experiments due to its importance as a model compound for photochemical reactions of various cyclic dieny systems. The locations and assignments of various singlet intravalence $\pi-\pi^*$ and Rydberg transitions have been a major subject of optical ultraviolet (UV) spectroscopic studies.^{1–3} Spin-forbidden singlet–triplet transitions have so far been investigated by electron energy-loss (EEL) spectroscopy.⁴ Ionization potentials (IPs) of CP have also been studied by photoelectron spectroscopy (PES).⁵

On the other hand, the experimental work has received relatively little support from *ab initio* calculations. Only a few theoretical interpretations of observed spectra based on *ab initio* calculations have been reported; symmetry-adapted cluster (SAC) and SAC–configuration interaction (SAC-CI) calculations⁶ in 1985, random phase approximation (RPA) calculations⁷ in 1991, complete active space second-order perturbation (CASPT2) calculations⁸ in 1993, and multireference Møller–Plesset (MRMP) perturbation calculations⁹ in 1996. The earliest SAC/SAC-CI study⁶ was a pioneering work, but because of the lack of the computational resources in those days, the limited basis sets and small active space impaired the quality of the results. The RPA calculations were not accurate enough to allow a full interpretation of the spectrum. In the latter CASPT2 and MRMP studies, fairly large basis sets were used; an atomic natural orbital (ANO) basis set augmented by ring-centered diffuse *s*-, *p*- and *d*-functions in CASPT2, and Dunning's cc-pVTZ basis set augmented by charge-centered diffuse *s*-, *p*-, and *d*-functions

in MRMP. These two similar theoretical methods gave very similar computational results and gave a relatively reasonable interpretation of singlet valence excitations lower than 8 eV. The ionized states of CP were investigated by, together with the excited states, the SAC/SAC-CI method,⁶ but not by the CASPT2 and MRMP methods.

In this study, we re-examined the singlet and triplet valence and Rydberg excitations and ionizations of CP using a better-quality basis set and a larger active space by the SAC/SAC-CI singles and doubles operator R_K^+ (SD-R) calculations. Furthermore, to give a theoretical interpretation of correlation (satellite) peaks characterized by the multielectron process in the PES spectrum,⁵ the SAC/SAC-CI general-R method¹⁰ was also employed for an accurate calculation of the inner-valence region of the ionization spectrum. In Sec. II, we outline the computational details. Calculated results and discussions are presented in Sec. III. A summary is given in Sec. IV.

II. COMPUTATIONAL DETAILS

The details of the SAC/SAC-CI method for calculating ground, excited, and ionized states of molecules have been presented elsewhere.^{10–18} An experimental ground-state equilibrium geometry of CP with C_{2v} symmetry¹⁹ was used and the CP ring was put on the *yz*-plane with *z* as the principle axis (Fig. 1). The calculated excitation energies are, therefore, vertical in nature.

Dunning's augmented correlation consistent basis set AUG-cc-pVDZ (Ref. 20) was used for C atoms, and cc-pVDZ (Ref. 21) was used for H atoms. Additionally, a set of diffuse functions (*3s3p3d*) selected from the studies of Kaufmann *et al.*²² was placed on the molecular center of gravity. These molecule-centered diffuse functions are the same as those used by Trofimov and Schirmer.²³ All SCF

^{a)} Author to whom correspondence should be addressed. Electronic mail: hiroshi@sbchem.kyoto-u.ac.jp

FIG. 1. Cyclopentadiene (C_5H_6) molecule.

calculations were performed with the GAUSSIAN 98 package.²⁴ The total number of basis functions is 185. This augmented basis set is adequate for describing widely different electronic structures of molecules in the ground state, singlet and triplet valence and Rydberg excited states and ionized states, and for describing valence and diffuse characters and different correlation effects.

The present SAC/SAC-CI calculations were performed with the local version of the SAC-CI module.¹⁸ The active space consists of 13 occupied orbitals and 167 virtual orbitals; only the $1s$ core MOs of C atoms were frozen. In the SAC ground-state calculation, all single-excitation and selected double-excitation operators S_I^+ were included in the linked term. The energy threshold λ_g for perturbation selection¹⁶ (PS) was 1.0×10^{-5} a.u. For the unlinked term, we included only the products of the double-excitation operators $S_I^+ S_J^+$ when the coefficients C_I and C_J , estimated by SD-CI in practice, were larger than 1.0×10^{-3} .

In the SAC-CI SD-R calculations of excited and ionized states, all single-excitation operators and selected double-excitation operators were included in the linked operators R_K^+ . Perturbation selection was performed as follows. First,

we selected the main reference configurations from SE-CI with a coefficient greater than 0.1, and then selected the double-excitation operator whose second-order perturbation matrix element with one of the main reference configurations is larger than the threshold λ_e (1.0×10^{-6} a.u.). In the unlinked term $\sum_{K,I} d_K C_I R_K^+ S_I^+ |0\rangle$, we included the double-excitation operators $\{S_I^+\}$ whose coefficients (C_I) were larger than 1.0×10^{-3} in the SAC ground-state calculation, and as the R_K^+ operator, we included the single- and double-excitation operators whose coefficients (taken from SD-CI in practice) were greater than 0.05.

In the SAC-CI general-R (Ref. 10) calculations of the outer- and inner-valence ionization potentials of CP, a preliminary SD-CI calculation was performed to select the operators for the exponential generation (EG) scheme¹⁷ by which we construct important triple and quadruple excitation operators required for a quantitative description of the multielectron process in the ionization spectrum. The generated higher-order excitation operators were further selected by the perturbation selection¹⁶ procedure with the energy threshold of 1.0×10^{-6} hartree. Therefore, all single (ionization) operators and selected higher operators (SDTQ) were included in the linked term of SAC-CI general-R equation. In the unlinked term, we included all the double-excitation operators $\{S_I^+\}$ in the SAC ground-state calculation, and as the R_K^+ operator, we included all single operators and selected higher-order operators up to quadruple.

Table I shows the previous⁶ and present SAC/SAC-CI SD-R computational conditions. Fifteen years ago, the computational resources were substantially smaller than those used here; the available main memory and disk space were very limited, though the previous calculations were certainly “big” by the standard of that time. With the tremendous advances in computer technology, we can use a much better basis set and much larger active space. This improvement, by itself, made the calculated results significantly better, even

TABLE I. Comparison of the previous and present SAC/SAC-CI computational conditions for cyclopentadiene.

Previous work (Ref. 6)					Present work		
Basis sets	DZ+Rydberg C: $[4s2p]$ H: $[2s]$ Rydberg ($2s2p1d$) on the molecular center of gravity				AUG-cc-pVDZ+Rydberg C: AUG-cc-pVDZ H: cc-pVDZ Rydberg ($3s3p3d$) on the molecular center of gravity		
Active space	71 ^a [5, ^b 54 ^c]				185 ^a [5, ^b 180 ^c]		
Energy threshold for perturbation selection (PS)	$\lambda_g = 4.0 \times 10^{-5}$, $\lambda_e = 5.0 \times 10^{-5}$				$\lambda_g = 1.0 \times 10^{-5}$, $\lambda_e = 1.0 \times 10^{-6}$		
Dimension of linked operators after PS	Singlet $A_1(G)^d$	Triplet	Ionized		Singlet	Triplet	Ionized
	2312				14 871		
	A_1 1285	1793	1130		70 030	65 910	6649°/7410 ^f
	A_2 3203	912	195		57 921	61 985	2610°/3999 ^f
	B_1 1678	1200	357		96 643	33 945	3697°/4789 ^f
	B_2 984	1734	888		47 262	55 232	6305°/7218 ^f

^aThe total number of MOs.

^bThe number of frozen orbitals, 5 means “ $1s$ -core.”

^cThe size of the active space (occupied MOs plus unoccupied MOs).

^dSAC ground state.

^eDimension in the SAC-CI SD-R method.

^fDimension in the SAC-CI general-R method.

though the basic method used here is the same as the previous one, i.e., the SAC/SAC-CI method. The improvements in the SAC/SAC-CI program and algorithm, etc., are also important, but for the comparisons in Table I, the improvements in the computer technology play the greatest role.

III. RESULTS AND DISCUSSION

With the basis set mentioned above, the SCF energy of CP is $-192.814\,698$ hartrees. The SAC ground-state energy is $-193.172\,258$ hartrees, and the correlation energy is $0.357\,559$ hartrees. The corresponding previous values were $-192.737\,32$, $-192.965\,84$, and $0.228\,52$ hartrees, respectively.

To distinguish between valence and Rydberg-type transitions, the second moments of the charge distribution $\langle r^2 \rangle$ and its components $\langle x^2 \rangle$, $\langle y^2 \rangle$, and $\langle z^2 \rangle$ were calculated. Since all of the $\pi-\pi^*$ transitions in CP belong either to A_1 or B_2 symmetry, the A_2 and B_1 Rydberg states are not perturbed by the valence $\pi-\pi^*$ transitions. The oscillator strength has also been used by several authors as an indicator of valence versus Rydberg character in the case of strong mixing of valence and Rydberg states. We will use all these knowledge to analyze the characters of the excited states.

The SCF occupied valence orbitals and some important virtual orbitals are shown in Table II, together with their orbital energies, symmetries, and characters. Table III shows singlet and triplet excitation energies, oscillator strengths and second moments calculated by the SAC/SAC-CI SD-R method. In Table IV, the singlet and triplet excitation energies calculated by the SAC-CI SD-R method are compared to the experimental values, and to the CASPT2 and MRMP results. Figure 2 shows a comparison of the experimental EEL spectrum⁴ and the theoretical SAC-CI spectrum of CP.

A. Singlet valence excited states

Roughly speaking, CP is regarded as two ethylene π systems with two occupied π and two antibonding π^* molecular orbitals (MO's). Therefore, there are four separate singlet excited states within these π MO's, and Mulliken²⁵ designated the four lowest $\pi-\pi^*$ singlet transitions in this simple model as $N \rightarrow V_1$, V_2 , V_3 , and V_4 in the order of the transition energy. All of these excitations are electric-dipole allowed, and the transitions $N \rightarrow V_1$, V_2 , V_3 , and V_4 are characterized as 1B_2 , 1A_1 , 1A_1 , and 1B_2 , respectively, due to their irreducible representations.

The lowest observed³ singlet transition, $N \rightarrow V_1(\tilde{X}{}^1A_1 \rightarrow {}^1B_2)$ designated as peak I in Fig. 2, begins at 4.75 eV and ends at 5.90 eV with the absorption maximum located at 5.34 eV. Frueholz *et al.*⁴ observed the intensity maximum of a broad transition at 5.26 eV by electron impact spectroscopy and assigned it to $N \rightarrow V_1(\tilde{X}{}^1A_1 \rightarrow {}^1B_2)$. Our current lowest singlet excited state was computed to be $1{}^1B_2$ at 5.54 eV with oscillator strength of 0.1305 , which is clearly related to the intense feature of peak I. The second moment $\langle r^2 \rangle$ is 72.35 a.u.², which clearly means a valence-excited state. The nature of the $1{}^1B_2$ state is $\pi-\pi^*$ excitations from the HOMO ($1a_2$: 18th MO) to the virtual π^* orbital (48th MO). There is very little mixing of Rydberg functions.

TABLE II. Hartree-Fock occupied valence molecular orbitals (MOs) and important unoccupied MOs.

MOs	Symmetry	Orbital energy (eV)	Nature
Occupied:			
Cores			
6	$4a_1$	-31.362	$\sigma(1s)$
7	$3b_2$	-26.302	$\sigma(2s)$
8	$5a_1$	-26.000	$\sigma(2s)$
9	$6a_1$	-20.571	$\sigma(2s)$
10	$4b_2$	-19.811	$\sigma(2s)$
11	$7a_1$	-18.944	$\sigma(2p)$
12	$1b_1$	-16.071	π
13	$5b_2$	-15.290	$\sigma(2p)$
14	$8a_1$	-14.386	$\sigma(2p)$
15	$9a_1$	-13.786	$\sigma(2p)$
16	$6b_2$	-13.617	$\sigma(2p)$
17 (next HOMO)	$2b_1$	-11.244	π
18 (HOMO)	$1a_2$	-8.368	π
Unoccupied			
19 (LUMO)	a_1	0.048	s -Rydberg
25	b_2	0.345	d_{yz} -Rydberg
29	a_1	0.494	s -Rydberg
31	b_2	0.561	p_y -Rydberg
33	a_1	0.955	s/d_{yz} -Rydberg
34	b_2	0.958	d_{yz} -Rydberg
48	b_1	2.938	π^*
51	b_2	3.696	σ^*
54	a_2	4.095	π^*
56	b_2	4.585	σ^*
58	b_2	4.861	σ^*
59	a_1	4.883	σ^*
63	b_1	6.069	π^*
66	b_2	7.344	σ^*
67	a_2	7.362	π^*

At 6.15 – 6.7 eV, there is another feature, peak II, as shown in Fig. 2. This region is characterized by Rydberg transitions, and therefore will be discussed in the next section.

The region with the most intense inelastic scattering in the electron impact spectrum occurs from approximately 7.2 – 8.7 eV, with the intensity maximum located at 7.88 eV (peak III in Fig. 2). Robin²⁶ suggested that the corresponding feature (8.06 eV) in the optical spectrum results from the superposition of the $N \rightarrow V_2(\tilde{X}{}^1A_1 \rightarrow {}^1A_1)$ and $N \rightarrow V_3(\tilde{X}{}^1A_1 \rightarrow {}^1A_1)$ excitations. Our present calculated results give the $6{}^1B_2$, $5{}^1A_1$, $7{}^1B_2$, $6{}^1A_1$, and $7{}^1A_1$ states in this region. Although there is extensive mixing of Rydberg and valence transitions in the $6{}^1B_2$, $7{}^1B_2$, $6{}^1A_1$, and $7{}^1A_1$ states, we assign these states to Rydberg transitions according to their larger second moments, weaker oscillator strengths, and existence of Rydberg-type main configurations. The $5{}^1A_1$ state is calculated at 8.17 eV with an oscillator strength of 0.5377 , which is the strongest transition in the entire energy region of interest. The second moment $\langle r^2 \rangle$ is 152.02 a.u.², which clearly shows a diffuse character (the $\langle r^2 \rangle$ of the ground state is 68.46 a.u.²), and there is strong Rydberg-mixing in this valence-excited state. The $5{}^1A_1$ state is described by the mixed $\pi-\pi^*$ excitations from HOMO ($1a_2$: 18th MO) and next HOMO ($2b_1$: 17th MO) to the virtual π^* orbitals, which corresponds, to some extent, to

TABLE III. Calculated excitation energies (in eV), oscillator strengths $f(r)$, and second moments for cyclopentadiene.

State	Nature ^a	ΔE^b	Pol ^c	$f(r)$	Second moment			
					$\langle x^2 \rangle$	$\langle y^2 \rangle$	$\langle z^2 \rangle$	$\langle r^2 \rangle$
1 1A_1	ground	0.0		0	25.51	22.02	20.93	68.46
1 1B_2	V^d	5.54	y	0.1305	26.92	21.43	24.00	72.35
1 1A_2	3s-R	5.58		0.0000	44.96	36.44	37.78	119.18
1 1B_1	3p _y -R	6.17	x	0.0270	38.98	71.87	35.02	145.87
2 1A_2	3p _x -R	6.21		0.0000	41.05	35.17	76.98	153.20
2 1B_2	3p _z -R	6.22	y	0.0311	75.85	32.90	34.13	142.88
3 1A_2	3d _{xx} -R	6.69		0.0000	51.40	71.26	36.07	158.73
2 1B_1	3d _{yz} -R	6.71	x	0.0003	38.39	65.04	67.72	171.15
2 1A_1	$V \sim \text{mix.}^d$	6.76	z	0.0009	30.88	27.03	23.69	81.60
4 1A_2	3d _{yy-zz} -R	6.77		0.0000	42.35	47.71	90.41	180.47
3 1B_2	3d _{xz} -R	6.93	y	0.0005	91.86	37.32	85.15	214.33
3 1A_1	3d _{xy} -R	6.96	z	0.0007	91.12	83.29	39.29	213.70
5 1A_2	4s-R	7.07		0.0000	166.20	88.91	88.85	343.96
3 1B_1	4p _y -R	7.24	x	0.0072	98.90	245.10	92.76	436.76
4 1B_2	4p _x -R	7.25	y	0.0078	249.18	89.72	91.93	430.83
6 1A_2	4p _z -R	7.27		0.0000	104.17	96.81	262.05	463.03
7 1A_2	4d _{xx} -R	7.42		0.0000	142.36	192.34	129.62	464.32
4 1B_1	4d _{yz} -R	7.46	x	0.0002	83.37	198.85	200.45	482.67
8 1A_2	4d _{yy-zz} -R	7.48		0.0000	86.10	167.37	238.59	492.06
5 1B_2	4d _{xz} -R	7.55	y	0.0001	217.01	78.96	208.40	504.37
4 1A_1	4d _{xy} -R	7.57	z	0.0010	216.60	208.70	80.41	505.71
9 1A_2	5s-R	7.60		0.0000	382.32	249.15	248.35	879.82
5 1B_1	5p _y -R	7.73	x	0.0083	143.20	374.50	136.59	654.29
6 1B_2	5p _x -R	7.74	y	0.0054	402.27	140.63	396.33	686.23
6 1B_1	3s-R	7.93	x	0.0095	54.28	68.06	48.27	170.61
7 1B_1	5d _{yz} -R	8.08	x	0.0000	82.30	195.47	196.58	474.35
5 1A_1	$V \sim \text{mix.}^d$	8.17	z	0.5377	61.92	54.32	35.78	152.02
7 1B_2	5d _{xz} -R	8.23	y	0.0000	188.92	69.64	181.80	440.36
6 1A_1	5d _{xy} -R	8.27	z	0.0966	155.70	144.22	61.47	361.39
8 1B_1	3p _z '-R	8.47	x	0.0122	37.81	38.86	66.97	143.64
7 1A_1	3p _x '-R	8.56	z	0.0216	78.38	41.77	34.09	154.24
9 1B_1	4p _z '-R	8.59	x	0.0038	34.11	48.24	71.26	138.81
8 1B_2	$\pi-\sigma^*$	9.07	y	0.0248	153.34	58.13	58.81	270.28
8 1A_1	3d _{xz} '-R	9.23	z	0.0290	95.53	40.91	85.19	221.63
9 1B_2	3d _{xy} '-R	9.25	y	0.0053	83.56	75.79	38.17	197.52
10 1B_2	4d _{xy} '-R	9.43	y	0.0104	63.91	54.32	41.16	159.39
9 1A_1	4p _x '-R	9.54	z	0.0016	249.88	92.56	91.76	434.20
1 3B_2	V^d	3.40		0.0	25.61	21.09	22.72	69.42
1 3A_1	V^d	5.18		0.0	25.70	22.40	22.08	70.18
1 3A_2	3s-R	5.56		0.0	45.35	34.80	37.92	118.07
1 3B_2	3p _y -R	6.19		0.0	40.47	75.77	36.42	152.66

^aV and R denote, respectively, valence and Rydberg excited state.^bTransition energy.^cPolarization direction of transition moment.^dSee text.

Robin's superposition of the $N \rightarrow V_2(\tilde{X}^1A_1 \rightarrow ^1A_1)$ and $N \rightarrow V_3(\tilde{X}^1A_1 \rightarrow ^1A_1)$ excitations. From this perspective, our current results support Robin's suggestion.

In the higher energy region, several transitions are observed in peak IV located at 8.7–9.4 eV of the spectrum shown in Fig. 2. The present SAC-CI calculations show that the excitation energies in this region were relatively basis set-dependent, and place 8 1B_2 , 8 1A_1 , 9 1B_2 , 10 1B_2 , and 9 1A_1 states at 9.07, 9.23, 9.25, 9.43, and 9.54 eV with intensities of 0.0248, 0.0290, 0.0053, 0.0104, and 0.0016, respectively. McDiarmid and Sabljic²⁷ experimentally assigned peak IV observed at 9.2 eV to the $N \rightarrow V_4(\tilde{X}^1A_1 \rightarrow ^1B_2)$ valence excitation. RPA calculations gave the second 1B_2 state at 10.46 eV with an oscillator strength of 0.0901. CASPT2

and MRMP gave no excited states in this region.

Finally, we discuss the valence excited state 2 1A_1 . This state was not found in the spectra. Our present calculations gave the 2 1A_1 state at 6.76 eV with a very weak oscillator strength of 0.0009 and a second moment of 81.60 a.u.², which clearly defines it as a valence excited state. Its main configuration was almost purely composed of single excitations, 17th MO $\rightarrow$ 48th MO and 18th MO $\rightarrow$ 54th MO, and there was little double-excitation mixing in this state. CASPT2 gave a value of 6.31 eV with an intensity of 0.0003 and its nature was reported to be a mixture of the configurations $2b_1 \rightarrow 3b_1$ and $1a_2 \rightarrow 2a_2$ and the double excitation of $0.43[(1a_2)^2 \rightarrow (3b_1)^2]$. MRMP gave 6.42 eV with an intensity of 0.0039, and its nature was almost the same as that by

TABLE IV. SAC-CI results compared with the experimental excitation energies (in eV) and other theoretical results for cyclopentadiene.

State	Nature ^a	SAC-CI		Expt. ^c	CASPT2 ^d		State	MRMP ^e		
		$f(r)$	ΔE^b		ΔE	$f(r)$		ΔE	$f(r)$	State
1 ¹ A ₁	ground	0	0.0							
1 ¹ B ₂	V ^f	0.1305	5.54	5.26,5.34	5.27	0.1475	1 ¹ B ₂	5.19	0.1460	1 ¹ B ₂
1 ¹ A ₂	3s-R	forbidden	5.58	5.68	5.65	forbidden	1 ¹ A ₂	5.62	forbidden	1 ¹ A ₂
1 ¹ B ₁	3p _y -R	0.0270	6.17	6.25	6.24	0.0306	1 ¹ B ₁	6.24	0.0174	1 ¹ B ₁
2 ¹ A ₂	3p _x -R	forbidden	6.21	6.26	6.30	forbidden	2 ¹ A ₂	6.27	forbidden	2 ¹ A ₂
2 ¹ B ₂	3p _z -R	0.0311	6.22	6.31	6.25	0.0248	2 ¹ B ₂	6.25	0.0149	2 ¹ B ₂
3 ¹ A ₂	3d _{xx} -R	forbidden	6.69		6.85	forbidden	3 ¹ A ₂	6.81	forbidden	3 ¹ A ₂
2 ¹ B ₁	3d _{yz} -R	0.0003	6.71		6.87	0.0003	2 ¹ B ₁	6.86	0.0000	2 ¹ B ₁
2 ¹ A ₁	V~mix. ^f	0.0009	6.76		6.31	0.0003	2 ¹ A ₁	6.42	0.0039	2 ¹ A ₁
4 ¹ A ₂	3d _{yy-zz} -R	forbidden	6.77		6.91	forbidden	4 ¹ A ₂	6.88	forbidden	4 ¹ A ₂
3 ¹ B ₂	3d _{xz} -R	0.0005	6.93	6.80	6.86	0.0101	3 ¹ B ₂	6.91	0.0105	3 ¹ B ₂
3 ¹ A ₁	3d _{xy} -R	0.0007	6.96		6.93	0.0208	3 ¹ A ₁	6.89	0.0003	3 ¹ A ₁
5 ¹ A ₂	4s-R	forbidden	7.07							
3 ¹ B ₁	4p _y -R	0.0072	7.24	7.05						
4 ¹ B ₂	4p _x -R	0.0078	7.25							
6 ¹ A ₂	4p _z -R	forbidden	7.27							
7 ¹ A ₂	4d _{xx} -R	forbidden	7.42							
4 ¹ B ₁	4d _{yz} -R	0.0002	7.46							
8 ¹ A ₂	4d _{yy-zz} -R	forbidden	7.48							
5 ¹ B ₂	4d _{xz} -R	0.0001	7.55							
4 ¹ A ₁	4d _{xy} -R	0.0010	7.57							
9 ¹ A ₂	5s-R	forbidden	7.60							
5 ¹ B ₁	5p _y -R	0.0083	7.73	7.61						
6 ¹ B ₂	5p _x -R	0.0054	7.74							
6 ¹ B ₁	3s-R	0.0095	7.93	7.84				7.80	0.0217	3 ¹ B ₁
7 ¹ B ₁	5d _{yz} -R	0.0000	8.08	8.03	7.95	0.0252	3 ¹ B ₁			
5 ¹ A ₁	V~mix. ^f	0.5377	8.17	7.88,8.06	7.89	0.4415	4 ¹ A ₁	8.11	0.5020	4 ¹ A ₁
7 ¹ B ₂	5d _{xz} -R	0.0000	8.23	8.19						
6 ¹ A ₁	5d _{xy} -R	0.0966	8.27	8.28						
8 ¹ B ₁	3p _z '-R	0.0122	8.47	8.36						
7 ¹ A ₁	3p _x '-R	0.0216	8.56							
9 ¹ B ₁	4p _z '-R	0.0038	8.59							
8 ¹ B ₂	π - σ^*	0.0248	9.07							
8 ¹ A ₁	3d _{xz} '-R	0.0290	9.23	9.2						
9 ¹ B ₂	3d _{xy} '-R	0.0053	9.25							
10 ¹ B ₂	4d _{xy} '-R	0.0104	9.43							
9 ¹ A ₁	4p _x '-R	0.0016	9.54							
1 ³ B ₂	V ^f	0.0	3.40	3.10	3.15					
1 ³ A ₁	V ^f	0.0	5.18	>4.7	4.90					
1 ³ A ₂	3s-R	0.0	5.56		5.63					
1 ³ B ₂	3p _y -R	0.0	6.19		6.2					

^aV and R denote, respectively, valence and Rydberg excited state.^bTransition energy.^cReferences 4 and 26.^dReference 8.^eReference 9.^fSee text.

CASPT2 except that Rydberg mixing was found. The reason for this difference between our result and other theoretical results is unclear. This excited state is not observed experimentally because of the weak oscillator strength, so it is difficult to make a conclusion experimentally. However, in the case of thiophene, which will be discussed in a later paper,²⁸ the 2 ¹A₁ state has been well reproduced by the SAC-CI SD-R method with regard to both excitation energy and oscillator strength, in comparison with the experimental data in the VUV spectrum.

B. Singlet Rydberg states

The lowest singlet Rydberg transition located at 5.63 eV with a rather weak feature was assigned experimentally by

Robin²⁶ and McDiarmid and Sabljic²⁷ to the symmetry-forbidden 1 ¹A₂(1a₂→3s) state. In the present calculations, the 1 ¹A₂ state was calculated at 5.58 eV. In the two-photon resonant multiphoton ionization (2RMPI) spectrum of Sabljic *et al.*²⁹ the group of transition which occurred in the region 6.0–6.4 eV was assigned to the 1a₂→3p Rydberg series, 6.25(3p_y), 6.26(3p_z), and 6.31(3p_x) eV. The present results for these states are 6.17 eV (1 ¹B₁:3p_y), 6.21 eV (2 ¹A₂:3p_x), and 6.22 eV (2 ¹B₂:3p_z) in a slightly different order. Note that the splitting of the 3p series of Rydberg is very small in both experimental and theoretical results.

Frueholz *et al.*⁴ also observed some apparent Rydberg transitions at 6.80, 7.05, 7.61, 7.84, 8.03, 8.19, 9.2 eV and a higher energy region, but these states were not experimen-

FIG. 2. EEL spectrum and calculated excitation energies and oscillator strengths of cyclopentadiene.

tally assigned. Our current SAC-CI calculations gave a large number of Rydberg excitations, as shown in Tables III and IV. The 3^1B_2 state $1a_2 \rightarrow 3d_{xz}$ transition was calculated at 6.93 eV with an intensity of 0.0005, the 3^1A_1 state $1a_2 \rightarrow 3d_{xy}$ was at 6.96 eV with an intensity of 0.0007, and they were both assigned to the 6.80 eV region in the spectrum. The 3^1B_1 state $1a_2 \rightarrow 4p_y$ transition at 7.24 eV with an intensity of 0.0072 and the 4^1B_2 state $1a_2 \rightarrow 4p_x$ at 7.25 eV with an intensity of 0.0078 were both assigned to the 7.05 eV region of the spectrum. In addition, in the higher energy region of 7.07–7.74 eV we obtained many Rydberg states as seen from Table IV, but these states were not reported in the CASPT2 and MRMP studies. We assigned the transition observed at the 7.84 eV region to the 6^1B_1 $2b_1 \rightarrow 3s$ state (denoted as $3s'$ to distinguish the $1a_2 \rightarrow 3s$ series, for convenience the convention is reserved in thereafter) that was calculated at 7.93 eV with an intensity of 0.0095, where the space symmetry is the same as in the MRMP assignment, 3^1B_1 . The transition observed in the 8.03 eV region is assigned to the $7^1B_1(5d_{yz})$ state calculated at 8.08 eV. In the even higher energy region, our assignments are shown in Table IV. Both CASPT2 and MRMP assigned the peaks at 6.80 and 7.05 eV to the $1a_2 \rightarrow 3d$ transitions, but CASPT2 assigned the 8.03 eV peak to the $2b_1 \rightarrow 3s$ transition, while MRMP assigned this transition to the peak at 7.84 eV; these assignments were not consistent.

C. Triplet excited states

The lowest-energy transition of CP observed by electron impact spectroscopy⁴ had an effective Franck–Condon region extending from 2.5 to 3.9 eV with an intensity maxi-

FIG. 3. PES spectrum and calculated ionization energies and monopole intensities of cyclopentadiene.

mum at 3.10 eV, which was similar to that obtained by Evans³⁰ using the oxygen perturbation technique. Frueholz *et al.*⁴ experimentally assigned this transition to the singlet–triplet excitation, $N \rightarrow T_1(\bar{X}^1A_1 \rightarrow ^3B_2)$. Our current calculations gave the 1^3B_2 state at 3.40 eV and its second moment was 69.42, which clearly showed it as a valence excited state. Its main configuration is $^3\pi-\pi^*$ excitation of $18 \rightarrow 48$.

The second triplet state has not yet been observed by spectroscopic studies. Frueholz *et al.*⁴ suggested that it was hidden behind the singlet valence excitation at 5.2 eV and estimated the excitation energy of the second triplet state to be above 4.7 eV. Our current results predict this state to be 1^3A_1 at 5.18 eV with a second moment of 70.18, which means it is a valence-excited state. Its main configuration is a mixture of the $^3\pi-\pi^*$ excitation of $17 \rightarrow 48$ and $18 \rightarrow 54$. CASPT2 also computed these two valence excited states at 3.15 (1^3B_2) and 4.90 eV (1^3A_1). In addition, the lowest 3A_2 and 3B_1 states have also been calculated at 5.56 and 6.19 eV, respectively, and these are Rydberg excitations in nature as seen from Table IV.

D. Ionized states

The electronic structure of the ground state of CP can be described with 18 molecular orbitals (MOs). They are given in Table II together with their natures and symmetries.

A photoelectron spectrum (PES) of CP has been measured by Derrick *et al.*⁵ up to 23 eV, containing both outer- and inner-valence regions: it is illustrated in the upper side of Fig. 3. Table V shows the SAC-CI general-R theoretical binding energies, monopole intensities, main configurations, and assignments of the ionization peaks in PES. Table V also shows the SAC-CI SD-R results and Koopmans values. The SAC-CI general-R method gave satisfactory results, which were better than the SD-R results. The Koopmans values deviate strongly from the experimental values except for those of the two lowest states. In Fig. 3, the experimental

TABLE V. Calculated ionization potentials, Koopmans energies, and experimental values of PES for cyclopentadiene.

State	SAC-CI General-R			PES (Ref. 5)	SAC-CI SD-R		Koopmans energies
	Main configuration (MC)	IP(eV)	Intensity ^a	IP(eV)	IP(eV)	MC	
1 ² A ₂	0.95 ^b (18→∞)	8.39	0.90	8.566	8.30	0.98(18→∞)	8.37 (1a ₂)
1 ² B ₁	0.93(17→∞)	10.80	0.87	10.724	10.53	0.97(17→∞)	11.24 (2b ₁)
1 ² B ₂	0.93(16→∞)	12.23	0.87	~12.2	11.97	0.96(16→∞)	13.62 (6b ₂)
1 ² A ₁	0.93(15→∞)	12.29	0.87	~12.6	11.95	0.96(15→∞)	13.79 (9a ₁)
2 ² A ₁	0.93(14→∞)	12.79	0.87	13.2	12.50	0.96(14→∞)	14.39 (8a ₁)
2 ² B ₂	0.93(13→∞)	13.58	0.86	13.8	13.33	0.96(13→∞)	15.29 (5b ₂)
2 ² B ₁	0.39(12→∞),0.99(18→∞;18→48)	13.78	0.18		15.91	0.40(12→∞) ^g	
3 ² B ₁	0.85(12→∞),0.46(18→∞;18→48)	15.26	0.72	14.8			16.07 (1b ₁)
3 ² A ₁	0.87(11→∞),0.23(18→∞;16→48)	16.67	0.76	16.4	16.57	0.93(11→∞)	18.94 (7a ₁)
3 ² B ₂	0.77(10→∞),0.41(18→∞;15→48)	17.28	0.62	17.5	17.23	0.93(10→∞)	19.81 (4b ₂)
4 ² A ₁	0.13(11→∞),0.87(18→∞;16→48)	17.43	0.02				
4 ² B ₂	0.40(10→∞),0.80(18→∞;15→48)	17.44	0.14				
5 ² A ₁	0.85(9→∞),0.20(18→∞;16→48)	17.98	0.76	18.4	17.89	0.93(9→∞)	20.57 (6a ₁)
5 ² B ₂	0.87(18→∞;14→48)+0.39(14→∞; 18→48)−0.31(18→∞;14→63)	18.07	0.03				
11 ² A ₁	0.65(16→∞;18→48)+0.21(16→∞; 18→63)−0.20(15→∞;18→67)	19.70	0.07				
13 ² B ₂	−0.72(14→∞;14→48)+0.24(14→∞; 18→63)+0.23(13→∞;17→48)	20.36	0.04				
15 ² A ₁	−0.60(13→∞;18→48)+0.31(14→∞; 17→63)+0.29(16→∞;18→48)	20.89	0.05				
18 ² B ₂	−0.46(16→∞;17→48)+0.35(16→∞; 18→67)−0.27(15→∞;18→48)	21.47	0.06				
21 ² B ₂	0.36(7→∞),0.37(18→∞;18→58) ^c	21.86	0.13	22.0			26.00 (5a ₁)
24 ² B ₂	−0.31(7→∞),0.58(18→∞;18→31),0.43 (18→∞;18→51)	22.51	0.10				
25 ² B ₂	0.39(7→∞),0.60(18→∞;18→34) ^e	22.65	0.15				
29 ² A ₁	0.63(8→∞),0.29(18→∞;18→33) ^d	22.70	0.41	22.3			26.30 (3b ₂)
28 ¹ B ₂	0.32(7→∞),0.37(18→∞;18→34) ^f	22.84	0.10				
29 ² B ₂	0.27(7→∞),0.40(18→∞;18→34),0.27 (18→∞;18→25)	22.87	0.07				

^aMonopole intensity including the final state and initial state configuration interactions. The state whose intensity larger than 0.01 is given.

^bThe main configuration whose absolute value of the coefficient larger than 0.2 is given.

^cAlso includes configurations: 0.31(18→∞;18→66).

^dAlso includes configurations: 0.25(18→∞;18→59).

^eAlso includes configurations: 0.38(18→∞;18→56).

^fAlso includes configurations: 0.25(17→∞;18→29).

^gAlso includes configurations: 1.00(18→48+18→∞) and 0.38(18→∞;18→63).

PES and the present theoretical spectrum calculated by the SAC-CI general-R method are compared.

The peaks at 8.566 and 10.724 eV were experimentally assigned⁵ to the outer two π MOs 1a₂ and 2b₁. Our present SAC-CI general-R calculations computed these two states at 8.39 and 10.80 eV, respectively. It is clear from the main configurations (shown in Table IV) of the two ionized states 1 ²A₂ and 1 ²B₁ that they are dominated by one-electron process, confirming the previously experimental assignment. The next sequential peaks at ~12.2, ~12.6, 13.2, and 13.8 eV in PES were assigned to the 1 ²B₂, 1 ²A₁, 2 ²A₁, and 2 ²B₂ states calculated at 12.23, 12.29, 12.79, and 13.58 eV, respectively, by the SAC-CI general-R method. These ionizations are also dominated by one-electron process, as seen from their main configurations. Note that the experimental peaks at ~12.2 and ~12.6 eV are overlapping ones with complicated vibrational structure.

In the energy region higher than 14.3 eV, other six sequential ionization peaks were observed at 14.8, 16.4, 17.5, 18.4, 22.0, and 22.3 eV. Our present SAC-CI general-R results calculations showed that these six ionizations were

more or less characterized by multielectron processes. The peak at 14.8 eV was assigned to the 3 ²B₁ state calculated at 15.26 eV with the main configurations that are a mixture of the single- and double-electron processes, 0.85(12→∞) and 0.46(18→48,18→∞), respectively. The peak at 16.4 eV was assigned to the 3 ²A₁ state calculated at 16.67 eV with the main configurations of 0.87(11→∞), 0.11(9→∞), and 0.23(18→48,16→∞). The peak at 17.5 eV was assigned to the 3 ²B₂ state calculated at 17.28 eV with the main configurations of 0.77(10→∞) and 0.41(15→48,18→∞). The peak at 18.4 eV was assigned to the 5 ²A₁ state calculated at 17.98 eV with the main configurations of 0.87(9→∞), 0.17(11→∞), and 0.19(18→48,16→∞). The peaks at about 22.0 and 22.3 eV lie in a broad peak in PES.⁵ Our present SAC-CI general-R calculations gave many ionized states with the natures of the multielectron process in this region. The 29 ²A₁ state calculated at 22.70 eV with a monopole intensity of 0.41 has the main configurations of 0.63(8→∞), 0.29(18→33,18→∞), and 0.25(18→59,18→∞). The 21 ²B₂ state calculated at 21.86 eV with a monopole intensity of 0.13 has the main configurations of 0.36(7→∞), 0.37(18→58,18

$\rightarrow\infty$), and $0.31(18\rightarrow 66,18\rightarrow\infty)$. The 25^2B_2 state calculated at 22.51 eV with a monopole intensity of 0.10 has the main configurations of $0.31(7\rightarrow\infty)$, $0.58(18\rightarrow 31,18\rightarrow\infty)$, and $0.43(18\rightarrow 51,18\rightarrow\infty)$. The 26^2B_2 state calculated at 22.65 eV with a monopole intensity of 0.15 has the main configurations of $0.39(7\rightarrow\infty)$, $0.60(18\rightarrow 34,18\rightarrow\infty)$, and $0.38(18\rightarrow 56,18\rightarrow\infty)$. The 28^2B_2 state calculated at 22.84 eV with a monopole intensity of 0.10 has the main configurations of $0.32(7\rightarrow\infty)$, $0.37(18\rightarrow 34,18\rightarrow\infty)$, and $0.25(18\rightarrow 29,17\rightarrow\infty)$. Thus our results show that the peaks at around 22.0 eV consist of a series of correlation peaks.

Note that the present SAC-CI SD-R results (shown in Table V) are generally in agreement with the experimental PES data except for the three ionization peaks at 14.8, 22.0, and 22.3 eV, which are dominated by multiple-electron processes. Since the SD-R calculations did not include higher-order excitation operators, the ionization potentials of the inner-valence region were not accurately computed. A comparison of the general-R results with the SD-R results obviously shows that the inclusion of a small number of but important higher-order excitation operators by the SAC-CI general-R method remarkably improves the description of the binding energies in the inner-valence region.

IV. CONCLUSIONS

In the present study, singlet and triplet excited states and ionized states of cyclopentadiene (CP) were reinvestigated by the SAC/SAC-CI method with an adequately extended basis set and a large active orbital space. The 40 low-lying singlet and triplet excited states were calculated and a reliable assignment for the UV and EEL spectra up to 8 eV was made using the calculated excitation energies and oscillator strengths. Valence and Rydberg excited states were discussed separately in detail. Furthermore, the SAC-CI general-R method with the combined use of the exponential generation and perturbation selection schemes for including higher-order excitation operators was applied to the ionized states of CP up to 23 eV. A comparison of the experimental PES spectrum and our present calculations shows a good agreement in terms of the binding energies and the peak intensities in the both outer- and inner-valence regions.

ACKNOWLEDGMENTS

We are grateful to Dr. J. Meller and Mr. K. Toyota for their helpful discussions on this subject. This research was

supported by a Grant-in-Aid for Scientific Research from the Ministry of Education, Science, Culture, and Sports.

- ¹G. Scheibe and H. Grieneisen, Z. Phys. Chem. Abt. B **25**, 52 (1934).
- ²L. W. Pickett, E. Paddock, and E. Sackter, J. Am. Chem. Soc. **63**, 1073 (1941).
- ³W. C. Price and A. D. Walsh, Proc. R. Soc. London, Ser. A **179**, 201 (1941).
- ⁴R. P. Frueholz, W. M. Flicker, O. A. Mosher, and A. Kuppermann, J. Chem. Phys. **70**, 2003 (1979).
- ⁵P. J. Derrick, L. Asbrink, O. Edqvist, B.-Ö. Jonsson, and E. Lindholm, Int. J. Mass Spectrom. Ion Phys. **6**, 203 (1971).
- ⁶H. Nakatsuji, O. Kitao, and T. Yonezawa, J. Chem. Phys. **83**, 723 (1985).
- ⁷V. Galasso, Chem. Phys. **153**, 13 (1991).
- ⁸L. Serrano-Andres, M. Merchán, I. Nebot-Gil, B. O. Roos, and M. Fulscher, J. Am. Chem. Soc. **115**, 6184 (1993).
- ⁹H. Nakano, T. Tsuneda, T. Hashimoto, and K. Hirao, J. Chem. Phys. **104**, 2312 (1996).
- ¹⁰H. Nakatsuji, Chem. Phys. Lett. **177**, 331 (1991).
- ¹¹H. Nakatsuji and K. Hirao, J. Chem. Phys. **68**, 2053 (1978).
- ¹²H. Nakatsuji, Chem. Phys. Lett. **59**, 362 (1978).
- ¹³H. Nakatsuji, Chem. Phys. Lett. **67**, 329 (1979).
- ¹⁴H. Nakatsuji, in *Computational Chemistry: Reviews of Current Trends*, edited by J. Leszczynski (World Scientific, Singapore, 1997), Vol. 2.
- ¹⁵K. Hirao and H. Nakatsuji, Chem. Phys. Lett. **79**, 292 (1981).
- ¹⁶H. Nakatsuji, Chem. Phys. **75**, 425 (1983).
- ¹⁷H. Nakatsuji, J. Chem. Phys. **83**, 5743 (1985).
- ¹⁸H. Nakatsuji, M. Hada, M. Ehara, J. Hasegawa, T. Nakajima, H. Nakai, O. Kitao, and K. Toyota, SAC/SAC-CI program system (SAC-CI96) for calculating ground, excited, ionized, and electron attached states and singlet to septet spin multiplicities.
- ¹⁹LANDOLT-BORNSTEIN, *Numerical Data and Functional Relationships in Science and Technology, New Series, Group II: Atomic and Molecular Physics*, edited by J. H. Callomon, E. Hirota, K. Kuchitsu, W. J. Lafferty, A. G. Maki, and C. S. Pote (Springer-Verlag, Berlin-Heidelberg, 1976), Vol. 7.
- ²⁰R. A. Kendall, T. H. Dunning, Jr., and R. J. Harrison, J. Chem. Phys. **96**, 6796 (1992).
- ²¹T. H. Dunning Jr., J. Chem. Phys. **90**, 1007 (1989).
- ²²K. Kaufmann, W. Baumeister, and M. Jungen, J. Phys. B **22**, 2223 (1989).
- ²³A. B. Trofimov and J. Schirmer, Chem. Phys. **214**, 153 (1997).
- ²⁴GAUSSIAN 98, Revision A.1, M. J. Frisch, G. W. Trucks, H. B. Schlegel et al., Gaussian, Inc., Pittsburgh, Pennsylvania, 1998.
- ²⁵R. S. Mulliken, J. Chem. Phys. **7**, 121 (1939).
- ²⁶M. B. Robin, *Higher Excited States of Polyatomic Molecules* (Academic, New York, 1975), Vol. II, pp. 166–176.
- ²⁷R. McDiarmid and A. Sabljic, J. Chem. Phys. **83**, 2147 (1985).
- ²⁸H. Nakatsuji, J. Wan, and M. Hada (unpublished).
- ²⁹A. Sabljic, R. McDiarmid, and A. Gedanken, J. Phys. Chem. **96**, 2442 (1992).
- ³⁰D. F. Evans, J. Chem. Soc., 1735 (1960).