TITLE:

Fine theoretical spectroscopy using
symmetry adapted cluster-configuration
interaction general-R method: Outer- and
inner-valence ionization spectra of CS₂ and
OCS

AUTHOR(S):

Ehara, M; Ishida, M; Nakatsuji, H

CITATION:

Ehara, M ...[et al]. Fine theoretical spectroscopy using symmetry adapted cluster-configuration interaction general-R method: Outer- and inner-valence ionization spectra of CS₂ and OCS. JOURNAL OF CHEMICAL PHYSICS 2002, 117(7): 3248-3255

ISSUE DATE:

2002-08-15

URL:

<http://hdl.handle.net/2433/50184>

RIGHT:

Copyright 2002 American Institute of Physics. This article may be downloaded for personal use only. Any other use requires prior permission of the author and the American Institute of Physics.

Fine theoretical spectroscopy using symmetry adapted cluster-configuration interaction general-*R* method: Outer- and inner-valence ionization spectra of CS₂ and OCS

Masahiro Ehara, Mayumi Ishida, and Hiroshi Nakatsuji^{a)}

Department of Synthetic Chemistry and Biological Chemistry, Graduate School of Engineering, Kyoto University, Kyoto 606-8501, Japan

(Received 25 January 2002; accepted 20 May 2002)

Fine theoretical spectroscopy has been presented by the SAC-CI (symmetry adapted cluster-configuration interaction) general-*R* method for the outer- and inner-valence ionization spectra of CS₂ and OCS. The SAC-CI general-*R* method simulated the experimental spectra quite accurately and the detailed assignments of the satellite peaks were given. For CS₂, four outer-valence satellites ²Π_u states were calculated, one of which was attributed to the recently observed peak (1'). Numerous ²Σ_u⁺ and ²Σ_g⁺ satellite peaks were obtained in the inner-valence region and some of them were dominantly described by triple electron processes; the quadruple *R*-operators were found to be important for describing these states in the general-*R* method. For OCS, the relative position of the main peaks was correctly reproduced and the higher *R*-operators were found to be important for the ordering of *A* and *B* states. In the energy region of 24–36 eV, continuous spectra of numerous ²Σ⁺ satellites were obtained, which reproduced the feature of the photoelectron spectrum. © 2002 American Institute of Physics. [DOI: 10.1063/1.1492798]

I. INTRODUCTION

Multiple-electron processes observed in ionization spectra have attracted considerable attention since they directly reflect the electron correlations of the molecules. They are called “satellites” and usually appear in the inner-valence region. But, for the strongly correlated system, they even appear in the outer-valence region. Recently, fine analyses of these peaks on peak positions, intensities, and ionization characters have become possible due to the cooperative interplays between qualified theoretical and experimental works. The developments in the high-resolution spectroscopies are remarkable as seen for synchrotron radiation photoelectron spectroscopy (SRPES), x-ray PES, electron momentum spectroscopy (EMS), and Penning ionization electron spectroscopy (PIES).

Among the extensive works on the satellite peaks, special attention has been paid to the correlation effects in the valence ionized states of CS₂.^{1–9} Allan *et al.*¹ observed both the outer- and inner-valence region using MgKα PES. The He II PES^{2,3} was recorded and the theoretical assignment was given by the 2ph-TDA Green's function method.³ The dipole (*e,2e*)⁴ and binary (*e,2e*)⁵ electron impact spectroscopy were also applied. Later, the details of the valence satellites were studied by the high-resolution SRPES using the photon energy dependence of the partial cross sections and asymmetry parameter and also with the help of theoretical spectrum of CIPSI method.⁷ In particular, it was pointed out that some of the satellite peaks were described by the three-electron processes. Recently, fine structure of the satellite peaks was analyzed by the He I PES, He II PES and SRPES combined with the algebraic diagrammatic construction (ADC) calculation.⁸ However, there are still some discrepan-

cies in the assignments of the satellite peaks. It should also be noted that the precise positions of the main peaks were determined by the high-resolution He II PES¹⁰ and the double ionization spectrum was investigated.¹¹

The valence ionized states of OCS have also been intensively studied by various spectroscopies, for examples, He I PES,¹⁰ He II PES,² MgKα PES,¹ and dipole (*e,2e*) for impact energies up to 40 eV.^{12–14} The latest experimental work is due to the SRPES¹⁵ with photon energies up to 110 eV. However, theoretical studies on the detailed analysis of the satellites have been very limited for this compound; only the SAC-CI (symmetry adapted cluster-configuration interaction) method⁹ and 2ph-TDA method¹² were applied to the inner-valence region. It was also criticized that theoretical spectra did not reproduce the experimental PES; the calculated inner-valence satellite peaks clustered separately against the experimental observation. Thus, it is worth presenting the accurate theoretical spectra and detailed assignments for the correlation peaks of these molecules.

In the series of the studies, we have investigated the valence ionization spectra of various molecules using the SAC-CI method. The SAC/SAC-CI method^{16–21} has been successfully applied to a number of molecular spectroscopies including ionization spectra.^{20–27} In particular, the SAC-CI general-*R* method^{28–31} has been shown to be a powerful tool for describing the multiple electron processes in high accuracy and in studying the large numbers of states appearing in the ionization spectra.^{32–38} Recently, we applied the general-*R* method to both the outer- and inner-valence ionization spectra of Group V and VI hydrides and gave the detailed characterizations of the satellite peaks.^{36,37} In this work, we investigate the valence ionization spectra of CS₂ and OCS by the SAC-CI general-*R* method aiming at the detailed and quantitative assignments of the satellite peaks appearing in the outer- and inner-valence regions.

^{a)} Author to whom correspondence should be addressed. Electronic mail: hiroshi@sbchem.kyoto-u.ac.jp

II. COMPUTATIONAL DETAILS

The basis set was selected as flexible as possible to describe the electron correlations of shake-up states, namely valence triple zeta (VTZ) of Ahlrichs;³⁹ (10s6p)/[6s3p] for C and O and (12s9p)/[7s5p] Gaussian type orbitals (GTOs) for S augmented with two polarization *d*-functions of $\zeta_d=2.314$, 0.645 for O, $\zeta_d=1.097$, 0.318 for C, and $\zeta_d=0.819$, 0.269 for S.^{40,41} The present basis set was examined for the low-lying valence ionized states with the additional polarization function and diffuse functions, and was confirmed to be sufficient as shown in the Appendix. The resultant SCF dimensions were 89 and 82 for CS₂ and OCS, respectively. The vertical ionization process was studied in this work and, therefore, molecular geometries were fixed to the experimental values;⁴² namely $R_{CS}=1.55545$ Å for CS₂ and $R_{OC}=1.16$ Å and $R_{CS}=1.56$ Å for OCS.

The valence ionization spectra of CS₂ and OCS were calculated by the SAC-CI general-*R* method in both outer- and inner-valence regions up to around double ionization thresholds. In the preliminary calculations, several shake-up states in the inner-valence region of CS₂ were found to be described by triples with considerable intensity; this feature was also pointed out in the CIPSI work.⁷ For producing the reliable spectrum in this energy region, we included *R*-operators up to quadruples in the general-*R* calculation and utilized the exponentially generated CI (EGCI) method^{29–31} for the reference-CI. In the case of OCS, most of the shake-up states were dominantly described by the doubles until around the double ionization threshold and, therefore, *R*-operators were included up to triples. The 1s orbitals of C and O and 1s, 2s, and 2p orbitals of S were kept as frozen core and all the other MOs were included in the active space; the active space of the SAC-CI consisted of 8 occupied and 70 unoccupied MOs for CS₂ and 8 occupied and 67 unoccupied MOs for OCS.

In order to reduce the computational effort, perturbation selection²² was performed in the state-selection scheme. The thresholds were determined so as to achieve the calculations including the sufficient order of *R*-operators yet with appropriate computational cost. For CS₂, the threshold of the linked terms for the ground state was set to $\lambda_g=5 \times 10^{-6}$ and the unlinked terms were adopted as the products of the important linked terms whose SDCI coefficients were larger than 0.005. For ionized state, the threshold of the double linked term was set to $\lambda_e=1 \times 10^{-7}$ and those of the triples and quadruples were $\lambda_e=1 \times 10^{-6}$ and 5×10^{-6} , respectively. The reference-EGCI calculations were performed to determine the primary states using small active space; triples were generated by the exponential-generation algorithm.^{29–31} The threshold of the exponential generation was set to $\lambda_{AA}=0.1$. For the main peaks of CS₂, better thresholds were used. For OCS, better thresholds were adopted, namely, $\lambda_g=1 \times 10^{-7}$ for SAC and $\lambda_e=1 \times 10^{-7}$ and 1×10^{-6} for doubles and triples of SAC-CI, respectively. The thresholds of the CI coefficients for calculating the unlinked operators in the SAC-CI method are 0.01 and 1×10^{-8} for the *R* and *S* operators, respectively.

Ionization cross-sections were calculated using the monopole approximation^{43,44} to estimate the relative intensi-

TABLE I. The reference-EGCI and SAC-CI general-*R* dimensions for the ionized states of CS₂ and OCS.

State	Total	Singles	Doubles	Triples	Quadruples
CS ₂					
Reference-EGCI dimension					
$2\Sigma_g^+$	5617	2	228	5387	...
$2\Pi_g$	5679	1	200	5478	...
$2\Sigma_u^+$	5650	2	228	5420	...
$2\Pi_u$	5586	1	200	5385	...
SAC-CI dimension after perturbation selection					
$2\Sigma_g^+$	86 399	2	608	42 523	43 266
$2\Pi_g$	82 227	1	518	35 630	46 078
$2\Sigma_u^+$	84 319	2	601	42 130	41 586
$2\Pi_u$	83 758	1	515	35 730	47 512
OCS					
SAC-CI dimension after perturbation selection					
$2\Sigma^+$	97 117	4	1164	95 949	...
2Π	83 562	2	986	82 574	...

ties of the peaks. Both initial- and final-state correlation effects were included.

The SAC/SAC-CI calculations were executed using the SAC96 program system,⁴⁵ which has been incorporated into the development version of the Gaussian suite of programs.⁴⁶

III. RESULTS AND DISCUSSION

A. CS₂

The outer- and inner-valence regions of the ionization spectra of CS₂ up to c.a. 28 eV were studied by the SAC-CI general-*R* method. The valence electronic configuration of CS₂ is described by

$$(\text{core})^{22}(5\sigma_g)^2(4\sigma_u)^2(6\sigma_g)^2(5\sigma_u)^2(2\pi_u)^4(2\pi_g)^4.$$

As noted in the previous section, several states in the inner-valence region were found to be substantially dominated by triples; namely ionization accompanied by two-electron ex-

FIG. 1. Valence ionization spectra of CS₂ by (a) SRPES (Ref. 8) and (b) SAC-CI general-*R* method.

TABLE II. Ionization potential (I.P.) (in eV), monopole intensity and main configuration of the valence ionized states of CS_2 .

Experimental I.P.						SAC-CI general-R			
SRPES ^a	PES ^b	PES ^c	Hell ^d	($e, 2e$) ^e		State	I.P. (eV)	Intensity	Main configurations ($ C > 0.3$)
No.									
X	10.00	10.11	...	...	...	$2\Pi_g$	9.70	0.841	$0.89(2\pi_g^{-1})$
A	12.90	12.89	...	...	...	$2\Pi_u$	12.76	0.693	$0.80(2\pi_u^{-1}) - 0.33(2\pi_g^{-2}3\pi_u)$
B	14.46	14.49	...	...	...	$2\Sigma_g^+$	14.38	0.804	$0.86(5\sigma_u^{-1})$
1	14.1	14.09	14.1	14.06	...	$2\Pi_u$	14.75	0.027	$0.52(2\pi_g^{-2}3\pi_u) + 0.44(2\pi_g^{-2}3\pi_u) - 0.39(2\pi_g^{-2}3\pi_u)$
						$2\Pi_u$	15.19	0.005	$0.78(2\pi_g^{-2}3\pi_u) + 0.56(2\pi_g^{-2}3\pi_u) + 0.39(2\pi_g^{-2}3\pi_u)$
1'	...	15.2	...	...	...	$2\Pi_u$	15.69	0.009	$0.57(2\pi_g^{-2}3\pi_u) - 0.56(2\pi_g^{-2}3\pi_u) - 0.48(2\pi_g^{-2}3\pi_u) + 0.42(2\pi_g^{-2}3\pi_u)$
C	16.15	16.20	...	...	...	$2\Sigma_g^+$	16.19	0.705	$0.81(7\sigma_g^{-1})$
2	17.2	17.0	17.1	17.06	...	$2\Pi_u$	18.24	0.115	$0.32(2\pi_u^{-1}) + 0.67(2\pi_g^{-2}3\pi_u) + 0.43(2\pi_g^{-2}3\pi_u) - 0.20(2\pi_g^{-2}3\pi_u^22\pi_u^{-1})$
3	19.15	18.81	18.36	18.97	19.1	$2\Sigma_g^+$	19.20	0.006	$0.48(2\pi_g^{-2}8\sigma_g) + 0.48(2\pi_g^{-2}8\sigma_g) - 0.39(5\sigma_u^{-1}3\pi_u2\pi_g^{-1}) - 0.37(5\sigma_u^{-1}3\pi_u2\pi_g^{-1})$
						$2\Sigma_g^+$	19.74	0.026	$0.54(5\sigma_u^{-1}3\pi_u2\pi_g^{-1}) + 0.53(5\sigma_u^{-1}3\pi_u2\pi_g^{-1}) + 0.44(5\sigma_u^{-1}3\pi_u2\pi_g^{-1})$
4	20.4	20.17	19.91	20.22	20.2	$2\Pi_g$	20.15	0.001	$0.44(2\pi_u^{-1}3\pi_u2\pi_g^{-1}) + 0.35(2\pi_g^{-1}3\pi_u2\pi_u^{-1}) + 0.34(2\pi_u^{-1}3\pi_u2\pi_g^{-1}) + 0.31(2\pi_g^{-1}3\pi_u2\pi_u^{-1}) - 0.60(2\pi_g^{-3}3\pi_u^2)$
						$2\Pi_g$	20.23	0.0002	$0.46(2\pi_g^{-1}3\pi_u2\pi_u^{-1}) + 0.43(2\pi_g^{-1}3\pi_u2\pi_u^{-1}) - 0.42(2\pi_u^{-1}3\pi_u2\pi_g^{-1}) + 0.38(2\pi_u^{-1}3\pi_u2\pi_g^{-1}) + 0.38(2\pi_u^{-1}3\pi_u2\pi_g^{-1}) - 0.36(2\pi_g^{-1}3\pi_u2\pi_u^{-1}) + 0.39(2\pi_g^{-3}3\pi_u^2) + 0.22(2\pi_g^{-3}3\pi_u^2)$
5	20.8	...	20.96	...	...	$2\Sigma_u^+$	20.88	0.005	$0.53(5\sigma_u^{-1}3\pi_u2\pi_u^{-1}) + 0.51(5\sigma_u^{-1}3\pi_u2\pi_u^{-1}) + 0.50(7\sigma_g^{-1}3\pi_u2\pi_g^{-1}) + 0.50(7\sigma_g^{-1}3\pi_u2\pi_g^{-1})$
						$2\Pi_u$	20.97	0.001	$0.39(2\pi_u^{-2}3\pi_u) + 0.38(2\pi_u^{-2}3\pi_u) + 0.92(2\pi_g^{-1}3\pi_u2\pi_u^{-2}) + 0.29(2\pi_g^{-1}3\pi_u2\pi_u^{-2}) + 0.25(2\pi_g^{-1}3\pi_u2\pi_u^{-2})$
						$2\Sigma_u^+$	21.10	0.003	$0.49(2\pi_g^{-1}3\pi_u7\sigma_g^{-1}) - 0.46(2\pi_g^{-1}3\pi_u7\sigma_g^{-1}) - 0.45(2\pi_u^{-2}3\pi_u) + 0.44(7\sigma_g^{-1}3\pi_u2\pi_g^{-1}) - 0.44(5\sigma_u^{-1}3\pi_u2\pi_u^{-1}) - 0.41(7\sigma_g^{-1}3\pi_u2\pi_g^{-1}) + 0.41(2\pi_u^{-1}3\pi_u5\sigma_u^{-1}) + 0.39(5\sigma_u^{-1}3\pi_u2\pi_u^{-1}) - 0.30(2\pi_g^{-2}3\pi_u5\sigma_u^{-1})$
6	21.35	21.25	21.34	21.35	21.3	$2\Sigma_u^+$	21.23	0.170	$0.39(4\sigma_u^{-1}) + 0.56(2\pi_g^{-2}7\sigma_u) - 0.39(2\pi_g^{-1}8\sigma_g2\pi_u^{-1})$
						$2\Sigma_u^+$	21.27	0.039	$0.61(2\pi_g^{-2}7\sigma_u) - 0.37(2\pi_g^{-1}8\sigma_g2\pi_u^{-1})$
						$2\Pi_g$	21.35	0.004	$0.38(2\pi_u^{-1}3\pi_u2\pi_g^{-1}) + 0.37(2\pi_g^{-1}3\pi_u2\pi_u^{-1}) + 0.48(2\pi_g^{-3}3\pi_u^2) + 0.38(2\pi_g^{-3}3\pi_u^2)$
						$2\Pi_g$	21.93	0.002	$0.44(2\pi_g^{-1}3\pi_u2\pi_u^{-1}) + 0.36(2\pi_u^{-1}3\pi_u2\pi_g^{-1}) + 0.66(2\pi_g^{-3}3\pi_u^2) + 0.22(2\pi_g^{-3}3\pi_u^2)$
7	21.6	21.6	21.76	21.56	...	$2\Sigma_u^+$	22.27	0.052	$0.21(4\sigma_u^{-1}) - 0.50(2\pi_g^{-1}8\sigma_g2\pi_u^{-1})$
8	22.4	22.4	22.69	...	...	$2\Sigma_u^+$	22.77	0.180	$0.40(4\sigma_u^{-1}) - 0.34(2\pi_g^{-2}7\sigma_u) - 0.33(5\sigma_u^{-1}3\pi_u2\pi_u^{-1}) - 0.32(2\pi_g^{-2}7\sigma_u) - 0.21(2\pi_g^{-2}3\pi_u5\sigma_u^{-1})$
						$2\Sigma_g^+$	23.33	0.010	$0.60(2\pi_g^{-2}9\sigma_g) + 0.60(2\pi_g^{-2}9\sigma_g)$
9	23.3	23.0	23.05	...	...	$2\Sigma_g^+$	23.60	0.077	$0.53(2\pi_g^{-1}3\pi_u5\sigma_u^{-1}) + 0.53(2\pi_g^{-1}3\pi_u5\sigma_u^{-1})$
						$2\Pi_u$	23.72	0.010	$0.39(2\pi_g^{-1}3\pi_u2\pi_u^{-1}) + 0.45(2\pi_g^{-3}3\pi_u^2) + 0.43(2\pi_g^{-3}3\pi_u^2)$
						$2\Sigma_u^+$	23.72	0.035	$0.30(5\sigma_u^{-1}3\pi_u2\pi_u^{-1}) + 1.00(2\pi_g^{-2}3\pi_u5\sigma_u^{-1}) + 0.33(2\pi_g^{-2}3\pi_u5\sigma_u^{-1}) + 0.32(2\pi_g^{-2}3\pi_u5\sigma_u^{-1}) + 0.30(2\pi_g^{-2}3\pi_u5\sigma_u^{-1})$
						$2\Pi_u$	24.17	0.008	$0.69(2\pi_g^{-2}3\pi_u) + 0.34(2\pi_u^{-2}3\pi_u) + 0.25(2\pi_g^{-2}3\pi_u2\pi_u^{-1}) - 0.23(2\pi_g^{-2}3\pi_u2\pi_u^{-1})$
10	24.1	24.0	23.91	23.96	23.9	$2\Sigma_g^+$	24.19	0.270	$0.49(6\sigma_g^{-1}) + 0.47(7\sigma_g^{-1}3\pi_u2\pi_u^{-1}) + 0.45(7\sigma_g^{-1}3\pi_u2\pi_u^{-1}) + 0.33(2\pi_u^{-1}3\pi_u7\sigma_g^{-1}) + 0.33(2\pi_u^{-1}3\pi_u7\sigma_g^{-1})$
						$2\Sigma_u^+$	24.59	0.009	$0.41(5\sigma_u^{-1}3\pi_u2\pi_u^{-1}) - 0.35(7\sigma_g^{-1}3\pi_u2\pi_g^{-1}) - 0.35(2\pi_g^{-2}6\sigma_u) - 0.45(2\pi_g^{-2}3\pi_u5\sigma_u^{-1}) - 0.30(2\pi_g^{-2}3\pi_u5\sigma_u^{-1})$
11	24.6	24.3	...	...	...	$2\Sigma_g^+$	24.77	0.025	$0.46(2\pi_u^{-1}7\sigma_u2\pi_g^{-1}) + 0.34(2\pi_u^{-1}7\sigma_u2\pi_g^{-1})$
						$2\Sigma_u^+$	24.81	0.011	$0.40(2\pi_g^{-2}6\sigma_u) - 0.35(2\pi_g^{-2}6\sigma_u) - 0.32(2\pi_g^{-1}3\pi_u7\sigma_g^{-1})$
12	25.25	25.2	...	...	...	$2\Sigma_u^+$	25.25	0.026	$0.38(2\pi_u^{-1}3\pi_u5\sigma_u^{-1}) - 0.34(5\sigma_u^{-1}3\pi_u2\pi_u^{-1})$
13	26.35	...	...	...	26.2	$2\Sigma_g^+$	25.83	0.101	$0.30(6\sigma_g^{-1}) - 0.35(7\sigma_g^{-1}3\pi_u2\pi_u^{-1}) - 0.32(7\sigma_g^{-1}3\pi_u2\pi_u^{-1})$
						$2\Sigma_g^+$	26.31	0.004	$0.64(2\pi_g^{-2}1\delta_g) + 0.55(2\pi_g^{-2}1\delta_g) + 0.38(2\pi_g^{-2}1\delta_g) - 0.30(2\pi_g^{-2}1\delta_g)$
						$2\Sigma_g^+$	26.46	0.010	$0.41(2\pi_u^{-2}8\sigma_g) + 0.32(2\pi_u^{-2}8\sigma_g) - 0.31(2\pi_g^{-1}7\sigma_u2\pi_u^{-1}) + 0.30(2\pi_g^{-2}1\delta_g)$
						$2\Sigma_g^+$	26.62	0.011	$0.61(2\pi_g^{-2}1\delta_g) + 0.41(2\pi_g^{-2}1\delta_g)$

TABLE II. (Continued.)

Experimental I.P.					SAC-CI general-R				
SRPES ^a	PES ^b	PES ^c	HeII ^d	(e,2e) ^e	State	I.P. (eV)	Intensity	Main configurations ($ C > 0.3$)	
No.									
14	27.4	...	...	...	2 Σ_u^+	27.02	0.073	0.25(4 σ_u^{-1}) + 0.31(2 $\pi_u^{-1}3\pi_u5\sigma_u^{-1}$) + 0.31(2 $\pi_u^{-1}3\pi_u5\sigma_u^{-1}$) - 0.34(2 $\pi_g^{-2}3\pi_u5\sigma_u^{-1}$) + 0.34(2 $\pi_g^{-2}3\pi_u5\sigma_u^{-1}$)	
					2 Σ_u^+	27.17	0.017	0.60(2 $\pi_g^{-2}1\delta_u$) + 0.47(2 $\pi_g^{-2}1\delta_u$) + 0.41(2 $\pi_g^{-2}1\delta_u$)	
					2 Σ_u^+	27.27	0.005	0.46(2 $\pi_u^{-1}6\sigma_u2\pi_g^{-1}$) - 0.33(2 $\pi_g^{-1}6\sigma_u2\pi_u^{-1}$)	
					2 Σ_u^+	27.40	0.013	0.45(2 $\pi_g^{-2}1\delta_g$) - 0.33(2 $\pi_g^{-2}1\delta_g$)	
					2 Σ_u^+	27.94	0.021	0.62(5 $\sigma_u^{-2}8\sigma_g$)	
					2 Σ_u^+	28.15	0.007	0.60(2 $\pi_g^{-2}3\pi_u5\sigma_u^{-1}$) - 0.32(2 $\pi_g^{-1}9\sigma_g2\pi_u^{-1}$) - 0.31(2 $\pi_g^{-1}9\sigma_g2\pi_u^{-1}$) + 0.30(2 $\pi_g^{-2}3\pi_u5\sigma_u^{-1}$)	
15	28.7	...	...	...	2 Σ_u^+	28.22	0.089	0.35(4 $\sigma_u^{-1}3\pi_u2\pi_g^{-1}$) + 0.33(2 $\pi_g^{-2}1\delta_g$)	
					2 Σ_u^+	28.28	0.009	0.42(2 $\pi_g^{-2}1\delta_u$) + 0.41(2 $\pi_g^{-1}1\delta_g2\pi_u^{-1}$) - 0.40(2 $\pi_u^{-1}1\delta_g2\pi_g^{-1}$) - 0.37(2 $\pi_g^{-2}1\delta_u$) + 0.36(2 $\pi_u^{-1}1\delta_g2\pi_g^{-1}$)	
					2 Σ_u^+	28.47	0.007	0.47(2 $\pi_g^{-1}1\delta_g2\pi_u^{-1}$) + 0.45(2 $\pi_g^{-2}1\delta_u$) + 0.38(2 $\pi_u^{-1}1\delta_g2\pi_g^{-1}$) - 0.36(2 $\pi_u^{-1}1\delta_g2\pi_g^{-1}$) - 0.24(2 $\pi_g^{-2}3\pi_u5\sigma_u^{-1}$)	
					2 Σ_u^+	28.67	0.042	0.42(2 $\pi_u^{-1}9\sigma_g2\pi_g^{-1}$) + 0.41(2 $\pi_u^{-1}9\sigma_g2\pi_g^{-1}$) + 0.40(2 $\pi_g^{-1}9\sigma_g2\pi_u^{-1}$) + 0.35(2 $\pi_g^{-1}9\sigma_g2\pi_u^{-1}$)	

^aReference 7.

^bReference 8: experiment was due to the SRPES and He II PES.

^cReference 6.

^dReference 2.

^eReference 5.

citation, therefore the EGCI including up to triples was performed for the reference-CI. The calculational dimensions for the reference-EGCI and resultant SAC-CI general-R were summarized in Table I. The former dimensions were around 5600 and the latter's after perturbation selection were almost 85 000 for each symmetry of $D_{\infty h}$. In Fig. 1, the SAC-CI general-R spectrum convoluting with Gaussian envelope for describing the Frank-Condon width and the resolution of spectrometer was compared with the experimental PES;⁸ the FWHM of Gaussian was adopted as 0.05* ΔE (in eV). In Table II, the results of IPs, monopole intensities, and detailed ionization characters for the valence ionized states, which have large intensity greater than 0.005, were presented with several experimental IPs.^{2,5-8}

Four main peaks in the outer-valence region were dominantly described by the single electron process and the previous SAC-CI SD-R⁹ calculation has already given accurate results. In the present work, their IPs were calculated as 9.70, 12.76, 14.38, and 16.19 eV for $X^2\Pi_g$, $A^2\Pi_u$, $B^2\Sigma_u^+$, and $C^2\Sigma_g^+$ states, respectively, which were compared with the experimental vertical IPs 10.080, 12.698, 14.477, and 16.190 eV by the high-resolution He I PES.¹⁰ The intensities of the $A^2\Pi_u$ and $C^2\Sigma_g^+$ states were considerably distributed to the shake-up states: the monopole intensities were calculated as 0.693 and 0.705 for these states. The relative intensities of the main peaks are almost the same as those of other theoretical calculations by the CIPSI⁷ and simplified ADC(4).⁸

Next, we discuss the spectrum of the shake-up states comparing with the high-resolution spectrum by the SRPES^{7,8} and He II PES.⁸ So far, three satellite peaks were experimentally identified in the outer-valence region of CS₂. These peaks were assigned to $^2\Pi_u$ states^{7,8} and, especially,

the second one (1') was recently observed at the foot of the main peak B.⁸ Accordingly, the present calculation gave four $^2\Pi_u$ states in this energy region at 14.75(1), 15.19, 15.69(1'), and 18.24(2) eV. These outer-valence satellites were calculated in the higher energy region in the previous SAC-CI SD-R study.⁹ They are characterized as (2 $\pi_g^{-2}3\pi_u$) and have the intensity mainly through the final-state correlation interaction with (2 π_u^{-1}) state. The shake-up states at 15.19 and 15.69 eV split when including an extensive basis set. The

FIG. 2. Valence ionization spectra of OCS by (a) SRPES (Ref. 15) and (b) SAC-CI general-R method.

TABLE III. Ionization potential (I.P.) (in eV), monopole intensity and main configuration of the valence ionized states of OCS.

Experimental I.P.					SAC-CI general-R				
HeI ^a PES	X-ray ^b PES	$(e,2e)^c$	$(e,2e)^d$	SRPES ^e		State	I.P. (eV)	Intensity	Main configurations ($ C >0.3$)
				No.					
11.19	...	...	...	...	...	X ² Π	11.02	0.857	0.90(3π ⁻¹)
15.08	...	...	...	...	...	A ² Π	15.39	0.669	0.79(2π ⁻¹) - 0.33(3π ⁻¹ 4π2π ⁻¹)
16.04	...	...	...	...	...	B ² Σ ⁺	15.75	0.809	0.86(9σ ⁻¹)
...	...	...	...	...	...	² Π	17.40	0.019	0.79(3π ⁻² 4π) + 0.40(3π ⁻² 4π)
17.96	...	...	...	...	...	C ² Σ ⁺	17.70	0.708	0.81(8σ ⁻¹)
...	...	20.5	20.2	1	19.9	² Σ ⁺	20.89	0.033	0.54(3π ⁻² 10σ) + 0.54(3π ⁻² 10σ)
...	...	...	...	...	...	² Π	20.99	0.079	0.26(2π ⁻¹) + 0.67(3π ⁻² 4π) + 0.50(3π ⁻² 4π)
...	...	...	21.7	...	...	² Π	22.07	0.005	0.61(2π ⁻¹ 4π3π ⁻¹) + 0.47(2π ⁻¹ 4π3π ⁻¹) + 0.43(3π ⁻¹ 4π2π ⁻¹) + 0.40(3π ⁻¹ 4π2π ⁻¹)
...	...	...	...	...	...	² Σ ⁺	22.35	0.033	0.55(9σ ⁻¹ 4π3π ⁻¹) + 0.55(9σ ⁻¹ 4π3π ⁻¹) + 0.48(9σ ⁻¹ 4π3π ⁻¹) + 0.48(9σ ⁻¹ 4π3π ⁻¹)
...	23.0	23.5	22.8	2	23.0	² Σ ⁺	23.69	0.174	0.38(7σ ⁻¹) + 0.45(8σ ⁻¹ 4π3π ⁻¹) + 0.45(8σ ⁻¹ 4π3π ⁻¹) - 0.32(8σ ⁻¹ 4π3π ⁻¹) - 0.31(3π ⁻¹ 4π8σ ⁻¹) + 0.31(3π ⁻¹ 4π8σ ⁻¹)
...	...	...	...	...	...	² Σ ⁺	24.13	0.098	0.33(3π ⁻² 11σ) + 0.33(3π ⁻² 11σ)
...	25.0	...	24.1	...	...	² Σ ⁺	24.85	0.132	0.35(7σ ⁻¹) - 0.38(9σ ⁻¹ 4π2π ⁻¹) - 0.38(9σ ⁻¹ 4π2π ⁻¹) + 0.31(3π ⁻² 11σ) + 0.31(3π ⁻² 11σ)
...	...	...	...	...	...	² Π	25.41	0.009	0.59(2π ⁻¹ 4π3π ⁻¹) + 0.55(2π ⁻¹ 4π3π ⁻¹)
...	...	...	26.0	3	26.3	² Π	25.83	0.061	0.25(2π ⁻¹) - 0.56(3π ⁻¹ 4π2π ⁻¹) - 0.49(3π ⁻¹ 4π2π ⁻¹) - 0.32(9σ ⁻¹ 10σ2π ⁻¹)
...	...	...	...	...	...	² Σ ⁺	26.24	0.014	0.40(3π ⁻¹ 10σ2π ⁻¹) + 0.40(3π ⁻¹ 10σ2π ⁻¹) + 0.33(2π ⁻¹ 10σ3π ⁻¹) + 0.33(2π ⁻¹ 10σ3π ⁻¹)
...	...	...	...	...	...	² Σ ⁺	26.60	0.026	0.35(2π ⁻¹ 10σ3π ⁻¹) + 0.35(2π ⁻¹ 10σ3π ⁻¹)
...	27.4	27.1	27.9	4	27.8	² Σ ⁺	27.48	0.068	0.34(3π ⁻¹ 4π9σ ⁻¹) + 0.34(3π ⁻¹ 4π9σ ⁻¹)
...	...	...	...	...	...	² Π	27.73	0.005	0.65(3π ⁻² 5π) + 0.45(3π ⁻² 5π) - 0.34(3π ⁻² 10σ)
...	...	...	...	...	...	² Σ ⁺	27.78	0.012	0.43(3π ⁻² 12σ) + 0.43(3π ⁻² 12σ) - 0.34(3π ⁻² 11σ) - 0.34(3π ⁻² 11σ)
...	...	...	29.0	...	...	² Σ ⁺	28.06	0.004	0.35(2π ⁻¹ 11σ3π ⁻¹) + 0.35(2π ⁻¹ 11σ3π ⁻¹)
...	...	...	...	...	...	² Σ ⁺	28.67	0.062	0.33(3π ⁻¹ 4π8σ ⁻¹) + 0.33(3π ⁻¹ 4π8σ ⁻¹)
...	...	...	...	...	...	² Σ ⁺	29.68	0.014	0.46(3π ⁻¹ 11σ2π ⁻¹) + 0.46(3π ⁻¹ 11σ2π ⁻¹) + 0.33(2π ⁻¹ 11σ3π ⁻¹)
...	...	...	...	...	...	² Σ ⁺	30.20	0.035	0.30(3π ⁻² 1δ) + 0.30(3π ⁻² 1δ)
...	31.0	30.0	30.9	5	30.4	² Σ ⁺	30.49	0.041	0.38(2π ⁻¹ 4π9σ ⁻¹) + 0.38(2π ⁻¹ 4π9σ ⁻¹) + 0.37(9σ ⁻² 10σ)
...	...	...	...	...	...	² Σ ⁺	30.89	0.031	0.45(3π ⁻² 13σ) + 0.45(3π ⁻² 13σ) - 0.37(8σ ⁻¹ 4π9σ ⁻¹)
...	...	...	...	...	...	² Π	31.14	0.005	0.46(9σ ⁻² 4π) - 0.38(8σ ⁻¹ 4π9σ ⁻¹) - 0.38(9σ ⁻¹ 4π8σ ⁻¹) - 0.32(2π ⁻² 4π)
...	...	...	...	...	...	² Σ ⁺	31.15	0.104	0.30(7σ ⁻¹) - 0.45(8σ ⁻¹ 10σ9σ ⁻¹)
...	...	...	...	...	...	² Σ ⁺	31.72	0.042	0.49(8σ ⁻¹ 4π2π ⁻¹) + 0.49(8σ ⁻¹ 4π2π ⁻¹)
...	...	...	...	...	...	² Π	31.89	0.007	0.31(3π ⁻² 6π) - 0.31(3π ⁻¹ 5π2π ⁻¹)
...	...	...	...	...	...	² Σ ⁺	32.13	0.007	0.53(9σ ⁻¹ 5π3π ⁻¹) + 0.53(9σ ⁻¹ 5π3π ⁻¹)
...	...	...	...	...	...	² Π	32.41	0.005	0.45(9σ ⁻¹ 11σ3π ⁻¹) - 0.39(9σ ⁻¹ 12σ3π ⁻¹) - 0.32(9σ ⁻¹ 10σ2π ⁻¹) - 0.32(8σ ⁻¹ 12σ3π ⁻¹)
...	...	...	...	...	...	² Σ ⁺	32.71	0.004	0.51(3π ⁻¹ 5π9σ ⁻¹) + 0.51(3π ⁻¹ 5π9σ ⁻¹) + 0.40(3π ⁻¹ 5π8σ ⁻¹) + 0.40(3π ⁻¹ 5π8σ ⁻¹)
...	...	...	32.6	...	...	² Σ ⁺	33.11	0.019	0.35(9σ ⁻² 10σ) - 0.33(8σ ⁻¹ 5π3π ⁻¹) - 0.33(8σ ⁻¹ 5π3π ⁻¹) - 0.32(8σ ⁻¹ 10σ9σ ⁻¹)
...	...	...	...	...	...	² Σ ⁺	33.33	0.076	0.26(6σ) - 0.28(7σ ⁻¹ 4π3π ⁻¹) - 0.28(7σ ⁻¹ 4π3π ⁻¹)
...	...	...	...	...	...	² Σ ⁺	33.54	0.006	0.34(9σ ⁻¹ 10σ8σ ⁻¹)
...	...	...	...	...	...	² Σ ⁺	33.58	0.016	0.39(9σ ⁻¹ 10σ8σ ⁻¹) + 0.31(3π ⁻¹ 12σ2π ⁻¹) + 0.31(3π ⁻¹ 12σ2π ⁻¹)
...	...	...	...	...	...	² Σ ⁺	33.81	0.029	0.28(7σ ⁻¹ 4π3π ⁻¹)
...	...	...	...	...	...	² Σ ⁺	33.82	0.012	0.27(2π ⁻¹ 1δ3π ⁻¹)
...	...	...	...	...	...	² Σ ⁺	33.93	0.018	0.38(2π ⁻¹ 1δ3π ⁻¹) - 0.32(2π ⁻¹ 1δ3π ⁻¹)
...	...	...	...	...	...	² Σ ⁺	34.08	0.014	0.30(2π ⁻¹ 1δ3π ⁻¹)
...	...	...	...	...	...	² Σ ⁺	34.08	0.009	0.33(3π ⁻² 14σ) + 0.30(2π ⁻¹ 1δ3π ⁻¹)
...	...	...	34.2	...	...	² Σ ⁺	34.27	0.033	0.38(9σ ⁻² 11σ) + 0.33(8σ ⁻¹ 10σ9σ ⁻¹)
...	...	...	...	...	...	² Σ ⁺	34.40	0.029	0.45(9σ ⁻¹ 6π3π ⁻¹) + 0.45(9σ ⁻¹ 6π3π ⁻¹)
...	...	...	...	...	...	² Σ ⁺	34.53	0.010	0.37(3π ⁻² 14σ) + 0.37(3π ⁻² 14σ)
...	...	...	...	...	...	² Σ ⁺	34.58	0.018	0.46(8σ ⁻¹ 11σ9σ ⁻¹) - 0.32(9σ ⁻² 10σ)

TABLE III. (Continued.)

Experimental I.P.					SAC-CI general- <i>R</i>				
HeI ^a PES	X-ray ^b PES	(<i>e</i> ,2 <i>e</i>) ^c	(<i>e</i> ,2 <i>e</i>) ^d	SRPES ^e No.	State	I.P. (eV)	Intensity	Main configurations ($ C >0.3$)	
...	35.7	35.8	35.8	6	35.7	$2\Sigma^+$	35.30	0.141	$0.36(6\sigma^{-1})$
						$2\Sigma^+$	35.42	0.018	$0.42(2\pi^{-2}10\sigma)+0.42(2\pi^{-2}10\sigma)$
						$2\Sigma^+$	35.94	0.015	$0.50(9\sigma^{-1}11\sigma8\sigma^{-1})$
						$2\Sigma^+$	36.21	0.007	$0.41(2\pi^{-1}13\sigma3\pi^{-1})+0.41(2\pi^{-1}13\sigma3\pi^{-1})$
						$2\Sigma^+$	36.41	0.009	$0.32(9\sigma^{-2}12\sigma)-0.31(8\sigma^{-1}11\sigma9\sigma^{-1})$ $-0.31(9\sigma^{-1}11\sigma8\sigma^{-1})$

^aReference 10.

^bReference 1.

^cReference 12.

^dReference 13.

^eReference 15.

present calculation still overestimated the IPs of these correlation peaks, especially for peak (2).

Third to fifth correlation peaks were observed at 19.15 (3), 20.4 (4), and 20.8 (5) eV in SRPES.⁷ The peak (3) was assigned to $2\Sigma_g^+$ state in the recent two SRPES works;^{7,8} however, the assignment of the peak (4) was contradictory as $2\Pi_g$ (Ref. 7) or $2\Sigma_u^+$ (Ref. 8). The former assignment was based on the angular distribution and the latter was due to the angle dependence of the cross section relative to the incident radiation with the help of ADC calculation. In Ref. 8, it was also proposed that the peak (4) is composed by more-than-one shake-up states since it has unusual vibrational progression and the peak (5) was not observed. In the present calculation, two $2\Sigma_g^+$ states were calculated at 19.20 and 19.74 eV, which were attributed to the peak (3), two $2\Pi_g$ states were obtained at 20.15 and 20.23 eV for the peak (4), and one $2\Sigma_u^+$ state at 20.88 eV for the peak (5). Therefore, for the peak (4), we support the assignment of Roy *et al.*⁷ and proposed that the overlapping states, which were observed as two components at least by Baltzer *et al.*,⁸ were interpreted as the $2\Pi_g$ and $2\Sigma_u^+$ states. Note that the contribution of triples was found to be considerably large for the $2\Pi_g$ states of the peak (4).

The prominent peak (6) was observed at 21.35 eV and assigned to $2\Sigma_u^+$ state.⁷ Correspondingly, the general-*R* calculation gave $2\Sigma_u^+$ states with considerable intensity at 21.23 and 21.27 eV and these states were characterized as $(2\pi_g^{-2}7\sigma_u)$ interacting with $(4\sigma_u^{-1})$ state. We also found two $2\Pi_g$ states in this region. In the energy region from peak (7) observed at 21.6 eV to peak (9) at 23.3 eV,⁷ many $2\Sigma_u^+$ and $2\Sigma_g^+$ states were predicted: $2\Sigma_u^+$ states were calculated at 22.27, 22.77, and 23.72 eV and $2\Sigma_g^+$ states were also obtained at 23.33 and 23.60 eV. We attribute these states to the peaks from (7) to (9). It should be noted that the spectrum in this energy region was calculated very differently between the SDT-*R* and SDTQ-*R* methods. This is because the triples have large weights for some shake-up states and the quadruple *R*-operators become important in such case.

The peak (10) observed at 24.1 eV again has large intensity in SRPES.⁷ This peak was clearly attributed to the $2\Sigma_g^+$ state calculated at 24.19 eV with large intensity of 0.27 origi-

nating in $(6\sigma_g)^{-1}$. The previous SAC-CI SD-*R* calculation⁹ also reproduced the positions of the prominent peaks (6) and (10) and, hence, the qualitative picture of the spectrum. As the shoulder of the peak (10), peak (11) was observed at 24.6 eV.⁷ For this shoulder, we obtained three shake-up states; one $2\Sigma_g^+$ state at 24.77 eV and two $2\Sigma_u^+$ states at 24.59 and 24.81 eV.

Above the peak (11), $2\Sigma_u^+$ and $2\Sigma_g^+$ states were calculated at 25.25 and 25.83 eV, which are the candidates of the peaks (12) and (13) observed at 25.25 and 26.35 eV, respectively.⁷ The intensity of the former state was calculated smaller than the latter, which accords with the feature of the experimental spectrum. Around and above the double ionization threshold determined at 27.3 eV,¹¹ many shake-up states were calculated and some of them were characterized as triple electron processes. The prominent states with relatively large intensity are $2\Sigma_u^+$ and $2\Sigma_g^+$ states calculated at 27.02, 28.22, and 28.67 eV. They are assigned to the peaks (14) and (15), which were observed at 27.4 and 28.7 eV.⁷ Many continuous shake-up states around this energy region were also attributed to these peaks. Above this energy, no significant peaks which have considerable intensity were calculated and the peaks above 30 eV observed in SRPES⁷ may be attributed to the doubly ionized states.

B. OCS

The ground state electronic configuration of OCS is written as

$$(\text{core})^{14}(6\sigma)^2(7\sigma)^2(8\sigma)^2(2\pi)^4(9\sigma)^2(3\pi)^4.$$

The valence-ionization spectrum up to ~ 36 eV, whose energy region has been measured by the SRPES,¹⁵ was examined by the SAC-CI general-*R* method; 80 $2\Sigma^+$ states and 40 2Π states were solved for calculating the spectrum. The photoion-photoion coincidence (PIPICO) and double charge transfer experiments with the CIPSI calculation showed that several double-ionized states exist in the energy region of 31.8–40.4 eV.¹¹ From the preliminary calculation, the shape of the theoretical spectrum was found to be less dependent on the inclusion of higher-than-triple *R*-operators and, there-

TABLE IV. Ionization potential (I.P.) (in eV) and monopole intensity for the low-lying valence ionized states of OCS calculated with the augmented basis sets.

State	VTZ+2 <i>d</i> (present)		VTZ+2 <i>d</i> +diffuse ^a		VTZ+3 <i>d</i> ^b	
	I.P.	Intensity	I.P.	Intensity	I.P.	Intensity
² Σ ⁺	15.75	0.809	15.85	0.808	15.89	0.808
	17.70	0.708	17.87	0.706	17.89	0.703
	20.89	0.033	21.04	0.031	20.97	0.030
	22.35	0.033	22.44	0.031	22.47	0.030
² Π	11.02	0.857	11.08	0.859	11.21	0.857
	15.39	0.669	15.54	0.662	15.56	0.662
	17.40	0.019	17.45	0.020	17.44	0.019
	20.99	0.079	21.00	0.082	21.07	0.074

^aDiffuse function of [*1s1p1d*] are added on each atom.

^bOne polarization function with $\zeta_d=1.25$ is added on S atom.

fore, we exploited the general-*R* calculations up to triple *R*-operators with the SD-CI reference states. The resultant SAC-CI dimensions were 97 117 and 83 562 for ²Σ⁺ and ²Π states, respectively, as shown in Table I. In Fig. 2, the theoretical spectrum was presented with the latest SRPES¹⁵ measured at a photon energy of 60 eV. The peaks of the theoretical spectrum were convoluted by Gaussians with FWHM of 0.08*Δ*E* (in eV). The calculated IPs, intensities, and the characterization of the ionized states were summarized in Table III in comparison with the IPs determined by the He I PES,¹⁰ x-ray PES,¹ dipole (*e,2e*),^{12,13} and SRPES.¹⁵

High-resolution He I PES¹⁰ reported the outermost four main peaks at 11.185, 15.078, 16.042, and 17.957 eV, for X²Π, A²Π, B²Σ⁺, and C²Σ⁺ states, respectively. The A²Π and B²Σ⁺ states exist very close in energy and the ordering of these states was not reproduced in the previous SAC-CI SD-*R* calculation.⁹ In the present work, the general-*R* method gave the correct ordering of the main peaks: four peaks were calculated at 11.02(*X*), 15.39(*A*), 15.75(*B*), and 17.70(*C*) eV. In parallel, we also executed the SD-*R* calculation with the same basis set and found that the inclusion of the higher-order *R*-operators was important for the ordering of *A* and *B* states. Comparing the SAC-CI coefficients between the general-*R* and SD-*R* methods, weights of doubles for *A* state by the former method were found to be larger than those by the latter. The monopole intensities of the *A* and *C* states were considerably distributed to the satellite peaks as found in their intensities, 0.669 and 0.708, respectively, which also shows the contribution of doubles is significant for these two states.

The characterization of the inner-valence spectrum of OCS is still not definitive as pointed out in the SRPES study.¹⁵ Though there are extensive experimental works,^{1,2,12–15} only two theoretical works, namely SAC-CI SD-*R* method⁹ and the 2ph-TDA method,¹² have examined the inner-valence region. First two correlation peaks were clearly observed in the SRPES at 19.9 and 23.0 eV.¹⁵ In the present results, both the ²Σ⁺ and ²Π states were calculated at 20.89 and 20.99 eV, respectively. They are the candidates for the first peak and are characterized as (3π^{−2}10σ) and (3π^{−2}4π). We also obtained the ²Π state (17.40 eV) with considerable intensity at the foot of main peak *C*, which has never been reported in the experimental work. For the sec-

ond peak at 23.0 eV, four ²Σ⁺ states calculated at 22.35, 23.69, 24.13, and 24.85 eV and one ²Π state at 22.07 eV are attributed. White *et al.*¹³ observed detailed spectrum in this energy region using the dipole (*e,2e*); three peaks at 21.7, 22.8, and 24.1 eV. We assign ²Π state calculated at 22.07 eV to the peak 21.7 eV, three ²Σ⁺ states obtained at 22.35, 23.69, and 24.13 eV to the peak at 22.8 eV, and ²Σ⁺ state at 24.85 eV to the peak at 24.1 eV.

The complex and continuous spectrum was observed in the energy region of 25–40 eV.¹⁵ In the theoretical works,^{9,12} many shake-up states associated with the ionization of 6σ and 7σ MOs were predicted, however, it was criticized that these calculated states were separately clustered at around 25 and 36 eV against the experimental observation.¹⁵ The present calculation gave continuous band of the satellite spectrum as seen in Fig. 2. Since numerous states were calculated, one-to-one assignment of the observed peaks would be difficult; however, we proposed our assignment with the shake-up states whose intensity was calculated to be larger than 0.01. For the peak (3) observed at 26.3 eV,¹⁵ two ²Π states at 25.41 and 25.83 eV and two ²Σ⁺ states at 26.24 and 26.60 eV are attributed. Next peak (4) observed at around 27.8 eV¹⁵ is assigned to the ²Σ⁺ states calculated at 27.48 and 27.78 eV. For the peak observed by dipole (*e,2e*)¹³ at 29.0 eV, two ²Σ⁺ states calculated at 28.67 and 29.68 eV are candidates. Numerous states were calculated from 30 to 32 eV, which were attributed to the band (5) measured around 30.4 eV.

Dipole (*e,2e*)¹³ observed two bands at 32.6 and 34.2 eV, for which the SAC-CI general-*R* method gave 15 ²Σ⁺ states from 32.13 to 34.58 eV. This energy region is above the double ionization threshold¹¹ and, therefore, these states may be observed as the auto-ionizing states with the continuum background. Finally, for the peak at 35.7 eV,¹⁵ ²Σ⁺ state with large intensity of 0.141 was calculated at 35.30 eV. In the previous SAC-CI study,⁹ many shake-up states associated with (6σ)^{−1} were calculated at around this energy region.

IV. CONCLUSION

The valence ionization spectra of CS₂ and OCS were studied by the SAC-CI general-*R* method in both outer- and inner-valence regions. The present calculations finely simu-

lated the shapes of the experimental spectra and gave the detailed and quantitative assignments of the satellite peaks.

For CS₂, the present results presented the assignments of the satellites up to peak (15) measured by the SRPES.^{7,8} Four $^2\Pi_u$ states were calculated in the outer-valence region and were attributed to the observed peaks (1), (1)' and (2).^{7,8} Three peaks, (3), (4), and (5), were assigned to the $^2\Sigma_g^+$, $^2\Pi_g$, and $^2\Sigma_u^+$ states, respectively. The peak (4) observed with two components at least by recent SRPES⁸ was interpreted as the $^2\Pi_g$ and $^2\Sigma_u^+$ states. The prominent peaks (6) and (10) were assigned as the $^2\Sigma_u^+$ and $^2\Sigma_g^+$ states, respectively, and, for the satellites between these states, contributions of triples were remarkable: quadruple *R*-operators were important for reproducing the shape of the spectrum, especially for the peaks (8) and (9). Numerous $^2\Sigma_u^+$ and $^2\Sigma_g^+$ states, some of which were described by triples, continued up to double ionization threshold. Above 28 eV, no significant peaks were calculated.

The present calculation has also given detailed theoretical information on the valence-ionization spectrum of OCS. The position of the main peaks was correctly reproduced, especially for the ordering of *A* and *B* states, which was not obtained in the previous SAC-CI SD-*R* study:⁹ the higher *R*-operators were found to be significant for this ordering. For the first outer-valence satellite observed by the SRPES¹⁵ and (*e*,2*e*),^{12,13} both $^2\Sigma^+$ and $^2\Pi$ states were calculated. The second satellite of the SRPES¹⁵ was attributed to four $^2\Sigma^+$ states and one $^2\Pi$ state, and these five shake-up states were also assigned to the three satellites measured by dipole (*e*,2*e*).¹³ Below the $C^2\Sigma^+$ state, one satellite peak of $^2\Pi$ state, which has not been observed by experiment, was predicted. For the energy region of 24–36 eV, continuous band of the satellite spectrum was calculated, which reproduced the feature of SRPES.¹⁵ Above the double ionization threshold, some satellite peaks were also obtained.

ACKNOWLEDGMENTS

This study has been supported by the Special Fund for Scientific Research from the Japanese Ministry of Education, Science, Culture, and Sports and a grant of Kyoto University VBL project.

APPENDIX: EXAMINATION OF THE BASIS SETS

The present basis set was examined for the low-lying valence-ionized states of OCS with the additional polarization function and diffuse functions. The SAC-CI general-*R* calculations with the augmented basis sets were performed for the first ten states of each symmetry and the results were compared in Table IV. Since the results with the VTZ plus double polarization function (VTZ2P) were obtained by the calculations with the selection of the large number of solutions, the IPs with the better basis sets were calculated slightly higher than the results with VTZ2P. Nevertheless, there was no remarkable difference for the calculated IPs and intensities. From this examination, the present basis set of VTZ2P was found to be sufficient for describing the electron correlations of the shake-up states and suitable for calculating

entire valence ionization spectra with the moderate computational cost.

- ¹C. J. Allan, U. Gelius, D. A. Allison, G. Johansson, H. Siegbahn, and K. Siegbahn, *J. Electron Spectrosc. Relat. Phenom.* **1**, 131 (1972).
- ²A. W. Potts and T. A. Williams, *J. Electron Spectrosc. Relat. Phenom.* **3**, 3 (1974).
- ³J. Schirmer, W. Domcke, L. S. Cederbaum, W. von Niessen, and L. Asbrink, *Chem. Phys. Lett.* **61**, 30 (1979).
- ⁴F. Carnovale, M. G. White, and C. E. Brion, *J. Electron Spectrosc. Relat. Phenom.* **24**, 63 (1981).
- ⁵R. T. Leung and C. E. Brion, *Chem. Phys.* **82**, 87 (1983).
- ⁶M. J. Hubin-Franskin, J. Delwiche, P. Natalis, G. Caprace, and D. Roy, *J. Electron Spectrosc. Relat. Phenom.* **18**, 295 (1980).
- ⁷P. Roy, I. Nenner, P. Millie, P. Morin, and D. Roy, *J. Chem. Phys.* **87**, 2536 (1987).
- ⁸P. Baltzer, B. Wannberg, M. Lundqvist, L. Karlsson, D. M. P. Holland, M. A. MacDonald, M. A. Hayes, P. Tomasello, and W. von Niessen, *Chem. Phys.* **202**, 185 (1996).
- ⁹H. Nakatsuji, *Chem. Phys.* **76**, 283 (1983).
- ¹⁰L. S. Wang, J. E. Reutt, Y. T. Lee, and D. A. Shirley, *J. Electron Spectrosc. Relat. Phenom.* **47**, 167 (1988).
- ¹¹P. Millie, I. Nenner, P. Archirel, P. Lablanquie, P. Fournier, and J. H. D. Eland, *J. Chem. Phys.* **84**, 1259 (1986).
- ¹²J. P. D. Cook, M. G. White, C. E. Brion, W. Domcke, L. S. Cederbaum, J. Schirmer, and W. von Niessen, *J. Electron Spectrosc. Relat. Phenom.* **22**, 261 (1981).
- ¹³M. G. White, K. T. Leung, and C. E. Brion, *J. Electron Spectrosc. Relat. Phenom.* **23**, 127 (1981).
- ¹⁴K. T. Leung and C. E. Brion, *Chem. Phys.* **96**, 241 (1985).
- ¹⁵D. M. P. Holland and M. A. Macdonald, *Chem. Phys.* **144**, 279 (1990).
- ¹⁶H. Nakatsuji and K. Hirao, *J. Chem. Phys.* **68**, 2053 (1978).
- ¹⁷H. Nakatsuji, *Chem. Phys. Lett.* **59**, 362 (1978).
- ¹⁸H. Nakatsuji, *Chem. Phys. Lett.* **67**, 329 (1979).
- ¹⁹H. Nakatsuji, *Chem. Phys. Lett.* **67**, 334 (1979).
- ²⁰H. Nakatsuji, *Computational Chemistry-Review of Current Trends* (World Scientific, Singapore, 1997), Vol. 2, pp. 62–124.
- ²¹H. Nakatsuji, *Acta Chim. Hung.* **129**, 719 (1992).
- ²²H. Nakatsuji, *Chem. Phys.* **75**, 425 (1983).
- ²³H. Nakatsuji, O. Kitao, and T. Yonezawa, *J. Chem. Phys.* **83**, 723 (1985).
- ²⁴H. Nakatsuji and S. Saito, *J. Chem. Phys.* **91**, 6205 (1989).
- ²⁵H. Nakatsuji, M. Ehara, M. H. Palmer, and M. F. Guest, *J. Chem. Phys.* **97**, 2561 (1992).
- ²⁶H. Nakatsuji and M. Ehara, *J. Chem. Phys.* **101**, 7658 (1994).
- ²⁷H. Nakatsuji, J. Hasegawa, and M. Hada, *J. Chem. Phys.* **104**, 2321 (1996).
- ²⁸H. Nakatsuji, *Chem. Phys. Lett.* **177**, 331 (1991).
- ²⁹H. Nakatsuji, *J. Chem. Phys.* **83**, 713 (1985).
- ³⁰H. Nakatsuji, *J. Chem. Phys.* **83**, 5743 (1985).
- ³¹H. Nakatsuji, *J. Chem. Phys.* **94**, 6716 (1991).
- ³²M. Ehara and H. Nakatsuji, *Chem. Phys. Lett.* **282**, 347 (1998).
- ³³J. Hasegawa, M. Ehara, and H. Nakatsuji, *Chem. Phys. Lett.* **230**, 23 (1998).
- ³⁴M. Ehara, P. Tomasello, J. Hasegawa, and H. Nakatsuji, *Theor. Chem. Acc.* **102**, 161 (1999).
- ³⁵M. Ehara and H. Nakatsuji, *Spectrochim. Acta, Part A* **55**, 487 (1998).
- ³⁶M. Ehara, M. Ishida, and H. Nakatsuji, *J. Chem. Phys.* **114**, 8990 (2001).
- ³⁷M. Ishida, M. Ehara, and H. Nakatsuji, *J. Chem. Phys.* **116**, 1934 (2002).
- ³⁸M. Ehara, M. Ishida, K. Toyota, and H. Nakatsuji, in *Reviews in Modern Quantum Chemistry*, edited by K. D. Sen (World Scientific, Singapore, 2002), pp. 293–319.
- ³⁹A. Schafer, H. Horn, and R. Ahlrichs, *J. Chem. Phys.* **97**, 2571 (1992).
- ⁴⁰T. H. Dunning, Jr., *J. Chem. Phys.* **90**, 1007 (1989).
- ⁴¹D. E. Woon and T. H. Dunning, Jr., *J. Chem. Phys.* **98**, 1358 (1993).
- ⁴²K. P. Huber and G. Herzberg, *Molecular Spectra and Molecular Structure, IV. Constants of Diatomic Molecules* (Van Nostrand, New York, 1979).
- ⁴³S. Süzer, S. T. Lee, and D. A. Shirley, *Phys. Rev. A* **13**, 1842 (1976).
- ⁴⁴R. I. Martin and D. A. Shirley, *J. Chem. Phys.* **64**, 3685 (1976).
- ⁴⁵H. Nakatsuji, M. Hada, M. Ehara, J. Hasegawa, T. Nakajima, H. Nakai, O. Kitao, and K. Toyota, SAC/SAC-CI program system (SAC-CI96) for calculating ground, excited, ionized, and electron-attached states having singlet to septet spin multiplicities (1996).
- ⁴⁶M. J. Frisch *et al.* GAUSSIAN 98, Revision A.5, Gaussian, Inc., Pittsburgh PA, 1998.