TITLE:

Radiative decay of a massive particle and the nonthermal process in primordial nucleosynthesis

AUTHOR(S):

Kawasaki, M; Kohri, K; Moroi, T

CITATION:

Kawasaki, M ...[et al]. Radiative decay of a massive particle and the nonthermal process in primordial nucleosynthesis. PHYSICAL REVIEW D 2001, 63(10): 103502.

ISSUE DATE:

2001-05-15

URL:

<http://hdl.handle.net/2433/50451>

RIGHT:

Copyright 2001 American Physical Society

Radiative decay of a massive particle and the nonthermal process in primordial nucleosynthesis

Masahiro Kawasaki

Research Center for the Early Universe, School of Science, University of Tokyo, Tokyo 113-0033, Japan

Kazunori Kohri

Yukawa Institute for Theoretical Physics, Kyoto University, Kyoto 606-8502, Japan

Takeo Moroi

Department of Physics, Tohoku University, Sendai 980-8578, Japan

(Received 22 December 2000; published 9 April 2001)

We consider the effects on big bang nucleosynthesis (BBN) of the radiative decay of a long-lived massive particle. If high-energy photons are emitted after the BBN epoch ($t \sim 1 - 10^3$ sec), they may change the abundance of the light elements through photodissociation processes, which may result in a significant discrepancy between standard BBN and observation. Taking into account recent observational and theoretical developments in this field, we revise our previous study constraining the abundance of the radiatively decaying particles. In particular, on the theoretical side it was recently claimed that the nonthermal production of ${}^6\text{Li}$, which is caused by the photodissociation of ${}^4\text{He}$, most severely constrains the abundance of the radiatively decaying particle. We will see however, that it is premature to emphasize the importance of the nonthermal production of ${}^6\text{Li}$ because (i) the theoretical computation of the ${}^6\text{Li}$ abundance has a large uncertainty due to the lack of a precise understanding of the ${}^6\text{Li}$ production cross section, and (ii) the observational data of the ${}^6\text{Li}$ abundance has large errors.

DOI: 10.1103/PhysRevD.63.103502

PACS number(s): 98.80.Cq, 26.35.+c, 98.80.Ft

I. INTRODUCTION

Big bang nucleosynthesis (BBN) is one of the most important tools used to probe the early universe. Because it is very sensitive to the condition of the universe from 10^{-2} sec to 10^{12} sec, we can indirectly check the history of the universe and impose constraints on hypothetical particles by observational light element abundances.

There are many models of modern particle physics beyond the standard model, e.g., supergravity or superstring theory, which predict unstable massive particles with masses of $\mathcal{O}(100 \text{ GeV})$, such as the gravitino, Polonyi field, moduli, and so on. They have long lifetimes because their interactions are suppressed by inverse powers of the gravitational scale. Consequently, these exotic particles may decay at about the BBN epoch ($T \lesssim 1 \text{ MeV}$). If the massive particles radiatively decay, the emitted high-energy photons induce the electromagnetic cascade process. If the decay occurs after the BBN starts, the light elements would be destroyed by the cascade photons and their abundances would be changed significantly. Comparing the theoretically predicted light element abundances with the observational ones, we can impose constraints on the energy density, the mass, and the lifetime of the parent massive particle [1–3].¹ In particular, Holtmann and the present authors [3] performed the maximum likelihood analysis including both theoretical and observational errors and obtained the precise constraints.

After Ref. [3] was published, several new observational data of light elements were reported. As for the ${}^4\text{He}$ abundance, it was still unclear whether the observational value of the primordial ${}^4\text{He}$ mass fraction Y is low (~ 0.234) [9,10] or high (~ 0.244) [11]. However, Fields and Olive considered the HeI absorption effect and reanalyzed the data [12] and obtained a relatively middle value of Y (~ 0.238). On the other hand, as for the primordial D/H, although low values of D/H ($\sim 10^{-5}$) [13] had been measured and regarded as the primordial abundance, a relatively high value of D/H ($\sim 10^{-4}$) was claimed again by Tytler *et al.* in the high redshift quasistellar object (QSO) absorption systems [14]. In their paper they stressed that while the data may be inadequate to definitely conclude it to be of a precise value, there is still a possibility of the high D/H.

On the theoretical side it was recently claimed that the severest constraint on the radiatively decaying particle may be from the nonthermal production of ${}^6\text{Li}$, which is a secondary ${}^6\text{Li}$ production due to the background ${}^4\text{He}$ and the energetic T or ${}^3\text{He}$ produced by the ${}^4\text{He}$ photodissociation.² However, the observational data of the primordial component of ${}^6\text{Li}$ has large uncertainties. In addition, precise experimental data for the nuclear cross sections are not available. Therefore, it is unclear how important the nonthermal ${}^6\text{Li}$ production is once we take account of these uncertainties.

With these new developments in theory and observation, we revise the previous constraint on the radiative decay of

¹There have been many articles concerning BBN and long-lived massive particles; see [4,5] for hadronic decaying particles, [6] for residual annihilations, and [7,8] for decaying neutrinos.

²Such a possibility of the secondary process had already been pointed out by the earlier works for hadronic decaying particles [5].

long-lived particles. We obtain the photon spectrum by solving the Boltzmann equation numerically [2]. In addition, we perform the Monte Carlo simulation which includes both the experimental and theoretical errors. Then, we estimate the confidence levels by performing the maximum likelihood method including both the theoretical and the observational errors.

This paper is organized as follows. In Sec. II we briefly review the current status of the observational data. In Sec. III we introduce the formulations for the photodissociation and nonthermal ${}^6\text{Li}$ production. In Sec. IV we compare the theoretical predictions with the observations. Section V is devoted to conclusions.

II. OBSERVATIONAL LIGHT ELEMENT ABUNDANCES

Here we summarize the current status of the observational light element abundances. The primordial D/H is measured in the high redshift QSO absorption systems. Recently new deuterium data was obtained from observation of QSO HS 0105+1619 at $z = 2.536$ [16]. It was found that the cloud is neutral and has a simple structure. Five Lyman series transitions caused by D and H were observed there. The reported value of the deuterium abundance was relatively low, $(\text{D}/\text{H})^{\text{obs}} = (2.54 \pm 0.23) \times 10^{-5}$. Combined with the previous “low D” data which were obtained by the clouds at $z = 3.572$ towards Q 1937–1009 and at $z = 2.504$ towards Q 1009+2956 [13], the primordial abundance is obtained as

$$\text{low D: } (\text{D}/\text{H})^{\text{obs}} = (3.0 \pm 0.4) \times 10^{-5}. \quad (1)$$

We call this value “low D.” On the other hand, Webb *et al.* observed a high deuterium abundance in relatively low redshift absorption systems at $z = 0.701$ towards Q 1718+4807 [17],

$$\text{high D: } (\text{D}/\text{H})^{\text{obs}} = (2.0 \pm 0.5) \times 10^{-4}. \quad (2)$$

Tytler *et al.* [14] also observed the clouds independently and obtained the similar value. Since Webb *et al.* and Tytler *et al.* did not obtain the full spectra of the Lyman series in their observations, the precise fitting of D/H based on the “high D” data might be inadequate. However, the possibility of “high D” have not been excluded yet. Therefore, we also consider the possibility of “high D” and include it in our analysis.

For ${}^3\text{He}$, we use the presolar measurements. In this paper, we do not rely upon models of galactic and stellar chemical evolution because of the large uncertainty in extrapolating back to the primordial abundance. But it is reasonable to assume that ${}^3\text{He}/\text{D}$ is an increasing function of the cosmic time, because D is the most fragile isotope and is always destroyed whenever ${}^3\text{He}$ is destroyed. Using the solar system data reanalyzed by Geiss [18],

$$r_{3,2,\odot}^{\text{obs}} \equiv ({}^3\text{He}/\text{D})_{\odot}^{\text{obs}} = 0.591 \pm 0.536, \quad (3)$$

where $\odot$ denotes the presolar abundance. We take this to be an upper bound on the primordial ${}^3\text{He}$ to D ratio $r_{3,2}^{\text{obs}}$,

$$r_{3,2}^{\text{obs}} \leq r_{3,2,\odot}^{\text{obs}}. \quad (4)$$

Although in the standard scenario the theoretical prediction satisfies the above constraint, ${}^4\text{He}$ photodissociation produces both D and ${}^3\text{He}$ and can raise the ${}^3\text{He}$ to D ratio [19]. Hence, we include this constraint into our analysis.

The primordial ${}^4\text{He}$ mass fraction Y is inferred from observation of recombination lines from the low metallicity extragalactic HII regions. Since ${}^4\text{He}$ is produced with the oxygen in stars, the primordial value is obtained to regress to the zero metallicity $\text{O}/\text{H} \rightarrow 0$ for the observational data. Recently, Fields and Olive [12] reanalyzed the data including the HeI absorption effect and they obtained

$$Y^{\text{obs}} = 0.238 \pm (0.002)_{\text{stat}} \pm (0.005)_{\text{syst}}, \quad (5)$$

where the first error is the statistical uncertainty and the second error is the systematic one. We adopt the above value as the observational Y .

The primordial ${}^7\text{Li}/\text{H}$ is observed in the Pop II old halo stars. We adopt the recent measurements by Bonifacio and Molaro [20]

$$\begin{aligned} \log_{10}[({}^7\text{Li}/\text{H})^{\text{obs}}] &= -9.76 \pm (0.012)_{\text{stat}} \pm (0.05)_{\text{syst}} \\ &\pm (0.3)_{\text{add}}. \end{aligned} \quad (6)$$

Here we have added the additional uncertainty for fear that the ${}^7\text{Li}$ in halo stars might have been supplemented (by production in cosmic ray interactions) or depleted (in stars) [21].

It is much more difficult to observe the primordial component of ${}^6\text{Li}$ because ${}^6\text{Li}$ is so much rarer than ${}^7\text{Li}$. Unfortunately, enough data have not been obtained to find the “Spite plateau” of ${}^6\text{Li}$. However, we can set an upper bound on ${}^6\text{Li}/{}^7\text{Li}$, since it is generally believed that the evolution of ${}^6\text{Li}$ is dominated by the production through $p, \alpha\text{-C, N, O}$ cosmic ray spallation (reactions of cosmic rays with the interstellar medium). Intrinsically the models of the nucleosynthesis through the cosmic ray spallation were motivated to simultaneously agree with the whole observational Li-Be-B abundances [22–24]. On the other hand, recently it was claimed that the observational ${}^6\text{Li}$ abundance in halo stars is too abundant from the point of view of the cosmic ray energy if ${}^9\text{Be}$ is fit by the model of the cosmic ray metal [25]. Therefore, there seems to be some uncertainties in the models of the cosmic ray spallation. In this situation, however, at least it would be safe to assume that ${}^6\text{Li}$ abundance increases as the metallicity increases. Today we observe only the ${}^6\text{Li}$ to ${}^7\text{Li}$ ratio in low-metallicity ($[\text{Fe}/\text{H}] \leq -2.0$) halo stars [26],

$$r_{6,7,\text{halo}}^{\text{obs}} \equiv ({}^6\text{Li}/{}^7\text{Li})_{\text{halo}}^{\text{obs}} = 0.05 \pm 0.02. \quad (7)$$

We take this value as an upper bound on the primordial value $r_{6,7}^{\text{obs}}$,

$$r_{6,7}^{\text{obs}} \leq r_{6,7,\text{halo}}^{\text{obs}}. \quad (8)$$

TABLE I. List of photodissociation processes, and the $1\text{-}\sigma$ uncertainty in the cross sections. Since there are no experimental data on photodissociation of ${}^7\text{Be}$, we assume that the rate, threshold, and uncertainty for reaction 13 is the same as for reaction 11, and the rate for reaction 14 is the sum of the rates for reactions 10 and 12.

	Photodissociation reactions	$1\text{-}\sigma$ uncertainty	Threshold energy	Ref.
1.	$\text{D} + \gamma \rightarrow p + n$	6%	2.2 MeV	[27]
2.	$\text{T} + \gamma \rightarrow n + \text{D}$	14%	6.3 MeV	[28,29]
3.	$\text{T} + \gamma \rightarrow p + 2n$	7%	8.5 MeV	[29]
4.	${}^3\text{He} + \gamma \rightarrow p + \text{D}$	10%	5.5 MeV	[30]
5.	${}^3\text{He} + \gamma \rightarrow n + 2p$	15%	7.7 MeV	[30]
6.	${}^4\text{He} + \gamma \rightarrow p + \text{T}$	4%	19.8 MeV	[30]
7.	${}^4\text{He} + \gamma \rightarrow n + {}^3\text{He}$	5%	20.6 MeV	[31,32]
8.	${}^4\text{He} + \gamma \rightarrow p + n + \text{D}$	14%	26.1 MeV	[33]
9.	${}^6\text{Li} + \gamma \rightarrow \text{anything}$	4%	5.7 MeV	[34]
10.	${}^7\text{Li} + \gamma \rightarrow 2n + \text{anything}$	9%	10.9 MeV	[34]
11.	${}^7\text{Li} + \gamma \rightarrow n + {}^6\text{Li}$	4%	7.2 MeV	[34]
12.	${}^7\text{Li} + \gamma \rightarrow {}^4\text{He} + \text{anything}$	9%	2.5 MeV	[34]
13.	${}^7\text{Be} + \gamma \rightarrow p + {}^6\text{Li}$	4%		
14.	${}^7\text{Be} + \gamma \rightarrow \text{anything except } {}^6\text{Li}$	9%		

III. PHOTODISSOCIATION AND NONTHERMAL PRODUCTION OF ${}^6\text{Li}$

A. Photodissociation

In order to discuss the effect of high-energy photons on BBN, we need the shape of the photon spectrum induced by the primary high-energy photons from the decay of the massive particle X . In the thermal bath (mixture of photons γ_{BG} , electrons e_{BG}^- , and nucleons N_{BG}), high energy photons lose their energy by the following cascade processes: double-photon pair creation ($\gamma + \gamma_{\text{BG}} \rightarrow e^+ + e^-$); photon-photon scattering ($\gamma + \gamma_{\text{BG}} \rightarrow \gamma + \gamma$); pair creation in nuclei ($\gamma + N_{\text{BG}} \rightarrow e^+ + e^- + N$); Compton scattering ($\gamma + e_{\text{BG}}^- \rightarrow \gamma + e^-$); inverse Compton scattering ($e^\pm + \gamma_{\text{BG}} \rightarrow e^\pm + \gamma$). In this study we numerically solved the Boltzmann equation including the above processes, and obtained the distribution function of photons, $f_\gamma(E_\gamma)$.

The cascade photons induce the photodissociation of the light elements, which modifies the result of standard BBN (SBBN). The evolutions of the light nuclei abundances are governed by the following Boltzmann equation:

$$\begin{aligned} \frac{dn_N}{dt} + 3Hn_N = & \left[\frac{dn_N}{dt} \right]_{\text{SBBN}} \\ & - n_N \sum_{N'} \int dE_\gamma \sigma_{N \rightarrow N'}(E_\gamma) f_\gamma(E_\gamma) \\ & + \sum_{N''} n_{N''} \int dE_\gamma \sigma_{N'' \rightarrow N}(E_\gamma) f_\gamma(E_\gamma), \end{aligned} \quad (9)$$

where n_N is the number density of the nuclei N , and $[dn_N/dt]_{\text{SBBN}}$ denotes the SBBN contribution to the Boltzmann equation. In Table I we list the photodissociation processes included in our computation. In this study the model

parameters are the baryon to photon ratio (η), the lifetime of X (τ_X), the mass of X (m_X), and the yield variable Y_X of X after electron-positron annihilation,

$$Y_X = n_X/n_\gamma, \quad (10)$$

where n_γ is the number density of the photon.³ In this paper we assume that X decays only into photons, i.e., $m_X Y_X$ corresponds to $\Delta\rho_\gamma/n_\gamma$. Then, the photodissociation rates depend on the combination $m_X Y_X$ which characterizes the amount of the energy of the injected photons $\Delta\rho_\gamma$ as far as m_X is much larger than 20 MeV [35].

B. Nonthermal ${}^6\text{Li}$ production

As pointed out by Jedamzik [15], both T and ${}^3\text{He}$ are produced through the photodissociation of ${}^4\text{He}$,

$${}^4\text{He} + \gamma \rightarrow \begin{cases} n + {}^3\text{He}, \\ p + \text{T}. \end{cases} \quad (11)$$

They are still energetic and have enough kinetic energies to produce ${}^6\text{Li}$ through the following processes with the background ${}^4\text{He}$:

$$\text{T} + {}^4\text{He} \rightarrow {}^6\text{Li} + n, \quad (12)$$

$${}^3\text{He} + {}^4\text{He} \rightarrow {}^6\text{Li} + p, \quad (13)$$

until they are stopped by the ionization loss through the plasma excitation in the electromagnetic plasma. The threshold energy of the ${}^6\text{Li}$ production is $E_{6^3\text{He}}^{\text{th}} = 4.03$ MeV for

³Note that in Ref. [3], $Y_X = n_X/n_\gamma$ is defined before electron-positron annihilation (e^+e^- ann.). Then they have a relationship $Y_X|_{\text{after } e^+e^- \text{ ann.}} = \frac{4}{11} Y_X|_{\text{before } e^+e^- \text{ ann.}}$.

^3He , and $E_6^{\text{th}} = 4.80$ MeV for T. Then, the abundance of ^6Li produced through the nonthermal (NT) process in Eq. (12) is governed by

$$\left[\frac{dn_{^6\text{Li}}}{dt} \right]_{\text{NT}} = n_{^4\text{He}} \int_{E_4^{\text{th}} + 4E_6^{\text{th}}}^{\infty} dE_{\gamma} \sigma_{^4\text{He}(\gamma,p)\text{T}}(E_{\gamma}) f_{\gamma}(E_{\gamma}) \times \int_{E_6^{\text{th}}}^{(E_{\gamma} - E_4^{\text{th}})/4} n_{^4\text{He}} \sigma_{\text{T}(\alpha,n)^6\text{Li}}(E) \left(\frac{dE}{dx} \right)^{-1} dE, \quad (14)$$

where $n_{^6\text{Li}}(n_{^4\text{He}})$ denotes the number density of ^6Li (^4He). $\sigma_{^4\text{He}(\gamma,p)\text{T}}(E_{\gamma})$ is the cross section of the ^4He photodissociation, E_4^{th} is the threshold energy of the photodissociation process, $f_{\gamma}(E_{\gamma})$ is the photon spectrum which is obtained by solving the Boltzmann equation, and $\sigma_{\text{T}(\alpha,n)^6\text{Li}}(E)$ is the cross section of the process in Eq. (12). dE/dx denotes the rate of the ionization loss while the charged particle T is running a distance dx in the electromagnetic plasma. The rate of the ionization loss is expressed by [36]

$$\frac{dE}{dx} = \frac{Z^2 \alpha}{\beta^2} \omega_p^2 \ln \left(\frac{\Lambda m_e \beta^2}{\omega_p} \right), \quad (15)$$

where ω_p^2 is the plasma frequency ($= 4\pi n_e \alpha / m_e$), n_e is the electron number density, m_e is electron mass, Z is the charge, $\Lambda \sim \mathcal{O}(1)$ is a constant and β is the velocity. The effect of the process (13) is described by replacing the suffix T with ^3He in Eq. (14).

We include the above two processes of the nonthermal ^6Li production in BBN code and compute the ^6Li abundance. In the computation we adopt the experimental cross section $\sigma_{\text{T}(\alpha,n)^6\text{Li}} = 35 \pm 1.4$ mb [37] commonly for the two processes. Because we have only one data point at the kinetic energy $E_T = 28$ MeV in the laboratory system, we assume that the cross section is constant for the whole energy region and neglect its energy dependence. Then, we integrate the second factor in Eq. (14) up to a high energy. One can easily find that there exists a serious problem in this procedure if it is compared to the case of the original photodissociation where the photodissociation rates steeply decrease as the energy increases. Because we have the experimental data for the ^4He photodissociation rates only up to about 100 MeV for the photon energy [31–33], we should interpolate the photodissociation rates in a high-energy region because of the mild dumping of the integrand. Then, the integration has a large uncertainty ($\sim 20\%$) when we change the upper limit of the integration from 500 MeV to 1 GeV.⁴ Therefore, in

this situation we adopt a 20% error for the nonthermal ^6Li production rates and perform the Monte Carlo simulation which includes them.⁵

C. Constraint from cosmic microwave background

In addition to the photodissociation process, there also exists another constraint. A radiative decay process releases a net photon energy into the electromagnetic plasma. The emitted photons should be thermalized soon, otherwise the photon spectrum deviates from the blackbody, which contradicts the observation of the cosmic microwave background (CMB) [38]. This leads to the following constraints:

$$m_X Y_X \lesssim 2.0 \times 10^{-12} \text{ GeV} \left(\frac{\tau_X}{10^{10} \text{ sec}} \right)^{1/2}, \quad (16)$$

for μ distortion [$1.8 \times 10^6 \text{ sec } (\Omega_B h^2 / 0.02)^{2/3} \lesssim \tau_X \lesssim 2.3 \times 10^9 \text{ sec } (\Omega_B h^2 / 0.02)$], and

$$m_X Y_X \lesssim 1.9 \times 10^{-12} \text{ GeV} \left(\frac{\tau_X}{10^{10} \text{ sec}} \right)^{1/2}, \quad (17)$$

for y distortion [$2.3 \times 10^9 \text{ sec } (\Omega_B h^2 / 0.02) \lesssim \tau_X \lesssim 10^{12} \text{ sec}$].

IV. COMPARISON WITH OBSERVATIONAL LIGHT ELEMENT ABUNDANCES

In Fig. 1 we plot the theoretically predicted ^6Li to ^7Li ratio ($\equiv r_{6,7}^{\text{th}}$) in $(\tau_X, m_X Y_X)$ plane. The solid line represents the model parameters which predict the observational mean value of $r_{6,7}^{\text{th}}$ and the dashed line denotes the observational $2\text{-}\sigma$ upper bound. From the figure, one may think that the mean value of the theoretical prediction constrains $m_X Y_X$ severely. We should bear in mind, however, that the theoretical prediction has a large uncertainty which comes from the errors of the production rates, and in addition the observational constraint also has a large error. To take account of these uncertainties systematically, we performed the maximum likelihood analysis [3] including both the theoretical and the observational errors. Here we assumed that the theoretical predictions of $(\text{D}/\text{H})^{\text{th}}$, Y^{th} , $\log_{10}[(^7\text{Li}/\text{H})^{\text{th}}]$, $r_{3,2}^{\text{th}} = (^3\text{He}/\text{D})^{\text{th}}$, and $r_{6,7}^{\text{th}}$ obey the Gaussian probability distribution functions (p.d.f.'s) with the widths given by the $1\text{-}\sigma$ errors. Concerning the observational values, $(\text{D}/\text{H})^{\text{obs}}$, Y^{obs} , and $\log_{10}[(^7\text{Li}/\text{H})^{\text{obs}}]$ are assumed to obey the Gaussian p.d.f.'s while we treat $r_{3,2}^{\text{obs}}$ and $r_{6,7}^{\text{obs}}$ as non-Gaussian variables [3].

In Fig. 2 we plot the results of the χ^2 fitting by using the method of the maximum likelihood analysis. The solid (dashed) line denotes the low D (high D) constraint. The

⁴In addition, there may be another larger uncertainty which comes from the differences of the method for the interpolation because we do not know the correct shape of the cross sections. In this case, the obtained constraint would be weaker.

⁵If the cross section $\sigma_{\text{T}(\alpha,n)^6\text{Li}}$ decreases at a high energy like other nuclear interactions, the ^6Li production is less important. As shown later, the resultant constraint is not changed even if we neglect the ^6Li production.

FIG. 1. Plot of ${}^6\text{Li}$ to ${}^7\text{Li}$ ratio in $(\tau_X, m_X Y_X)$ plane for various baryon to photon ratio ($\eta = n_B/n_\gamma$) in (a) $\eta = 2 \times 10^{-10}$, (b) $\eta = 4 \times 10^{-10}$, (c) $\eta = 5 \times 10^{-10}$, and (d) $\eta = 6 \times 10^{-10}$. The solid line denotes the observational mean value of ${}^6\text{Li}/{}^7\text{Li}$ and the dashed line denotes the observational $2\text{-}\sigma$ upper bound.

dotted line denotes the upper bound from the CMB constraint. In the figure, the region below the lines is consistent with the observations. The constraint from the CMB is almost always weaker than that from BBN. The main feature of the difference between high D and low D is that the low D constraint is severer than high D for a relatively long lifetime case ($\tau_X \gtrsim 3 \times 10^6$ sec). That is because the high D constraint modestly allows for the overproduction of ${}^3\text{He}$ accompanying the ${}^4\text{He}$ photodissociation. On the other hand, the high D constraint is more stringent for shorter lifetimes since the D dissociation is more important than the ${}^4\text{He}$ photodissociation.

FIG. 2. Plot of the contour of the confidence level in $(\tau_X, m_X Y_X)$ plane. The solid (dashed) line denotes the 95% C.L. for low D (high D) projected on an η axis. The dotted line denotes the upper bound which comes from CMB constraint.

The obtained upper bound does not change our earlier results as much [3]. It became slightly weaker because we included the η dependence for the photodissociation rates ($\Gamma \propto 1/\eta$) in this analysis.⁶ We find that the nonthermally produced ${}^6\text{Li}$ mildly contributes to the bound.⁷ The main reason is that both the theoretical computation and observational data have very large uncertainties which amount to about 30–40 %.

Assuming that the parent massive particle is the gravitino and that it dominantly decays into a photon and a photino ($\psi_{3/2} \rightarrow \tilde{\gamma} + \gamma$), the lifetime $\tau_{3/2}$ is related to the gravitino mass $m_{3/2}$ as

$$\tau_{3/2} \simeq 4 \times 10^5 \text{ sec} \times (m_{3/2}/1 \text{ TeV})^{-3}. \quad (18)$$

Assuming that the gravitino is produced through the thermal scattering in the reheating process after inflation,⁸ we relate the abundance $Y_{3/2} = n_{3/2}/n_\gamma$ of the gravitino with the reheating temperature T_R [2],

$$Y_{3/2} \simeq 1.1 \times 10^{-11} \times (T_R/10^{10} \text{ GeV}). \quad (19)$$

In Fig. 3 we plot the upper bound on the reheating temperature after inflation at 95% C.L. as a function of the gravitino mass. Here we can read off the constraint by using the relationship of the scaling, $\Delta \rho_\gamma/n_\gamma = \frac{1}{2} m_{3/2} Y_{3/2} (= m_X Y_X)$ because we assumed X decays into two photons. From the figure we can obtain the upper bound on the reheating temperature

$$m_{3/2} = 100 \text{ GeV} \quad (\tau_{3/2} \simeq 4 \times 10^8 \text{ sec}): T_R \lesssim 1 \times 10^7 \text{ GeV},$$

$$m_{3/2} = 1 \text{ TeV} \quad (\tau_{3/2} \simeq 4 \times 10^5 \text{ sec}): T_R \lesssim 1 \times 10^9 \text{ GeV},$$

⁶The η dependence is understood as follows. The soft photons produced in the electromagnetic cascade scatter off the background electrons and nucleons and lose their energy. Thus the number density of soft photons with energy larger than the threshold decreases as a scattering rate which is proportional to η . Therefore, the photodissociation rates are proportional to $1/\eta$.

⁷Tritium is unstable with a lifetime $\tau_T = 5.614 \times 10^8$ sec, and decays into ${}^3\text{He}$ whose charge is two. Thus, because ${}^3\text{He}$ is prone to stop ping much easier than T by the ionization loss, we might overestimate the ${}^6\text{Li}$ production in parameter regions where the stopping time $\tau_{stop} = \int_E^0 (dE/dt)^{-1} dE \simeq 2.5 \times 10^9 \text{ sec} (T/\text{eV})^{-3} (E/\text{MeV})^{3/2}$ is longer than the lifetime of tritium, i.e., for $T \lesssim 1.7$ eV. Therefore, at a long lifetime $\tau_X \gtrsim 5 \times 10^{11}$ sec, our constraint might become weaker by about a factor of two. However, it is expected that the effect would not change the result significantly because the ${}^3\text{He}$ overproduction gives a severer constraint there.

⁸Although these days it was claimed that gravitinos are also produced in the preheating epoch nonthermally [39–41], we do not consider such processes here because there are some ambiguities on the estimations and they depend on various model parameters. If the nonthermal production is effective, however, the obtained constraint might be more severe.

FIG. 3. Plot of the contour of the confidence level in $(m_{3/2}, T_R)$ plane. The solid (dashed) line denotes the 95% C.L. for low D (high D). The dotted line denotes the upper bound which comes from CMB constraint.

$$m_{3/2} = 3 \text{ TeV} \quad (\tau_{3/2} \approx 1 \times 10^4 \text{ sec}): T_R \lesssim 9 \times 10^{11} \text{ GeV}, \quad (20)$$

at 95% C.L.

V. CONCLUSION

In this paper we have studied the effects on primordial nucleosynthesis of the radiative decay of a long-lived massive particle X using new observational data. We have also considered the nonthermal ${}^6\text{Li}$ production caused by ener-

getic T and ${}^3\text{He}$ produced by the ${}^4\text{He}$ photodissociation. We obtained the photon spectrum through the electromagnetic cascade process by solving a set of Boltzmann equations numerically. In addition, to estimate the theoretical errors we performed Monte Carlo simulation including the theoretical uncertainties which come from those of nuclear reaction rates. To obtain the degree of agreements between theory and observation, we performed the maximum likelihood method and the χ^2 fitting including both the observational and theoretical errors.

As a result we have obtained the upper bound on the abundance $m_X Y_X$ as a function of its lifetime τ_X . The result does not change our previous works significantly. This is because the theoretical and observational errors for ${}^6\text{Li}$ are significantly large, and it contributes to the constraints more weakly than the ${}^3\text{He}$ overproduction accompanying the ${}^4\text{He}$ photodissociation. Therefore, we have concluded that it is premature to emphasize the importance of the nonthermal production of ${}^6\text{Li}$.

We have also applied the results obtained by a generic radiatively decaying particle to gravitino $\psi_{3/2}$, and we have got the upper bound on the reheating temperature after primordial inflation as a function of the mass, $T_R \lesssim 10^7 - 10^9 \text{ GeV}$ for $m_{3/2} = 100 \text{ GeV} - 1 \text{ TeV}$ (95% C.L.).

ACKNOWLEDGMENTS

The work of M.K. is supported by Priority Area 707 ‘‘Supersymmetry and Unified Theory of Elementary Particles.’’ The work of T.M. is supported by the Grant-in-Aid for Scientific Research from the Ministry of Education, Science, Sports and Culture of Japan No. 12047201.

- [1] J. Ellis, G. B. Gelmini, J. L. Lopez, D. V. Nanopoulos, and S. Sarkar, Nucl. Phys. **B373**, 399 (1992).
- [2] M. Kawasaki and T. Moroi, Prog. Theor. Phys. **93**, 879 (1995).
- [3] E. Holtmann, M. Kawasaki, K. Kohri, and T. Moroi, Phys. Rev. D **60**, 023506 (1999).
- [4] M. H. Reno and D. Seckel, Phys. Rev. D **37**, 3441 (1988).
- [5] S. Dimopoulos, R. Esmailzadeh, L. J. Hall, and G. D. Starkman, Astrophys. J. **330**, 545 (1988); Nucl. Phys. **B311**, 699 (1989).
- [6] J. A. Frieman, E. W. Kolb, and M. S. Turner, Phys. Rev. D **41**, 3080 (1990); J. S. Hagelin and R. D. J. Parker, Nucl. Phys. **B329**, 464 (1990); J. S. Hagelin, R. D. J. Parker, and A. Hankey, Phys. Lett. B **215**, 397 (1988); S. Hannestad, *ibid.* **452**, 23 (1999); P. McDonald, R. J. Scherrer, and T. P. Walker, Phys. Rev. D **63**, 023001 (2001).
- [7] M. Kawasaki, P. Kernan, H.-S. Kang, R. J. Scherrer, G. Steigman, and T. P. Walker, Nucl. Phys. **B419**, 105 (1994); M. Kawasaki, K. Kohri, and K. Sato, Phys. Lett. B **430**, 132 (1998).
- [8] S. Dodelson, G. Gyuk, and M. S. Turner, Phys. Rev. D **49**, 5068 (1994); S. Hannestad, *ibid.* **57**, 2213 (1998); A. D. Dolgov, S. H. Hansen, S. Pastor, and D. V. Semikoz, Phys. Lett. B **548**, 385 (1999).
- [9] B. E. J. Pagel *et al.*, Mon. Not. R. Astron. Soc. **255**, 325 (1992).
- [10] K. A. Olive, G. Steigman, and E. D. Skillman, Astrophys. J. **483**, 788 (1997).
- [11] Y. I. Izotov, T. X. Thuan, and V. A. Lipovetsky, Astrophys. J., Suppl. Ser. **108**, 1 (1997); Y. I. Izotov and T. X. Thuan, Astrophys. J. **500**, 188 (1998).
- [12] B. D. Fields and K. A. Olive, Astrophys. J. **506**, 177 (1998).
- [13] S. Burles and D. Tytler, Astrophys. J. **507**, 732 (1998).
- [14] D. Tytler, S. Burles, L. Lu, X.-M. Fan, and A. Wolfe, Astron. J. **117**, 63 (1999).
- [15] K. Jedamzik, Phys. Rev. Lett. **84**, 3248 (2000).
- [16] J. M. O’Meara, D. Tytler, D. Kirkman, N. Suzuki, J. X. Prochaska, D. Lubin, and A. M. Wolfe, astro-ph/0011179.
- [17] J. K. Webb *et al.*, Nature (London) **388**, 250 (1997).
- [18] J. Geiss, in *Origin and Evolution of the Elements*, edited by N. Prantzos, E. Vangioni-Flam, and M. Cassé (Cambridge University Press, Cambridge, 1993), p. 89.
- [19] G. Sigl, K. Jedamzik, D. N. Schramm, and V. S. Berezinsky, Phys. Rev. D **52**, 6682 (1995).
- [20] P. Bonifacio and P. Molaro, Mon. Not. R. Astron. Soc. **285**, 847 (1997).
- [21] B. D. Fields, K. Kainulainen, K. A. Olive, and D. Thomas, New Astron. **1**, 77 (1996).

- [22] D. K. Duncan, D. L. Lambert, and M. Lemke, *Astrophys. J.* **401**, 584 (1992); M. Cassé, R. Lehoucq, and E. Vangioni-Flam, *Nature (London)* **373**, 318 (1995); R. Ramaty, B. Kozlovsky, and R. E. Lingenfelter, *Astrophys. J. Lett.* **438**, L21 (1995).
- [23] M. Lemoine, D. N. Schramm, J. W. Truran, and C. J. Copi, *Astrophys. J.* **478**, 554 (1997).
- [24] B. D. Fields and K. A. Olive, *New Astron.* **4**, 255 (1999).
- [25] R. Ramaty, S. T. Scully, R. E. Lingenfelter, and B. Kozlovsky, *Astrophys. J.* **534**, 747 (2000).
- [26] V. V. Smith, D. L. Lambert, and P. E. Nissen, *Astrophys. J.* **408**, 262 (1993); L. M. Hobbs and J. A. Thorburn, *ibid.* **491**, 772 (1997); V. V. Smith, D. L. Lambert, and P. E. Nissen, *ibid.* **506**, 405 (1998); R. Cayrel *et al.*, *Astron. Astrophys.* **343**, 923 (1999).
- [27] R. D. Evans, *The Atomic Nucleus* (McGraw-Hill, New York, 1955).
- [28] R. Pfeiffer, *Z. Phys.* **208**, 129 (1968).
- [29] D. D. Faul, B. L. Berman, P. Mayer, and D. L. Olson, *Phys. Rev. Lett.* **44**, 129 (1980).
- [30] A. N. Gorbunov and A. T. Varfolomeev, *Phys. Lett.* **11**, 137 (1964).
- [31] J. D. Irish, R. G. Johnson, B. L. Berman, B. J. Thomas, K. G. McNeill, and J. W. Jury, *Can. J. Phys.* **53**, 802 (1975).
- [32] C. K. Malcom, D. V. Webb, Y. M. Shin, and D. M. Skopik, *Phys. Lett.* **47B**, 433 (1973).
- [33] Yu. M. Arkatov, P. I. Vatsset, V. I. Voloshchuk, V. A. Zolenko, I. M. Prokhorets, and V. I. Chimil', *Yad. Fiz.* **19**, 1172 (1974) [*Sov. J. Nucl. Phys.* **19**, 598 (1974)].
- [34] B. L. Berman, *At. Data Nucl. Data Tables* **15**, 319 (1975).
- [35] M. Kawasaki and T. Moroi, *Astrophys. J.* **452**, 506 (1995).
- [36] J. D. Jackson, *Classical Electrodynamics* (Wiley, New York, 1975).
- [37] J. A. Koepke and R. E. Brown, *Phys. Rev. C* **16**, 18 (1977).
- [38] D.J. Fixsen *et al.*, *Astrophys. J.* **473**, 576 (1996).
- [39] R. Kallosh, L. Kofman, A. Linde, and A. V. Proeyen, *Phys. Rev. D* **61**, 103503 (2000).
- [40] G. F. Giudice, A. Riotto, and I. Tkachev, *J. High Energy Phys.* **08**, 009 (1999).
- [41] D. H. Lyth, *Phys. Lett. B* **476**, 356 (2000).