

Characterization of *Asparagus officinalis* hinokiresinol synthase

Masaomi Yamamura

Research Institute for Sustainable Humanosphere, Kyoto University

Norlignans, which have a diphenylpentane (C6-C5-C6) structure, were found in coniferous trees and some monocotyledonous plants including *Asparagus officinalis* [1,2]. Their biosynthesis is of interest especially in relation to heartwood formation, a metabolic event specific to woody plants, because norlignans are deposited specifically in tree heartwood regions. This implies that the molecular mechanisms can be a clue to help us elucidate mechanisms for metabolic events specific to woody plants.

Hinokiresinol is the simplest norlignan compound, it is therefore a good target for the study of norlignan biosynthesis. Hinokiresinol has two geometrical (*E*)- and (*Z*)-isomers, which are phytoalexins of *Cryptomeria japonica* (Japanese cedar) and *A. officinalis*, respectively. Recently, biosynthetic processes and enzymatic formation of (*E*)- and (*Z*)-hinokiresinols were demonstrated using *C. japonica* and *A. officinalis* cells, respectively [2-4].

In the present study, cDNAs encoding *A. officinalis* (*Z*)-hinokiresinol synthase were cloned for the first time and their recombinant proteins were characterized biochemically.

References

- [1] T. Umezawa: *Regul. Plant Growth Dev.*, 2001, 36, 57.
- [2] S. Suzuki, T. Umezawa, and M. Shimada: *J. Chem. Soc. Perkin.*, 1, 2001, 3252.
- [3] S. Suzuki, T. Nakatsubo, T. Umezawa, and M. Shimada: *Chem. Commun.*, 2002, 1088.
- [4] S. Suzuki, M. Yamamura, T. Nakatsubo, T. Hattori, M. Shimada, and T. Umezawa: *Chem. Commun.*, 2004, 2838.