

De Novo Synthesis of Veratryl Alcohol by *Coriolus versicolor**

Shingo KAWAI**, Toshiaki UMEZAWA**,
and Takayoshi HIGUCHI**

(Received September 1, 1986)

Abstract—Veratryl alcohol was found in ligninolytic culture of *Coriolus versicolor*. The structure of veratryl alcohol synthesized *de novo* was confirmed in comparison with ¹H-NMR spectra of the authentic dimethoxybenzyl alcohols (veratryl alcohol and its isomers).

1. Introduction

Lignin biodegradation has been greatly elucidated in recent years¹⁻⁵). Ligninolytic enzyme (lignin peroxidase, ligninase) was purified from the culture filtrate of *Phanerochaete chrysosporium*⁶⁻⁹) and the reaction mechanism of lignin peroxidase *via* aryl cation radical was proposed^{10,11}). This enzyme activity was enhanced by the addition of veratryl alcohol, a secondary metabolite of *P. chrysosporium*¹²).

We previously reported the degradation of non-phenolic β -O-4 lignin substructure model compounds in ligninolytic culture of *Coriolus versicolor* and suggested that a similar lignin peroxidase is excreted by *C. versicolor*¹³⁻¹⁵). In the present paper we report *de novo* synthesis of veratryl alcohol by *C. versicolor* and discuss the role of veratryl alcohol in lignin biodegradation.

2. Materials and Methods

2.1 Culture Conditions and Extraction

Coriolus versicolor Ps4a was maintained on 2% malt agar slants. Experimental culture (20 ml in 300 ml-Erlenmeyer flasks) were inoculated with a small mycelial mat from the slant and grown without agitation at 30°C. The culture medium was prepared as described previously¹³).

The 7-day-old cultures (28 cultures) were flushed with sterile oxygen and incubated under the same conditions for 3 days. The whole cultures were combined, acidified with 1N HCl to pH 2 and extracted with 1 liter of ethyl acetate. The

* A part of this paper was presented at the 29th symposium on lignin in Tokyo, Oct. 1984.

** Research Section of Lignin Chemistry.

organic layer was washed with saturated NaCl solution, dried over anhydrous Na_2SO_4 and concentrated to dryness.

2.2 Syntheses of Authentic Compounds

2,3-Dimethoxybenzyl alcohol was prepared from 2-hydroxy-3-methoxybenzaldehyde (*o*-vanillin, Nakarai Chemicals Ltd.) *via* the following two steps; (i) methyl iodide/ K_2CO_3 in DMF at room temperature, and (ii) NaBH_4 in methanol at 0°C .

$^1\text{H-NMR}$ (CDCl_3) δ (ppm); 3.87 (3H, s, $-\text{OCH}_3$), 3.89 (3H, s, $-\text{OCH}_3$), 4.70 (2H, s, $-\text{CH}_2-$), 6.82–7.12 (3H, m, aromatic-H).

2,4-Dimethoxybenzyl alcohol was prepared from 2,4-dihydroxybenzaldehyde (β -resorcyraldehyde, Nakarai Chemicals Ltd.) *via* the following two steps; (i) methyl iodide/ K_2CO_3 in DMF at room temperature, and (ii) NaBH_4 in methanol at 0°C .

$^1\text{H-NMR}$ (CDCl_3) δ (ppm); 3.80 (3H, s, $-\text{OCH}_3$), 3.83 (3H, s, $-\text{OCH}_3$), 4.60 (2H, s, $-\text{CH}_2-$), 6.40–6.46 (2H, m, aromatic- $\text{H}_{3,5}$), 7.16 (1H, d, $J=9.0$, aromatic- H_6).

2,5-Dimethoxybenzyl alcohol was prepared from 2,5-dimethoxybenzaldehyde (Nakarai Chemicals Ltd.) by reduction with NaBH_4 in methanol at 0°C .

$^1\text{H-NMR}$ (CDCl_3) δ (ppm); 3.77 (3H, s, $-\text{OCH}_3$), 3.81 (3H, s, $-\text{OCH}_3$), 4.65 (2H, s, $-\text{CH}_2-$), 6.76–6.90 (3H, m, aromatic-H).

3,4-Dimethoxybenzyl alcohol (veratryl alcohol) was commercially available (Tokyo Chemical Industry Co., Ltd.).

$^1\text{H-NMR}$ (CDCl_3) δ (ppm); 3.87 (3H, s, $-\text{OCH}_3$), 3.88 (3H, s, $-\text{OCH}_3$), 4.61 (2H, s, $-\text{CH}_2-$), 6.78–6.94 (3H, m, aromatic-H).

2.3 Instrument

$^1\text{H-NMR}$ spectra were obtained with a Varian XL-200 FT-NMR spectrometer (200 MHz) using tetramethylsilane as an internal standard. Chemical shifts and coupling constants are given in δ values (ppm) and Hz, respectively.

3. Results and Discussion

The extracts were submitted to TLC (Kiesel gel 60, F_{254} , Merck, developing solvent: CH_2Cl_2). Veratryl alcohol was isolated and its structure was identified by $^1\text{H-NMR}$. The $^1\text{H-NMR}$ spectra of the metabolic veratryl alcohol and authentic compounds are shown in Fig. 1. Possibility of 3,5-dimethoxybenzyl alcohol is ruled out, because the protons of the two methoxyl groups of the metabolite have different chemical shifts in $^1\text{H-NMR}$ spectrum, while the chemical shifts of methoxyl groups of 3,5-dimethoxybenzyl alcohol are identical. From the $^1\text{H-NMR}$ spectra shown in Fig. 1, it is clear that the metabolic product (A) is veratryl alcohol (B) and not other isomers (C–E).

Russell *et al.*¹⁶⁾ found veratraldehyde in a culture of *C. versicolor*. *De novo* syn-




Fig. 1. $^1\text{H-NMR}$ spectra of metabolic veratryl alcohol (A) and authentic compounds (B-E).

thesis of veratryl alcohol by *P. chrysosporium* was reported previously by Lundquist and Kirk¹⁷⁾. Afterwards many papers were published in relation to physiological and biochemical role of veratryl alcohol in lignin biodegradation by *P. chrysosporium*. Shimada *et al.*¹⁸⁾ reported the biosynthesis of veratryl alcohol in relation to lignin degradation by *P. chrysosporium*. It was demonstrated that addition of veratryl alcohol to the culture of *P. chrysosporium* increased the ligninolytic activity and the production of lignin peroxidase in the culture^{12,19)}. It was further reported that the oxidation of monomethoxylated aromatic monomers²⁰⁾ and 2-keto-4-thiomethyl butyric acid (KTBA)²¹⁾ by lignin peroxidase of *P. chrysosporium* was enhanced by veratryl alcohol.

Our previous investigations¹³⁻¹⁵⁾ showed that the main degradation products of non-phenolic β -O-4 lignin substructure model compounds by *C. versicolor* were similar to those obtained by lignin peroxidase of *P. chrysosporium*^{6-9,22,23)}, suggesting that a similar lignin peroxidase is excreted by *C. versicolor*. These results also suggest that veratryl alcohol enhances the ligninolytic activity and the production of the lignin peroxidase by *C. versicolor*.

Acknowledgement

This research was partly supported by a Grant-in-Aid for Scientific research (No. 59760124) from the Ministry of Education of Japan.

References

- 1) C.-L. CHEN and H.-m. CHANG: "Biosynthesis and Biodegradation of Wood Components", (T. HIGUCHI ed) Academic Press, Florida, p.535 (1985).
- 2) T. HIGUCHI: *ibid.* p. 557.
- 3) T.K. KIRK and M. SHIMADA: *ibid.* p. 579.
- 4) M.S.A. LEISOLA and A. FIECHTER: "Advances in Biotechnological Processes vol. 5", (A. MIZRAHI and A.L. VAN WEZAL eds.) Alan R. Liss, New York, p. 59 (1985).
- 5) P.J. HARVEY, H.E. SHOEMAKER and J.M. PALMER: *Ann. Proc. Phytochem. Soc. Eur.*, **26**, 249 (1985).
- 6) M. TIEN and T.K. KIRK: *Science*, **221**, 661 (1983).
- 7) M. TIEN and T.K. KIRK: *Proc. Natl. Acad. Sci. USA*, **81**, 2280 (1984).
- 8) J.K. GLENN, M.A. MORGAN, M.B. MAYFIELD, M. KUWAHARA and M.H. GOLD: *Biochem. Biophys. Res. Commun.*, **114**, 1077 (1983).
- 9) M.H. GOLD, M. KUWAHARA, A.A. CHIU and J.K. GLENN: *Arch. Biochem. Biophys.*, **234**, 353 (1984).
- 10) P.J. KERSTEN, M. TIEN, B. KALYANARAMAN and T.K. KIRK: *J. Biol. Chem.*, **260**, 2609 (1985).
- 11) K.E. HAMMEL, M. TIEN, B. KALYANARAMAN and T.K. KIRK: *J. Biol. Chem.*, **260**, 8348 (1985).
- 12) B.D. FAISON and T.K. KIRK: *Appl. Environ. Microbiol.*, **49**, 299 (1985).
- 13) S. KAWAI, T. UMEZAWA and T. HIGUCHI: *Agric. Biol. Chem.*, **49**, 2325 (1985).
- 14) S. KAWAI, T. UMEZAWA and T. HIGUCHI: *Appl. Environ. Microbiol.*, **50**, 1505 (1985).
- 15) S. KAWAI, T. UMEZAWA and T. HIGUCHI: *FEBS Lett.*, in press.
- 16) J.D. RUSSELL, M.E.K. HENDERSON and V.C. FARMER: *Biochim. Biophys. Acta*, **52**, 565 (1961).
- 17) K. LUNDQUIST and T.K. KIRK: *Phytochem.*, **17**, 1676 (1978).
- 18) M. SHIMADA, F. NAKATSUBO, T.K. KIRK and T. HIGUCHI: *Arch. Microbiol.*, **129**, 321 (1981).
- 19) M.S.A. LEISOLA, D.C. ULMER, R. WALDNER and A. FIECHTER: *J. Biotechnol.*, **1**, 331 (1984).
- 20) P.J. HARVEY, H.E. SHOEMAKER and J.M. PALMER: *FEBS Lett.*, **195**, 242 (1986).
- 21) V. RANGANATHAN, K. MIKI and M.H. GOLD: *Arch. Biochem. Biophys.*, **241**, 304 (1985).
- 22) T. UMEZAWA, M. SHIMADA, T. HIGUCHI and K. KUSAI: *FEBS Lett.*, **205**, 287 (1986).
- 23) T. UMEZAWA and T. HIGUCHI: *FEBS Lett.*, **205**, 293 (1986).