

Title	On cubic polynomials with a parabolic fixed point of a capture type(Complex Dynamics and its Related Fields)
Author(s)	Yazawa, Hikaru
Citation	数理解析研究所講究録 (2006), 1494: 99-105
Issue Date	2006-05
URL	<a href="http://hdl.handle.net/2433/58294">http://hdl.handle.net/2433/58294</a>
Right	
Type	Departmental Bulletin Paper
Textversion	publisher

# On cubic polynomials with a parabolic fixed point of a capture type

学習院 大学院 自然科学研究科 数学専攻  
谷澤 晃 (Hikaru Yazawa)  
Faculty of Science, Gakushuin University

## Abstract

We consider the location of each critical point of a cubic polynomial map with a parabolic fixed point. We show that, for any given number of iterations, there exists a cubic polynomial map with a parabolic fixed point such that the immediate parabolic basin contains just one of the critical points and the image of another critical point under the specified number of iterations.

## 1 Introduction

Let  $f$  be any cubic polynomial. If  $f$  has a parabolic fixed point  $\alpha$ , then a cycle of Fatou components of  $f$  is called the immediate parabolic basin for  $\alpha$  if the cycle contains a parabolic petal for  $\alpha$ .

Roughly speaking, in this note we consider the dynamically location of each critical points of  $f$  with a parabolic fixed point whose basin contains both the critical points. We denote by  $c_0$  and  $c_1$  the critical points of  $f$ . Using the Haissinsky pinching deformation, we prove the following result:

**Theorem 1.1.** For any positive integer  $n$ , there exists a cubic polynomial map  $f$  with a parabolic fixed point such that the immediate parabolic basin contains  $c_0$  and  $f^{on}(c_1)$ , and does not contain  $f^{ok}(c_1)$  for any integer  $k$  with  $0 \leq k < n$ .

Now, suppose that  $f$  has a parabolic fixed point, and the parabolic basin contains  $c_0$  and  $c_1$ . By analogy with Milnor [3], we shall define the types of this parabolic fixed point. For  $j = 0, 1$ , we denote by  $U_j$  the Fatou component which contains  $c_j$ . Without loss of generality, we may assume that  $U_0$  is contained in the immediate basin of the parabolic fixed point. Following from [3], there exist four possibilities as follows.

**Case 1:** The Fatou component is adjacent, i.e.,  $U_0 = U_1$ .

**Case 2:** The Fatou component is bitransitive. Namely,  $U_0 \neq U_1$ , and moreover there exist the smallest positive integers  $p, q > 0$  such that  $f^{\circ p}(U_0) = U_1$  and  $f^{\circ q}(U_1) = U_0$ .

**Case 3:** The immediate parabolic basin captures  $U_1$ . Namely, the immediate parabolic basin does not contain  $U_1$ , but  $f^{\circ k}(U_1)$  for some integer  $k \geq 1$ .

**Case 4:** Each of  $U_0$  and  $U_1$  is contained in the disjoint cycle of the immediate parabolic basin. Namely,  $U_0$  and  $U_1$  is contained in the immediate parabolic basin, and it follows that  $f^{\circ n}(U_0) \cap f^{\circ m}(U_1) = \emptyset$  for any integers  $n, m \geq 0$ .

We define the types of the parabolic fixed point  $\alpha$  as follows:

**Definition 1.2.** In Case 1, 2, 3 or 4, we say that  $\alpha$  is a parabolic fixed point of an *adjacent*, *bitransitive*, *capture*, or *disjoint* type, respectively.

We will consider the type of the parabolic fixed point a the cubic polynomial map obtained by the Haissinsky pinching deformation, which is illustrated in the next section.

## 2 The Haissinsky Pinching deformation

Suppose that  $f$  is any cubic polynomial map with an attracting fixed point  $\alpha$ . Let  $B_f(\alpha)$  be the attracting basin for  $\alpha$ . We consider the Haissinsky pinching deformation of  $f$  defined by pinching curves in  $B_f(\alpha)$ .

Following from [1], for any integer  $q \geq 1$ , there exist a smooth open arc  $\gamma$  and a neighborhood  $U \subset B_f(\alpha)$  of  $\gamma$  satisfying the following conditions.

- $\bar{\gamma} \setminus \gamma$  consists of the attracting fixed point  $\alpha$  and a repelling periodic point  $\beta$  of period  $q$ .
- $f^{\circ q}(\gamma) = \gamma$ ,  $f^{\circ q}(U) = U$ , and  $f^{\circ q}|_U$  is univalent.
- $f^{\circ n}(U) \cap f^{\circ m}(U) = \emptyset$  for any  $0 \leq n < m < q$ .
- There exist a number  $\sigma > 0$  and a conformal map  $\Phi_\sigma : U \rightarrow \{|z| < \pi\}$  such that  $\Phi_\sigma \circ f^{\circ q}(z) = \Phi_\sigma(z) + \sigma$  for all  $z \in U$ .

We call the union  $S := \bigcup_{k \geq 0} f^{\circ -k}(\bar{\gamma})$  the *support of pinching*, and define  $S_0 := \bigcup_{k \geq 0} f^{\circ k}(\bar{\gamma})$ . It follows from [1] that we have a sequence of quasiconformal maps  $(h_t)_{t \geq 0}$  satisfying the following conditions.

- $h_t$  converges uniformly on  $\widehat{\mathbb{C}}$  to a local quasiconformal map  $h_\infty$  on  $\widehat{\mathbb{C}} \setminus S$ .
- $f_t := h_t \circ f \circ h_t^{-1}$  converges uniformly on  $\widehat{\mathbb{C}}$  to a cubic polynomial  $f_\infty$ .
- $h_\infty(\alpha)$  is a parabolic fixed point of  $f_\infty$ .
- $h_\infty(S_0) = h_\infty(\alpha)$ .

For further details, see [1] or [2].

### 3 Proof of Theorem 1.1

We first prove the following lemma needed later.

**Lemma 3.1.** Let  $n$  be any positive integer, and let  $\lambda$  be any complex number in  $\mathbb{D} \setminus \{0\}$ . Then there exists a cubic polynomial  $f$ , with  $f^{on}(c_1) = c_0$ , such that  $f$  has an attracting fixed point of multiplier  $\lambda$  whose attracting basin is simply connected.

**Proof.** Consider a monic and centered cubic polynomial

$$P_{A,B}(z) = z^3 - 3Az + \sqrt{B}, \quad (A, B) \in \mathbb{C}^2.$$

Suppose that  $P_{A,B}$  has a fixed point of multiplier  $\lambda$ . Then the fixed point is  $\alpha_{A,\lambda} := \sqrt{A + \lambda/3}$ , and hence,  $P_{A,B}$  is affine conjugate to the cubic polynomial map

$$Q_{A,\lambda}(z) = z^3 + 3\alpha_{A,\lambda}z^2 + \lambda z$$

with critical points  $c_{A,\lambda}^\pm := -\alpha_{A,\lambda} \pm \sqrt{A}$ .

Suppose that  $\lambda \in (-1, 0)$ , and the parameter  $A$  is any real number  $> -\lambda/3$  such that the attracting basin for zero is simply connected.

For each integer  $k \geq 0$ , we denote by  $z_{A,\lambda}(k)$  the unique point on  $\mathbb{R}_+$  such that  $Q_{A,\lambda}^k(z_{A,\lambda}(k)) = c_{A,\lambda}^+$ . For any integer  $k > 0$  and for any real number  $A'$  with  $A' > A$ , we have  $z_{A,\lambda}(k) < z_{A,\lambda}(k+1)$  and  $z_{A,\lambda}(k) > z_{A',\lambda}(k)$ . Thus since  $Q_{A,\lambda}(c_{A,\lambda}^-) \rightarrow +\infty$  as  $A \rightarrow +\infty$ , for any integer  $n > 0$  there exists a real number  $A$  such that  $Q_{A,\lambda}^{on}(c_{A,\lambda}^-) = c_{A,\lambda}^+$ .

Let  $\lambda'$  be any complex number in  $\mathbb{D} \setminus \{0\}$ . Then it follows from [5] that there exists a quasiconformal map  $h$  such that the cubic polynomial map  $g := h \circ Q_{A,\lambda} \circ h^{-1}$  has an attracting fixed point with multiplier  $\lambda'$ .  $\square$

We use the Haissinsky pinching deformation of  $f$  obtained from this lemma.

**Proof of Theorem 1.1.** Without loss of generality, we may assume that  $f(z) = z^3 + 3\alpha_{A,\lambda}z^2 + \lambda z$ ,  $c_0 = c_{A,\lambda}^+$  and  $c_1 = c_{A,\lambda}^-$ .

Suppose that  $\lambda$  is any real number with  $-1 < \lambda < 0$ , and  $A$  is a real number  $> -\lambda/3$  such that the attracting basin for zero is simply connected. Recall that  $B_f(0)$  is the attracting basin for zero. Let  $\varphi_f$  be the Koenigs map such that  $\varphi_f(0) = 0$ , and  $\varphi_f(z) = \lambda z$  for all  $z \in B_f(0)$ . We may assume that  $\varphi_f(c_0) = 1$ .

Define the half-line  $\hat{\gamma} := i\mathbb{R}^+$ , so that  $\hat{\gamma}$  is periodic of period two under the iterates of the map  $L(z) := \lambda z$ . We denote by  $\gamma$  the connected component of the preimage of  $\hat{\gamma}$  under  $\varphi_f$  whose closure contains zero. Thus, we have the support of pinching  $S := \bigcup_{k \geq 0} f^{-k}(\bar{\gamma})$ , and denote by  $f_\infty$  the limit of the Haissinsky pinching deformation of  $f$  defined by  $S$ .

Let  $n$  be any positive integer. From Lemma 3.1, we have a parameter  $A$  such that  $f^{\circ n}(c_1) = c_0$ . For each integer  $k \geq 1$ , we denote by  $\alpha(k)$  the point on  $\mathbb{R}_+$  such that  $f^{\circ k}(\alpha(k)) = 0$ , and by  $S_{\alpha(k)}$  the connected component of  $S$  which contains  $\alpha(k)$ .

At first consider the case  $n \geq 2$ . Since for each integer  $k \geq 1$  the component  $S_{\alpha(k)}$  separates the origin and  $f^{\circ k}(c_1)$ , it follows that  $f_\infty$  has a parabolic fixed point of a capture type.

Next, consider the case  $n = 1$ . Since no connected component of  $S$  separates the origin and  $c_1$ , it follows that  $f_\infty$  has a parabolic fixed point of a bitransitive type.

In order to obtain a polynomial with a parabolic fixed point of a capture type, we will use the Branner-Hubbard deformation of  $f$  obtained by wringing the almost complex structure on the attracting basin for zero (cf. [5]). In particular, we consider the Branner-Hubbard deformation which does not change the multiplier of the origin.

Let  $s = 1 + 2\pi i / \log \lambda$ , and let  $l$  be the quasi-conformal map defined as  $l(z) := z|z|^{s-1}$ .

Recall that  $\varphi_f$  is the Koenigs map defined on  $B_f(0)$ . We define the holomorphic map  $\psi_f : \mathbb{D} \rightarrow \mathbb{C}$  as the inverse map of  $\varphi_f$  such that  $\psi_f(0) = 0$ .

Let  $\sigma_0$  be the standard almost complex structure of  $\hat{\mathbb{C}}$ , and let  $\sigma$  be the almost complex structure defined as follows:

$$\sigma = \begin{cases} \sigma_0 & \text{on } \hat{\mathbb{C}} \setminus B_f(0) \\ (l \circ \varphi_f)^*(\sigma_0) & \text{on } \psi_f(\mathbb{D}) \\ (l \circ \varphi_f \circ f^{\circ k})^*(\sigma_0) & \text{on } f^{-k}(\psi_f(\mathbb{D})) \setminus f^{-k+1}(\psi_f(\mathbb{D})), \end{cases} \quad (1)$$

where  $k$  is an integer  $\geq 1$ .

From the Measurable Riemann Mapping Theorem, we obtain the quasi-conformal map  $h$  such that  $h^*\sigma_0 = \sigma$ . Suppose that  $h(0) = 0$ ,  $h(1) = 1$

and  $h(\infty) = \infty$ . Then, we obtain a cubic polynomial map  $g = h \circ f \circ h^{-1}$  with the attracting fixed point zero. It follows from [5] that the multiplier is  $g'(0) = h(\lambda) = \lambda |\lambda|^{s-1} = \lambda$ , and that the Koenigs map  $\varphi_g = l \circ \varphi_f \circ h^{-1}$ .

Following from the argument similar to the above discussion, we define  $S' \subset B_g(0)$  as the support of the pinching deformation, and denote by  $g_\infty$  the limit of the pinching deformation of  $g$  defined by the support  $S'$ .

There exists a cycle of connected components of  $B_f(0) \setminus h^{-1}(S')$  under the iterates of  $f$ ,

If  $c_1$  is not contained in this cycle, then one of the critical points of  $g_\infty$  is not contained in the immediate parabolic basin of  $g_\infty$ .

We consider the inverse image of  $i\mathbb{R}$  under  $\varphi \circ h^{-1}$ . We introduce a preliminary definition as follows. For any point  $z$  of the backward orbit of the origin, we denote by  $D_f(z; r)$  the connected component of the set  $\{w : |\varphi_f(w)| < r\}$  which contains the point  $z$ .

Since  $f$  has no critical point in the open set  $D_f(0; |\lambda|^{-1})$  except  $c_0$ , it follows that  $f$  maps  $D_f(0; |\lambda|^{-1}) \setminus \{c_0\}$  to  $D_f(0; 1) \setminus \{c_0\}$  in two-to-one correspondence. Thus  $f$  has the unique preimage  $\alpha'$  of the origin such that  $\alpha' \neq 0$  and  $\alpha' \in D_f(0; |\lambda|^{-1}) \setminus \{c_0\}$ .

We extend  $\psi_f$  to the conformal map  $\psi_{f,0}$  defined on  $\mathbb{D}(0; |\lambda|^{-1}) \setminus [1, |\lambda|^{-1})$  to a subset of  $D_f(0; |\lambda|^{-1})$ . Moreover, we define  $\psi_{f,1}$  as the conformal map defined on  $\mathbb{D}(0; |\lambda|^{-1}) \setminus [1, |\lambda|^{-1})$  such that  $\varphi_f \circ \psi_{f,1} \equiv \text{identity map}$  and  $\psi_{f,1}(0) = \alpha'$ .

The end points of the image of the set  $\{yi \mid -|\lambda|^{-1} < y < |\lambda|^{-1}\}$  under  $\psi_{f,0} \circ h^{-1}$  is contained in the boundary of  $\psi_{f,1}(\mathbb{D}(0; |\lambda|^{-1}) \setminus [1, |\lambda|^{-1}))$ . Hence, the connected component of the preimage of  $i\mathbb{R}$  under  $\varphi_f \circ h^{-1}$  which contains zero passes through the boundary of  $\psi_{f,1}(\mathbb{D}(0; |\lambda|^{-1}) \setminus [1, |\lambda|^{-1}))$ , and does not separate  $c_0$  and  $c_1$ . On the other hand, the connected component of the preimage of  $i\mathbb{R}$  under  $\varphi_f \circ h^{-1}$  which contains  $\alpha'$  separates  $c_0$  and  $c_1$ . Therefore, the cycle of the Fatou components of  $g$  does not contain one of the critical points of  $g$ , and hence  $g_\infty$  has a parabolic fixed point of a capture type.  $\square$

## 4 Notes

Consider the family of cubic polynomials  $P_{A,B}(z) := z^3 - 3Az + \sqrt{B}$  with  $P_{A,B}(-\sqrt{A}) = \sqrt{A}$ . We have  $B = A(1 - 2A)^2$ . The connectedness locus of the family of  $P_{A,A(1-2A)^2}(z) = z^3 - 3Az + \sqrt{A} - 2A\sqrt{A}$ ,  $A \in \mathbb{C}$ , is showed in Figure 2.




Figure 1: Sketch for the pinching curves.




Figure 2: The connectedness locus of the family of cubic polynomials  $P_{A, A(1-2A)^2}$ ,  $A \in \mathbb{C}$ .

$P_{A,A(1-2A)^2}$  is affine conjugate to the cubic polynomial map

$$F_A(z) := (P_{A,A(1-2A)^2}(\sqrt{A}z + \sqrt{A}) - \sqrt{A})/\sqrt{A} = Az^3 + 3Az^2 - 4A.$$

Suppose that  $0 < |A| < 1/4$ . Then the map  $F_A$  satisfies the inequality  $|F_A(z) + 4A| < |4A|$ , that is,  $F_A$  maps the disk of radius  $|F_A(0)|$  centered at  $F_A(0)$  into itself. Hence  $F_A$  has an attracting fixed point in the disk.

Let  $\alpha_A$  be the attracting fixed point.

**Proposition 4.1.** If  $A$  turns around the origin once, then the multiplier of the attracting fixed point of  $F_A$  turns around the origin twice.

**Proof.** Let  $D$  be the disk of radius  $|F_A(0)|$  centered at  $F_A(0)$ . If  $A$  turns around the origin once, then the center of  $D$  turns around the origin once.

Set  $0 < r < 1/4$ ,  $\theta \in [0, 1]$ , and  $A = re^{2\pi i\theta}$ . Since the radius of  $D$  is the constant  $|F_A(0)|$ , the attracting fixed point  $\alpha_A$  also turns around the origin once. Thus the multiplier  $F'_A(\alpha_A) = 3A\alpha_A(\alpha_A + 2)$  turns around the origin twice.  $\square$

## References

- [1] P. Haissinsky, *Pincement de polynômes*, Comment. Math. Helv. **77** (2002), no. 1, 1–23.
- [2] T. Kawahira, *Note on dynamically stable perturbations of parabolics*, RIMS Kokyuroku, 1447 (2005), pp 90 - 107.
- [3] J. Milnor, *Remarks on iterated cubic maps*, Experiment. Math. **1** (1992), no. 1, 5–24.
- [4] J. Milnor, *Dynamics in one complex variable*, Friedr. Vieweg & Sohn, Braunschweig, 2000.
- [5] C. Petersen, Tan Lei, *Branner-Hubbard motions and attracting dynamics*, preprint.
- [6] L. Tan, *On pinching deformations of rational maps*, Ann. Sci. École Norm. Sup. (4) **35** (2002), no. 3, 353–370.