TITLE:

An algorithm of computing b -functions

AUTHOR(S):

Oaku, Toshinori

CITATION:

Oaku, Toshinori. An algorithm of computing b -functions. 数理解析研究所講究録 1996, 941: 52-56

ISSUE DATE:

1996-03

URL:

<http://hdl.handle.net/2433/60136>

RIGHT:

8.

An algorithm of computing b -functions

大阿久 俊則 (横浜市大・理)

8.1 Introduction

Let $f(x) \in K[x] = K[x_1, \dots, x_n]$ be a polynomial with coefficients in a field K of characteristic zero. Let us denote by

$$\hat{\mathcal{D}}_n := K[[x_1, \dots, x_n]](\partial_1, \dots, \partial_n)$$

the ring of differential operators with formal power series coefficients with $\partial_i = \partial/\partial x_i$ and $\partial = (\partial_1, \dots, \partial_n)$. (If K is a subfield of the field $\mathbf{C}$ of complex numbers, then we can use the ring $\mathcal{D}_n$ of differential operators with convergent power series coefficients instead of $\hat{\mathcal{D}}_n$. This makes no difference in the definition below.) Let s be a parameter. Then the (local) b -function (or the Bernstein-Sato polynomial) $b_f(s)$ associated with $f(x)$ is the monic polynomial of the least degree $b(s) \in K[s]$ satisfying

$$P(s, x, \partial)f(x)^{s+1} = b(s)f(x)^s$$

with some $P(s, x, \partial) \in \hat{\mathcal{D}}_n[s]$.

We present an algorithm of computing the b -function $b_f(s)$ for an arbitrary $f(x) \in K[x]$. A system *Kan* of N. Takayama [T2] is available for actual execution of our algorithm.

An algorithm of computing $b_f(s)$ was first given by M. Sato et al. [SKKO] when $f(x) \in \mathbf{C}[x]$ is a relative invariant of a prehomogeneous vector space. J. Briançon et al. [BGMM], [M] have given an algorithm of computing $b_f(s)$ for $f(x) \in \mathbf{C}\{x\}$ with isolated singularity. Also note that T. Yano [Y] worked out many interesting examples of b -functions systematically.

8.2 Algorithm

Notation

- K : a field of characteristic 0;
- $A_{n+1} := K[t, x_1, \dots, x_n] \langle \partial_t, \partial_1, \dots, \partial_n \rangle$ ($\partial_t := \partial/\partial t$, $\partial_i = \partial/\partial x_i$)
- $\prec_F$: a (total) order on $\mathbb{N}^{2n+2}$ with $\mathbb{N} := \{0, 1, 2, \dots\}$ that satisfies the following conditions:
 - (A-1) $\alpha \succ_F \beta \implies \alpha + \gamma \succ_F \beta + \gamma$ ($\forall \alpha, \beta, \gamma \in \mathbb{N}^{2n+2}$);
 - (A-3) $\nu - \mu > \nu' - \mu' \implies (\mu, \nu, \alpha, \beta) \succ_F (\mu', \nu', \alpha', \beta')$ ($\forall \mu, \nu, \mu', \nu' \in \mathbb{N}$, $\forall \alpha, \beta, \alpha', \beta' \in \mathbb{N}^n$);
 - (A-4) $(\mu, \mu, \alpha, \beta) \succeq_F (0, 0, 0, 0)$ ($\forall \mu \in \mathbb{N}$, $\forall \alpha, \beta \in \mathbb{N}^n$),
 where $(\mu, \nu, \alpha, \beta)$ corresponds to the ‘monomial’ $t^\mu x^\alpha \partial_t^\nu \partial^\beta$. Note that $\succ_F$ does not satisfy
 - (A-2) $\alpha \succeq_F 0$ ($\forall \alpha \in \mathbb{N}^{2n+2}$).

For each integer m , define a K -subspace of A_{n+1} by

$$F_m(A_{n+1}) := \{P = \sum_{\mu, \nu, \alpha, \beta} a_{\mu, \nu, \alpha, \beta} t^\mu x^\alpha \partial_t^\nu \partial^\beta \in A_{n+1} \mid a_{\mu, \nu, \alpha, \beta} = 0 \text{ if } \nu - \mu > m\}.$$

If $P \neq 0$, its F -order $\text{ord}_F(P)$ is defined as the minimum integer m such that $P \in F_m(A_{n+1})$.

Then

$$\hat{\sigma}(P) = \hat{\sigma}_m(P) := \sum_{\nu - \mu = m} a_{\mu, \nu, \alpha, \beta} t^\mu x^\alpha \partial_t^\nu \partial^\beta$$

is called the *formal symbol* of P . We define $\psi(P)(s) \in A_n[s]$ by

$$\hat{\sigma}_0(t^m P) = \psi(P)(t \partial_t) \quad \text{if } m \geq 0,$$

$$\hat{\sigma}_0(\partial_t^{-m} P) = \psi(P)(t \partial_t) \quad \text{if } m < 0.$$

Definition 1 For $i, j, \mu, \nu, \mu', \nu' \in \mathbb{N}$, $\alpha, \beta, \alpha', \beta' \in \mathbb{N}^n$, an order $\prec_H$ on $\mathbb{N}^{2n+3}$ is defined by

$$(i, \mu, \nu, \alpha, \beta) \succ_H (j, \mu', \nu', \alpha', \beta') \iff (i > j)$$

$$\text{or } (i = j \text{ and } (\mu + \ell, \nu, \alpha, \beta) \succ_F (\mu' + \ell', \nu', \alpha', \beta'))$$

$$\text{or } (i = j, (\nu, \alpha, \beta) = (\nu', \alpha', \beta'), \mu > \mu')$$

with $\ell, \ell' \in \mathbb{N}$ s.t. $\nu - \mu - \ell = \nu' - \mu' - \ell'$, where $(i, \mu, \nu, \alpha, \beta)$ corresponds to $t^i x_0^\mu x^\alpha \partial^\beta$. This definition is independent of the choice of ℓ, ℓ' , and $\succ_H$ satisfies (A-1) and (A-2).

In the following algorithm, we also use an order $\prec$ on $\mathbb{N}^{2n+1}$ satisfying (A-1), (A-2) (with $2n+2$ replaced by $2n+1$) and

- (A-5) if $|\beta| > |\beta'|$, then $(\mu, \alpha, \beta) \succ (\mu', \alpha', \beta')$ for any $\mu, \mu' \in \mathbb{N}$ and $\alpha, \beta, \alpha', \beta' \in \mathbb{N}^n$,

where (μ, α, β) corresponds to $s^\mu x^\alpha \partial^\beta$.

Algorithm 2

Input: $f(x) \in K[x]$;

1. Let $\mathbf{G}$ be a Gröbner basis of the left ideal of $A_{n+1}[x_0]$ generated by $t - x_0 f(x)$ and $\partial_i + x_0(\partial f / \partial x_i) \partial_i$ ($i = 1, \dots, n$) with respect to $\prec_H$;
2. Compute a Gröbner basis $\mathbf{H}$ of the left ideal of $A_n[s]$ generated by $\psi(\mathbf{G}) := \{\psi(P(1)) \mid P(x_0) \in \mathbf{G}\}$ w.r.t. an order satisfying (A-1), (A-2), (A-5);
3. Let J be the ideal of $K[x, s]$ generated by $\mathbf{H} \cap K[x, s] = \{f_1(x, s), \dots, f_k(x, s)\}$;
4. Compute the monic generator $f_0(s)$ of the ideal of $K[s]$ generated by $f_1(0, s), \dots, f_k(0, s)$ by Gröbner basis or GCD computation; if $f_0(s) = 1$, then put $b(s) := 1$ and quit;
5. Compute the factorization $f_0(s) = (s - s_1)^{\mu_1} \dots (s - s_m)^{\mu_m}$ in $\overline{K}[s]$ ($\overline{K}$: the algebraic closure of K);
6. Put $\overline{J} := \overline{K}[x, s]J$.

For $i := 1$ to m do

- By computing the ideal quotient $\overline{J} : (s - s_i)^\ell$ for $\ell = \mu_i, \mu_i + 1, \dots$ repeatedly, determine the least $\ell \geq \mu_i$ such that $\overline{J} : (s - s_i)^\ell$ contains an element which does not vanish at $(x, s) = (0, s_i)$. Denote this ℓ by ℓ_i ;
7. Put $b(s) := (s - s_1)^{\ell_1} \dots (s - s_m)^{\ell_m}$;

Output: $b_f(s) := b(-s - 1) \in K[s]$;

Remark 3 A theorem of Kashiwara [K] states that the roots of $b_f(s)$ are negative rational numbers. Hence in steps 5 and 6, there is no need of field extension.

We have implemented the steps 1 and 2 of the above algorithm in Kan/sm1 [T2], and the steps 3–7 in Risa/Asir [NS]. In the following table, the timing data refer to the computation time of steps 1 and 2, which are naturally the most expensive part of our algorithm.

$f(x)$	$b_f(s)$	timing data by Kan on S-4/20
$x^3 - y^2$	$(s+1)(s+\frac{5}{6})(s+\frac{7}{6})$	0.2s
$(x^3 - y^2)^2$	$(s+1)(s+\frac{1}{12})(s+\frac{5}{12})(s+\frac{1}{2})(s+\frac{7}{12})(s+\frac{11}{12})$	0.7s
$x^5 - y^2$	$(s+1)(s+\frac{7}{10})(s+\frac{9}{10})(s+\frac{11}{10})(s+\frac{13}{10})$	0.2s
$x^5 + y^5$	$(s+1)^2(s+\frac{2}{5})(s+\frac{3}{5})(s+\frac{4}{5})(s+\frac{6}{5})(s+\frac{7}{5})(s+\frac{8}{5})$	0.8s
$x^5 + y^5 + x^3y^3$	$(s+1)^2(s+\frac{2}{5})(s+\frac{3}{5})(s+\frac{4}{5})(s+\frac{6}{5})(s+\frac{7}{5})$	180s
$x^3y + y^3 + z^2$	$(s+1)(s+\frac{35}{18})(s+\frac{31}{18})(s+\frac{29}{18})$ $\times (s+\frac{3}{2})(s+\frac{25}{18})(s+\frac{23}{18})(s+\frac{19}{18})$	5s
$x^5 + y^3 + z^2$	$(s+1)(s+\frac{59}{30})(s+\frac{53}{30})(s+\frac{49}{30})(s+\frac{47}{30})$ $\times (s+\frac{43}{30})(s+\frac{41}{30})(s+\frac{37}{30})(s+\frac{31}{30})$	7s
$x^3 + y^2z^2$	$(s+1)(s+\frac{5}{6})^2(s+\frac{7}{6})^2(s+\frac{4}{3})(s+\frac{5}{3})$	0.5s
$x^3 + y^3 - 3xyz$	$(s+1)^3(s+\frac{4}{3})(s+\frac{5}{3})$	2.5s
$x^3 + xyz$	$(s+1)^3(s+\frac{4}{3})(s+\frac{5}{3})$	0.5s
$x^4 + y^2z^2 + x^3y^3$	$(s+1)^3(s+\frac{3}{4})^2(s+\frac{5}{6})^2(s+\frac{7}{6})^2(s+\frac{5}{4})^2$ $\times (s+\frac{11}{12})(s+\frac{13}{12})(s+\frac{4}{3})(s+\frac{17}{12})(s+\frac{3}{2})$ $\times (s+\frac{19}{12})(s+\frac{5}{3})(s+\frac{7}{4})$	180s

In the above table, the last four examples have non-isolated singularities. Hence, as far as the author knows, no algorithm has been known for computing b -functions for these polynomials. See [Y, pp. 198–200] for estimates of the b -functions of $x^3 + y^2z^2$, $x^3 + y^3 - 3xyz$, $x^3 + xyz$.

Acknowledgement: The author would like to express his gratitude to Professor N. Takayama of Kobe University for kind assistance in using Kan, without which implementation of our algorithm

would have been much more difficult.

参考文献

- [Be] Bernstein, I. N.: Modules over a ring of differential operators. *Functional Anal. Appl.* **5** (1971), 89–101.
- [Bj] Björk, J.E.: *Rings of Differential Operators*. North-Holland, Amsterdam, 1979.
- [BGMM] Briançon, J., Granger, M., Maisonobe, Ph., Miniconi, M.: Algorithme de calcul du polynôme de Bernstein: cas non dégénéré. *Ann. Inst. Fourier* **39** (1989), 553–610.
- [G] Galligo, A.: Some algorithmic questions on ideals of differential operators. *Lecture Notes in Comput. Sci.* vol. 204, pp. 413–421, Springer, Berlin, 1985.
- [K] Kashiwara, M.: B -functions and holonomic systems –Rationality of roots of b -functions. *Invent. Math.* **38** (1976), 33–53.
- [M] Maisonobe, P.: D -modules: an overview towards effectivity. *Computer Algebra and Differential Equations* (ed. E. Tournier), Cambridge University Press, 1994, pp. 21–55.
- [NS] Noro, M., Shimoyama, T.: *Asir user's manual*, Edition 3.0 for Asir-950831. (<ftp://endeavor.fujitsu.co.jp>) ISIS, Fujitsu Laboratories Limited, 1995
- [O1] Oaku, T.: Algorithms for finding the structure of solutions of a system of linear partial differential equations. *Proceedings of International Symposium on Symbolic and Algebraic Computation* (eds J. Gathen, M. Giesbrecht), pp. 216–223, ACM, New York, 1994.
- [O2] Oaku, T.: Algorithmic methods for Fuchsian systems of linear partial differential equations. *J. Math. Soc. Japan* **47** (1995), 297–328.
- [O3] Oaku, T.: An algorithm of computing b -functions. Preprint.
- [SKKO] Sato, M., Kashiwara, M., Kimura, T., Oshima, T.: Micro-local analysis of prehomogeneous vector spaces. *Invent. Math.* **62** (1980), 117–179.
- [T1] Takayama, N.: An approach to the zero recognition problem by Buchberger algorithm. *J. Symbolic Comput.* **14** (1992), 265–282.
- [T2] Takayama, N.: Kan: A system for computation in algebraic analysis. <http://www.math.s.kobe-u.ac.jp>, 1991—.
- [Y] Yano, T.: On the theory of b -functions. *Publ. RIMS, Kyoto Univ.* **14** (1978), 111–202.