

A FUZZY RELATIONAL EQUATION IN DYNAMIC FUZZY SYSTEMS

M. KURANO(Chiba University), M. YASUDA(Chiba University),
J. NAKAGAMI(Chiba University) & Y. YOSHIDA(Kitakyushu University)

Abstract : For a dynamic Fuzzy system, the fundamental method is to analysis its recursive relation of the fuzzy states. It is similar as the Bellman equation is the important tool in the dynamic programming. Here we will consider the existence and the uniqueness of solution of the fuzzy relational equation. Two examples which satisfies our conditions, are given to illustrate the results.

1 Introduction and notations

We use the notations in [4]. Let X be a compact metric space. We denote by 2^X the collection of all subsets of X , and denote by $\mathcal{C}(X)$ the collection of all closed subsets of X . Let ρ be the Hausdorff metric on 2^X . Then it is well-known ([3]) that $(\mathcal{C}(X), \rho)$ is a compact metric space. Let $\mathcal{F}(X)$ be the set of all fuzzy sets $\tilde{s} : X \rightarrow [0, 1]$ which are upper semi-continuous and satisfy $\sup_{x \in X} \tilde{s}(x) = 1$. Let $\tilde{q} : X \times X \rightarrow [0, 1]$ be a continuous fuzzy relation on X .

In this paper, we consider the existence and uniqueness of solution $\tilde{p} \in \mathcal{F}(X)$ in the following fuzzy relational equation (1.1) for given a continuous fuzzy relation \tilde{q} on X (see [4]) :

$$\tilde{p}(y) = \sup_{x \in X} \{\tilde{p}(x) \wedge \tilde{q}(x, y)\}, \quad y \in X, \quad (1.1)$$

where $a \wedge b = \min\{a, b\}$ for real numbers a and b . We define a map $\tilde{q}_\alpha : 2^X \rightarrow 2^X$ ($\alpha \in [0, 1]$) by

$$\tilde{q}_\alpha(D) := \begin{cases} \{y \mid \tilde{q}(x, y) \geq \alpha \text{ for some } x \in D\} & \text{for } \alpha \neq 0, D \in 2^X, D \neq \emptyset, \\ \text{cl}\{y \mid \tilde{q}(x, y) > 0 \text{ for some } x \in D\} & \text{for } \alpha = 0, D \in 2^X, D \neq \emptyset, \\ X & \text{for } 0 \leq \alpha \leq 1, D = \emptyset, \end{cases} \quad (1.2)$$

where cl denotes the closure of a set. Especially, we put $\tilde{q}_\alpha(x) := \tilde{q}_\alpha(\{x\})$ for $x \in X$. We note that $\tilde{q}_\alpha : \mathcal{C}(X) \rightarrow \mathcal{C}(X)$.

Lemma 1.1 ([4, Lemma 2]). *For each $\alpha \in [0, 1]$, the map $\tilde{q}_\alpha : \mathcal{C}(X) \rightarrow \mathcal{C}(X)$ is continuous with respect to ρ .*

For $\tilde{s} \in \mathcal{F}(X)$, the α -cut \tilde{s}_α , $\alpha \in [0, 1]$ is defined by

$$\tilde{s}_\alpha := \{x \in X \mid \tilde{s}(x) \geq \alpha\} \quad (\alpha \neq 0) \quad \text{and} \quad \tilde{s}_0 := \text{cl}\{x \in X \mid \tilde{s}(x) > 0\}.$$

Lemma 1.2.

(i) *For $\tilde{s} \in \mathcal{F}(X)$, \tilde{s} satisfies (1.1) if and only if*

$$\tilde{q}_\alpha(\tilde{s}_\alpha) = \tilde{s}_\alpha, \quad \alpha \in [0, 1]. \quad (1.3)$$

(ii) *We suppose that a family of subsets $\{D_\alpha \mid \alpha \in [0, 1]\} (\subset \mathcal{C}(X))$ satisfies the following conditions (a), (b) and (c):*

- (a) $D_\alpha \subset D_{\alpha'}$ for $0 \leq \alpha' < \alpha \leq 1$;
- (b) $\lim_{\alpha' \uparrow \alpha} D_{\alpha'} = D_\alpha$ for $\alpha \neq 0$;
- (c) $\tilde{q}_\alpha(\tilde{s}_\alpha) = \tilde{s}_\alpha$ for $\alpha \in [0, 1]$.

Then $\tilde{s}(x) := \sup_{\alpha \in [0, 1]} \{\alpha \wedge 1_{D_\alpha}(x)\}$, $x \in X$, satisfies $\tilde{s} \in \mathcal{F}(X)$ and (1.1), where 1_D denotes the characteristic function of a set $D \in 2^X$.

Proof. (i) is trivial. (ii) is from (i) and [4, Lemma 3]. \square

2 The existence of solutions

For $\alpha \in [0, 1]$ and $x \in X$, a sequence $\{\tilde{q}_\alpha^k(x)\}_{k=1,2,\dots}$ is defined iteratively by

$$\tilde{q}_\alpha^0(x) := \{x\}, \quad \tilde{q}_\alpha^1(x) := \tilde{q}_\alpha(x) \quad \text{and} \quad \tilde{q}_\alpha^{k+1}(x) := \tilde{q}_\alpha(\tilde{q}_\alpha^k(x)) \quad \text{for } k = 1, 2, \dots$$

Then, let $G_\alpha(x) := \bigcup_{k=1}^{\infty} \tilde{q}_\alpha^k(x)$ and

$$F_\alpha(x) := \bigcup_{k=0}^{\infty} \tilde{q}_\alpha^k(x) = \{x\} \cup G_\alpha(x). \quad (2.1)$$

We now consider a class of invariant points for this iteration procedure, that is, $x \in G_\alpha(x)$. So put

$$R_\alpha := \{x \in X \mid x \in G_\alpha(x)\} \quad \text{for } \alpha \in [0, 1]. \quad (2.2)$$

Each state of R_α is called as an “ α -recurrent” state and it is studied by [7]. The following properties (i) and (ii) holds clearly:

- (i) $\tilde{q}_\alpha(F_\alpha(x)) = G_\alpha(x)$ for $\alpha \neq 0$ and $x \in X$;
- (ii) $R_\alpha \subset R_{\alpha'}$ for $0 \leq \alpha' < \alpha \leq 1$.

Lemma 2.1. *If $z \in R_1$, the following (i) and (ii) hold:*

- (i)
$$\tilde{q}_\alpha(F_\alpha(z)) = F_\alpha(z) \quad \text{for } \alpha \in [0, 1]; \quad (2.3)$$
- (ii) $F_\alpha(z) \subset F_{\alpha'}(z)$ for $0 \leq \alpha' < \alpha \leq 1$.

Proof. Since $z \in R_1 \subset R_\alpha$, we have

$$\tilde{q}_\alpha(F_\alpha(z)) = G_\alpha(z) = F_\alpha(z).$$

So, we obtain (i). (ii) is trivial. \square

For $z \in R_1$, we define

$$\hat{F}_\alpha(z) := \bigcap_{\alpha' < \alpha} \text{cl}\{F_{\alpha'}(z)\} \quad (\alpha \neq 0) \quad \text{and} \quad \hat{F}_0(z) := \text{cl}\{F_0(z)\}, \quad (2.4)$$

where $\text{cl}\{F_\alpha(z)\}$ denotes the closure of $F_\alpha(z)$.

Lemma 2.2. *If $z \in R_1$, the following (i), (ii) and (iii) hold:*

- (i) $\tilde{q}_\alpha(\hat{F}_\alpha(z)) = \hat{F}_\alpha(z)$ for $\alpha \in [0, 1]$;
- (ii) $\hat{F}_\alpha(z) \subset \hat{F}_{\alpha'}(z)$ for $0 \leq \alpha' < \alpha \leq 1$;
- (iii) $\hat{F}_\alpha(z) = \lim_{\alpha' \uparrow \alpha} \hat{F}_{\alpha'}(z)$ for $\alpha \neq 0$.

Proof. (ii) is trivial from Lemma 2.1 and (iii) is also trivial from the definition. To prove (i), let $\alpha = 0$. From Lemma 2.1(i), we have $\tilde{q}_\alpha(F_0(z)) = F_0(z)$. By the continuity of \tilde{q} , we can check $\tilde{q}_\alpha(\text{cl}\{F_0(z)\}) = \text{cl}\{F_0(z)\}$ in similar way to the proof of [4, Lemma 1]. Therefore, $\tilde{q}_0(\hat{F}_\alpha(z)) = \hat{F}_0(z)$.

Let $\alpha > 0$ and $y \in \tilde{q}_\alpha(\hat{F}_\alpha(z))$. By Lemma 1.1, we have

$$y \in \bigcap_{\alpha' < \alpha} \tilde{q}_{\alpha'}(\text{cl}\{F_{\alpha'}(z)\}) = \bigcap_{n=1}^{\infty} \tilde{q}_{\alpha'}(\text{cl}\{F_{(\alpha-1/n)\vee 0}(z)\}).$$

From the continuity of \tilde{q} , for $n \geq 1$, there exists $x_n \in F_{(\alpha-1/n)\vee 0}(z)$ such that $\tilde{q}(x_n, y) \geq \alpha - 1/n$. By Lemma 2.1(i),

$$y \in \tilde{q}_{(\alpha-1/n)\vee 0}(F_{(\alpha-1/n)\vee 0}(z)) = F_{(\alpha-1/n)\vee 0}(z) \subset \text{cl}\{F_{(\alpha-1/n)\vee 0}(z)\} \quad \text{for all } n \geq 1.$$

So, $y \in \hat{F}_\alpha(z)$. Therefore, we obtain

$$\tilde{q}_\alpha(\hat{F}_\alpha(z)) \subset \hat{F}_\alpha(z).$$

While, we have

$$\text{cl}\{F_{\alpha'}(z)\} \subset \tilde{q}_{\alpha'}(\text{cl}\{F_{\alpha''}(z)\}) \quad \text{for } \alpha'' < \alpha' < \alpha.$$

Then

$$\hat{F}_\alpha(z) = \bigcap_{\alpha' < \alpha} \text{cl}\{F_{\alpha'}(z)\} \subset \bigcap_{\alpha' < \alpha} \tilde{q}_{\alpha'}(\text{cl}\{F_{\alpha''}(z)\}) = \tilde{q}_\alpha(\text{cl}\{F_{\alpha''}(z)\}) \quad \text{for } \alpha'' < \alpha.$$

So, we get

$$\hat{F}_\alpha(z) \subset \bigcap_{\alpha'' < \alpha} \tilde{q}_{\alpha'}(\text{cl}\{F_{\alpha''}(z)\}) = \tilde{q}_\alpha \left(\bigcap_{\alpha'' < \alpha} \text{cl}\{F_{\alpha''}(z)\} \right) = \tilde{q}_\alpha(\hat{F}_\alpha(z)).$$

Therefore, we can obtain (i). \square

Let $z \in R_1$. Since $\{\hat{F}_\alpha(z) \mid \alpha \in [0, 1]\}$ satisfies the conditions (a) – (c) of Lemma 2.1(ii), we obtain the following theorem.

Theorem 2.1.

- (i) If $R_1 \neq \emptyset$, then there exists a solution of (1.1).
- (ii) Define a fuzzy state

$$\tilde{s}^z(x) := \sup_{\alpha \in [0,1]} \left\{ \alpha \wedge 1_{\hat{F}_\alpha(z)}(x) \right\}, \quad x \in X. \tag{2.5}$$

Then $\tilde{s}^z \in \mathcal{F}(X)$ satisfies (1.1).

Assume that $R_1 \neq \emptyset$. We introduce an equivalent relation \sim on R_α as follows: For $z_1, z_2 \in R_\alpha$,

$$z_1 \sim z_2 \quad \text{means that} \quad z_1 \in F_\alpha(z_2) \text{ and } z_2 \in F_\alpha(z_1).$$

Then we could identify the states of R_α which is equivalent with respect to \sim , and so put

$$R_\alpha^\sim := R_\alpha / \sim.$$

Lemma 2.3. For $z_1, z_2 \in R_1$,

$$z_1 \sim z_2 \quad \text{if and only if} \quad F_\alpha(z_1) = F_\alpha(z_2) \text{ for all } \alpha \in [0, 1].$$

Proof. Let $z_1 \sim z_2$. Then, we have $z_1 \in F_1(z_2) \subset F_\alpha(z_2)$ for any $\alpha \in [0, 1]$. From the definition (2.1) of $F_\alpha(z_1)$, we obtain $F_\alpha(z_1) \subset F_\alpha(z_2)$. Since we have $F_\alpha(z_2) \subset F_\alpha(z_1)$ similarly, $F_\alpha(z_1) = F_\alpha(z_2)$ holds. The reverse proof is trivial. \square

From Theorem 2.1 and Lemma 2.3, the number of solutions of (1.1) is greater than or equals to the number of “1-recurrent” sets. To consider the class of solution (1.1), let $P := \{\tilde{p} \in \mathcal{F}(X) \mid \tilde{p} \text{ is a solution of (1.1)}\}$. Then P has the following property:

Theorem 2.2. Let $\tilde{p}^k \in P$ ($k = 1, 2, \dots, l$). Then:

(i) Put

$$\tilde{p}(x) := \max_{k=1,2,\dots,l} \tilde{p}^k(x), \quad x \in X.$$

Then $\tilde{p} \in P$.

(ii) Let $\{\alpha^k \in [0, 1] \mid k = 1, 2, \dots, l\}$ satisfy $\max_{k=1,2,\dots,l} \alpha^k = 1$. Put

$$\tilde{p}(x) := \max_{k=1,2,\dots,l} \{\alpha^k \wedge \tilde{p}^k(x)\} \quad \text{for } x \in X.$$

Then $\tilde{p} \in P$.

Proof. (ii) Taking the α -cut of $\tilde{p} \in \mathcal{F}(X)$, we have

$$\tilde{p}_\alpha := \bigcup_{k:\alpha^k \geq \alpha} \tilde{p}_\alpha^k.$$

Then,

$$\tilde{q}_\alpha(\tilde{p}_\alpha) = \tilde{q}_\alpha \left(\bigcup_{k:\alpha^k \geq \alpha} \tilde{p}_\alpha^k \right) = \bigcup_{k:\alpha^k \geq \alpha} \tilde{q}_\alpha(\tilde{p}_\alpha^k) = \bigcup_{k:\alpha^k \geq \alpha} \tilde{p}_\alpha^k = \tilde{p}_\alpha.$$

Therefore, we obtain (ii) from Lemma 1.2(i). (i) is proved similarly. \square

3 The uniqueness of solutions

In this section, we discuss the uniqueness of a solution of the equation (1.1) under convexity and compactness. Let B be a convex subset of \mathcal{R}^n and $C_c(B)$ the class of all closed and convex subsets of B . Throughout this section, we assume that the state space X is a convex and compact subset of \mathcal{R}^n . The fuzzy set $\tilde{s} \in \mathcal{F}(X)$ is called convex if its α -cut \tilde{s}_α is convex for each $\alpha \in [0, 1]$. Let $\mathcal{F}_c(X) := \{\tilde{s} \in \mathcal{F} \mid \tilde{s} \text{ is convex}\}$.

Let, applying Kakutani's fixed point theorem ([2]), we have the following.

Lemma 3.1. *Let $\alpha \in [0, 1]$ and $\tilde{q}(x)$ is convex for each $x \in X$. Then, for any $A \in C_c(X)$ with $A = \tilde{q}_\alpha(A)$, there exists an $x \in X$ such that $\tilde{q}(x, x) \geq \alpha$.*

Proof. The map $\tilde{q}_\alpha : A \rightarrow C_c(A)$ with $\tilde{q}_\alpha(x) \in C_c(A)$ for all $x \in A$ is continuous from Lemma 1.1, so Kakutani's fixed point theorem guarantees the existence of an element $x \in A$ such that $x \in \tilde{q}_\alpha(x)$, which implies $\tilde{q}(x, x) \geq \alpha$. This completes the proof.

As a consequence, we have a property of the convex solution of (1.1).

Proposition 3.1. *Let $p \in \mathcal{F}_c(X)$ be a solution of (1.1). Then, for each $\alpha \in [0, 1]$, there exists an $x \in p_\alpha$ with $q(x, x) \geq \alpha$.*

Proof. By Lemma 1.2, $\tilde{p}_\alpha = \tilde{q}_\alpha(\tilde{p}_\alpha)$ for each $\alpha \in [0, 1]$. Thus, Lemma 3.1 clearly proves the desired result. \square

Now, we give sufficient conditions for the uniqueness of a convex solution of (1.1). Let $U_\alpha := \{x \in X \mid q(x, x) \geq \alpha\}$.

Assumption A. The following A1 – A3 holds.

- A1. The set $U_\alpha, \alpha = 1$ is one point set, say u . That is, $U_1 = \{u\}$.
- A2. $U_\alpha \subset F_\alpha(u)$ for each $\alpha \in [0, 1]$, where u is in the above A1 and $F_\alpha(u)$ is defined by (2.1).
- A3. If $A = \tilde{q}_\alpha(A)$ for any $A \in F_c(X), 0 \leq \alpha \leq 1$, then

$$A = \bigcup_{x \in U_\alpha \cap A} F_\alpha(A)$$

Theorem 3.1. Under Assumption A, the equation (1.1) has a unique solution in $\mathcal{F}_c(X)$.

Proof. Let $\tilde{p}, \tilde{p}' \in \mathcal{F}_c(X)$ be solution of (1.1). By Lemma 3.1, $\tilde{p}_1 \cap U_1 \neq \emptyset$ and $\tilde{p}'_1 \cap U_1 \neq \emptyset$. Since U_1 is one pointset, $u \in \tilde{p}_1$ and $u \in \tilde{p}'_1$. Thus, by A2 and A3, $\tilde{p}_1 = F_1(u)$ and $\tilde{p}'_1 = F_1(u)$, which implies $\tilde{p}_1 = \tilde{p}'_1$. We now show that $\tilde{p}_\alpha = \hat{F}_\alpha(u)$ for each $0 < \alpha \leq 1$. Since $u \in \tilde{p}_{\alpha'} = \tilde{q}_{\alpha'}(\tilde{p}_{\alpha'})$ and \tilde{p}_α is closed, it holds $F_{\alpha'}(u) \subset \tilde{p}_{\alpha'}$. Therefore,

$$\hat{F}_\alpha(u) = \bigcup_{\alpha' < \alpha} \text{cl}\{F_{\alpha'}(u)\} \subset \bigcup_{\alpha' < \alpha} \tilde{p}_{\alpha'}(u) = \tilde{p}_\alpha.$$

On the other hand, we have

$$\begin{aligned} \tilde{p}_\alpha &= \bigcup_{x \in U_\alpha \cap \tilde{p}_\alpha} \text{cl}\{F_\alpha(x)\}, & \text{from A3} \\ &\subset \bigcup_{x \in F_\alpha \cap \tilde{p}_\alpha} \text{cl}\{F_\alpha(x)\}, & \text{from A2} \\ &\subset \bigcup_{x \in \hat{F}_\alpha} \text{cl}\{F_\alpha(x)\}. \end{aligned}$$

From that $x \in \hat{F}_\alpha$ means $\hat{F}_\alpha(x) \subset \hat{F}_\alpha(u)$, it holds that

$$\tilde{p}_\alpha \subset \text{cl}\{F_\alpha(u)\} \subset \hat{F}_\alpha(u).$$

The above shows $\tilde{p}_\alpha = \hat{F}_\alpha(u)$. Similarly $\tilde{p}'_\alpha = \hat{F}_\alpha(u)$. Thus, $\tilde{p}_\alpha = \tilde{p}'_\alpha$. This completes the proof. \square

4 Numerical example

Here two numerical examples are given to comprehend computational aspect of this paper.

Example 1. Let $X = [0, 1]$. For any $g; [0, 1] \rightarrow [0, 1]$, let

$$\tilde{q}(x, y) := (1 - |y - g(x)|) \vee 0.$$

We assume that $g(\cdot)$ is strictly increasing and there exists a unique $x_0 \in [0, 1]$ with $x_0 = g(x_0)$. Under the above condition, $R_1 = \{x_0\}$ and for each $\alpha \in [0, 1]$,

$$U_\alpha = [\underline{x}_\alpha, \bar{x}_\alpha],$$

when $\underline{x}_\alpha, \bar{x}_\alpha$ is a unique solution of $x = g(x) - (1 - \alpha)$, $x = g(x) + (1 - \alpha)$ respectively and $\underline{x}_\alpha = 0, \bar{x}_\alpha = 1$ if the solution does not exist in $[0, 1]$.

Clearly, U_α equals a unique solution of the equation $A = \tilde{p}_\alpha(A)$ in $C_c([0, 1])$, so that Assumption A in Section 3 holds in this case. Thus, by Theorem 3.1, we have

$$\tilde{s}(x) = \sup_{\alpha \in [0, 1]} \{\alpha \wedge I_{U_\alpha}(x)\} \quad (4.1)$$

is a unique convex solution of (1.1). For a concrete example such as $g(x) = (2x^2 + 1)/4$, then it is seen that $R_1 = \{(2 - \sqrt{2})/2\}$ and


$$\underline{x}_\alpha = \begin{cases} 1 - \sqrt{5/2 - 2\alpha} \\ 1, \end{cases} \vee 0, \quad \begin{matrix} 3/4 < \alpha \\ 3/4 \leq \alpha \leq 1. \end{matrix}$$

$$\bar{x}_\alpha = \begin{cases} 1 - \sqrt{2\alpha - 3/2}, \\ 1, \end{cases}$$

By (4.1), the unique solution is as follows (Fig.1):

$$\tilde{s}(x) = \begin{cases} -x^2/2 + x + 3/4, & 0 \leq x \leq 1 - \sqrt{2}/2 \\ x^2/2 - x + 5/4, & 1 - \sqrt{2}/2 < x \leq 1 \end{cases}$$

Fig.1 The unique solution \tilde{s} .



Example 2. This example has two peaks for the fuzzy relation. Let $X = [0, 1]$ and

$$\tilde{q}(x, y) = (1 - |y - (x^2 + 1)/4|) \vee (1 - |y - (x^2 + 2)/4|).$$

Then, $R_1 = \{a, b\}$, where $a = 2 - \sqrt{3}$, $b = 2 - \sqrt{2}$. By simple calculation, we get

$$\hat{F}_\alpha(a) = [\underline{x}_\alpha^a, \bar{x}_\alpha^a] \quad \text{and} \quad \hat{F}_\alpha(b) = [\underline{x}_\alpha^b, \bar{x}_\alpha^b]$$

for $\alpha \in [0, 1]$, where

$$\underline{x}_\alpha^a = \begin{cases} 0, & 0 \leq \alpha \leq 3/4 \\ 2 - \sqrt{7 - 4\alpha}, & 3/4 < \alpha \leq 1 \end{cases}$$

$$\bar{x}_\alpha^a = \begin{cases} 1, & 0 \leq \alpha \leq 7/8 \\ 2 - \sqrt{4\alpha - 1}, & 7/8 < \alpha \leq 1 \end{cases}$$

$$\underline{x}_\alpha^b = \begin{cases} 0, & 0 \leq \alpha \leq 7/8 \\ 2 - \sqrt{6 - 4\alpha}, & 7/8 < \alpha \leq 1 \end{cases}$$


$$\bar{x}_\alpha^b = \begin{cases} 1, & 0 \leq \alpha \leq 3/4 \\ 2 - \sqrt{4\alpha - 2}, & 3/4 < \alpha \leq 1 \end{cases}$$

By Theorem 2.1, the solution of (1.1) are given as follows(Fig2.):

$$\tilde{s}^a(x) = \begin{cases} \frac{-x^2 + 2x + 3}{4}, & 0 \leq x \leq 2 - \sqrt{3} \\ \frac{x^2 - 4x + 5}{4}, & 2 - \sqrt{3} \leq x \leq 2 - \sqrt{10}/2 \\ 7/8, & 2 - \sqrt{10}/2 \leq x \leq 1 \end{cases}$$

$$\tilde{s}^b(x) = \begin{cases} 7/8, & 0 \leq x \leq 2 - \sqrt{10}/2 \\ \frac{-x^2 + 4x + 2}{4}, & 2 - \sqrt{10}/2 \leq x \leq 2 - \sqrt{2} \\ \frac{x^2 - 4x + 6}{4}, & 2 - \sqrt{2} \leq x \leq 1 \end{cases}$$

Fig.2 The unique solutions \tilde{s}^a and \tilde{s}^b .



References

- [1] R.E.Bellman and L.A.Zadeh, Decision-making in a fuzzy environment, *Management Sci. Ser B.* **17** (1970) 141-164.
- [2] N.Dunford and J.T.Schwartz, *Linear Operators, Part 1: General Theory* (Interscience Publishers, New York, 1958).
- [3] K.Kuratowski, *Topology I* (Academic Press, New York, 1966).
- [4] M.Kurano, M.Yasuda, J.Nakagami and Y.Yoshida, A limit theorem in some dynamic fuzzy systems, *Fuzzy Sets and Systems* **51** (1992) 83-88.
- [5] V.Novák, *Fuzzy Sets and Their Applications* (Adam Hilder, Bristol-Boston, 1989).
- [6] Y.Yoshida, M.Yasuda, J.Nakagami and M.Kurano, A limit theorem in some dynamic fuzzy systems with a monotone property, *accepted in Fuzzy Sets and Systems*.
- [7] Y.Yoshida, The recurrence of dynamic fuzzy systems, *RIFIS Technical Report 80* (1994), Research Institute of Fundamental Information Science, Kyushu University 33 Fukuoka 812, Japan.