TITLE:

Polynomiality of Primal-Dual Algorithms for Semidefinite Linear Complementarity Problems Based on the Kojima-Shindoh-Hara Family of Directions

AUTHOR(S):

Monteiro, Ronato D.C.; Tsuchiya, Takashi

CITATION:

Monteiro, Ronato D.C. ...[et al]. Polynomiality of Primal-Dual Algorithms for Semidefinite Linear Complementarity Problems Based on the Kojima-Shindoh-Hara Family of Directions. 数理解析研究所講究録 1997, 1004: 138-152

ISSUE DATE:

1997-06

URL:

<http://hdl.handle.net/2433/61438>

RIGHT:

Polynomiality of Primal-Dual Algorithms for Semidefinite Linear Complementarity Problems Based on the Kojima-Shindoh-Hara Family of Directions

Renato D.C. Monteiro*

Takashi Tsuchiya†

August 6, 1996

Abstract

Kojima, Shindoh and Hara proposed a family of search directions for the semidefinite linear complementarity problem (SDLCP) and established polynomial convergence of a feasible short-step path-following algorithm based on a particular direction of their family. The question of whether polynomiality could be established for any direction of their family thus remained an open problem. This paper answers this question in the affirmative by establishing the polynomiality of primal-dual interior-point algorithms for SDLCP based on any direction of the Kojima, Shindoh and Hara family of search directions. We show that the polynomial iteration-complexity bounds of two well-known algorithms for linear programming, namely the short-step path-following algorithm of Kojima et al. and Monteiro and Adler, and the predictor-corrector algorithm of Mizuno et al., carry over to the context of SDLCP.

keywords: Semidefinite programming, interior-point methods, polynomial complexity, path-following methods, primal-dual methods.

AMS 1991 subject classification: 65K05, 90C25, 90C30.

1 Introduction

Several authors have discussed generalizations of interior-point algorithms for linear programming (LP) to the context of semidefinite programming (SDP) and the more general semidefinite linear complementarity problem (SDLCP). The landmark work in this direction is due to Nesterov and Nemirovskii [19, 20] where a general approach for using interior-point methods for solving convex programs is proposed based on the notion of self-concordant functions. (See their book [22] for a comprehensive treatment of this subject.) They show that the problem of minimizing a linear function over a convex set can be solved in “polynomial time” as long as a self-concordant barrier function for the convex set is known. In particular, Nesterov and Nemirovskii show that linear

---

*School of Industrial and Systems Engineering, Georgia Institute of Technology, Atlanta, Georgia 30332, USA. (e-mail: monteiro@isye.gatech.edu). The work of this author was based on research supported by the Office of Naval Research under grant N00014-94-1-0340.

†The Institute of Statistical Mathematics, 4-6-7 Minami-Azabu, Minato-Ku, Tokyo, 106, Japan. (e-mail: tsuchiya@sun312.ism.ac.jp). This research is supported in part by the Grant-in-Aid for Encouragement of Young Scientists (A) 08780227 of the Ministry of Science, Education and Culture of Japan.

programs, convex quadratic programs with convex quadratic constraints, and semidefinite programs all have explicit and easily computable self-concordant barrier functions, and hence can be solved in “polynomial time”. On the other hand, Alizadeh [1] extends Ye’s projective potential reduction algorithm [30] for LP to SDP and argues that many known interior point algorithms for LP can also be transformed into algorithms for SDP in a mechanical way. Since then many authors have proposed interior-point algorithms for solving the SDP problem and SDLCP, including Alizadeh, Haeberly and Overton [2], Freund [3], Helmberg, Rendl, Vanderbei and Wolkowicz [4], Jarre [5], Kojima, Shida and Shindoh [8], Kojima, Shindoh and Hara [10], Lin and Saigal [11], Luo, Sturm and Zhang [12], Monteiro [14, 15], Monteiro and Zhang [18], Nesterov and Nemirovskii [21], Nesterov and Todd [24, 23], Potra and Sheng [25], Sturm and Zhang [26], Tseng [28], Vandenberghe and Boyd [29], and Zhang [31]. Most of these more recent works are concentrated on primal-dual methods.

The first algorithms for SDP and SDLCP that are extensions of primal-dual algorithms for LP, such as the long-step path-following algorithm of Kojima, Mizuno and Yoshise [7], the short-step path-following algorithm of Kojima, Mizuno and Yoshise [6] and Monteiro and Adler [16, 17], and the predictor-corrector algorithm of Mizuno, Todd and Ye [13], use one of the following three search directions: i) the Alizadeh, Haeberly and Overton (AHO) direction proposed in [2], ii) the KSH/HRVW/M direction independently proposed by Kojima, Shindoh and Hara [10] and Helmberg, Rendl, Vanderbei and Wolkowicz [4], and later rediscovered by Monteiro [14] via a formulation based on a scaling and symmetrization of the Newton equation, and iii) the Nesterov and Todd (NT) direction introduced in [23, 24].

To unify the above directions, two family of search directions have been proposed in the literature. The first family, proposed by Kojima, Shindoh and Hara [10], is known to contain the KSH/HRVW/M and NT directions but not the AHO direction. The second family, namely the Monteiro and Zhang (MZ) family, formally introduced by Zhang [31] to generalize a symmetric formulation of the KSH/HRVW/M direction proposed by Monteiro [14], contains all three search directions above. Proofs that the NT direction is a member of both the KSH family and the MZ family can be found in Kojima, Shida and Shindoh [9] and Todd, Toh and Tütüncü [27], respectively.

Unified convergence analyses for the MZ family have been given by Monteiro and Zhang [18] and Monteiro [15]. In the paper [18], iteration-complexity bounds are derived for long-step primal-dual path-following methods based on a subclass of the MZ family of search directions, which contains the KSH/HRVW/M and NT directions but not the AHO direction. In particular, it is shown that the corresponding algorithms based on the KSH/HRVW/M and NT directions perform $\mathcal{O}(n^{3/2}L)$ and $\mathcal{O}(nL)$ iterations, respectively, to reduce the duality gap by a factor of at least $2^{-\mathcal{O}(L)}$. (The $\mathcal{O}(n^{3/2}L)$ iteration-complexity bound for the KSH/HRVW/M direction was in fact obtained earlier by Monteiro [14].) More recently, Monteiro [15] proves the polynomiality of short-step path following algorithms and Mizuno-Todd-Ye predictor-corrector type algorithms based on any member of the MZ family, thus obtaining as a by-product the important result that Frobenius-norm type algorithms based on the AHO direction are polynomial.

Unified analysis for the KSH family of search directions are provided in Kojima, Shindoh and Hara [10]. This paper deals with primal-dual path-following algorithms for the semidefinite linear complementarity problem based on the KSH family of search directions and establishes the polynomiality of: 1) a feasible short-step path-following method based on a *special* member of their family, namely the KSH/HRVW/M direction and; 2) a (feasible and infeasible) potential reduction algorithm based on *any* search direction of their family. The question of whether polynomiality

of algorithm 1) can be established for any direction of the KSH family was thus left as an open problem.

In this paper, we answer the above question in the affirmative. Using new techniques recently proposed by Monteiro [15], we prove the polynomial convergence of two feasible primal-dual algorithms based on a narrow (or Frobenius norm) neighborhood of the central path, namely: a short-step path-following method which is an extension of the LP method of Kojima, Mizuno and Yoshise [6] and Monteiro and Adler [16, 17], and a predictor-corrector algorithm similar to the LP one of Mizuno, Todd and Ye [13].

This paper is organized as follows. In Section 2, we introduce the SDLCP problem and motivate the search directions used by the algorithms studied in this paper. In Section 3, we state and prove the technical results used in the polynomial convergence analysis of the algorithms of Section 4. In Section 4, we establish the polynomiality of two primal-dual feasible algorithms: the short-step path-following algorithm in Subsection 4.1 and the predictor-corrector algorithm in Subsection 4.2. We give some concluding remarks in Section 5.

1.1 Notation and terminology

The following notation is used throughout the paper. The superscript T denotes transpose. $\mathbb{R}^p$ denotes the p -dimensional Euclidean space. The set of all $p \times q$ matrices with real entries is denoted by $\mathbb{R}^{p \times q}$. The set of all symmetric $p \times p$ matrices is denoted by $\mathcal{S}^p$. For $Q \in \mathcal{S}^p$, $Q \succeq 0$ means Q is positive semidefinite and $Q \succ 0$ means Q is positive definite. The trace of a matrix $Q \in \mathbb{R}^{p \times p}$ is denoted by $\text{Tr } Q \equiv \sum_{i=1}^p Q_{ii}$. For a matrix $Q \in \mathbb{R}^{p \times p}$ with all real eigenvalues, we denote its eigenvalues by $\lambda_i[Q]$, $i = 1, \dots, p$, and its smallest eigenvalue by $\lambda_{\min}[Q]$. Given P and Q in $\mathbb{R}^{p \times q}$, the inner product between them in the vector space $\mathbb{R}^{p \times q}$ is defined as $P \bullet Q \equiv \text{Tr } P^T Q$. The Euclidean norm and its associated operator norm are both denoted by $\|\cdot\|$; hence, $\|Q\| \equiv \max_{\|u\|=1} \|Qu\|$ for any $Q \in \mathbb{R}^{p \times p}$. The Frobenius norm of $Q \in \mathbb{R}^{p \times p}$ is $\|Q\|_F \equiv (Q \bullet Q)^{1/2}$. $\mathcal{S}_+^p$ and $\mathcal{S}_{++}^p$ denote the set of all matrices in $\mathcal{S}^p$ which are positive semidefinite and positive definite, respectively. $\mathcal{S}_\perp^p$ denote the set of all skew-symmetric matrices in $\mathbb{R}^{p \times p}$. Since $\mathcal{S}^p + \mathcal{S}_\perp^p = \mathbb{R}^{p \times p}$ and $U \bullet V = 0$ for every $U \in \mathcal{S}^p$ and $V \in \mathcal{S}_\perp^p$, it follows that $\mathcal{S}_\perp^p$ is the orthogonal complement of $\mathcal{S}^p$ with respect to the inner product $\bullet$.

2 Description of the problem and preliminary discussion

In this section, we introduce the semidefinite linear complementarity problem and the assumptions made in our presentation. We also describe the family of search directions introduced by Kojima, Shindoh and Hara [10] and give a short proof for the existence and uniqueness of these directions.

Let $\mathcal{L}$ be an affine subspace of $\mathcal{S}^n \times \mathcal{S}^n$ whose dimension is $n(n+1)/2$. Let

$$\begin{aligned}\mathcal{L}_+ &\equiv \mathcal{L} \cap (\mathcal{S}_+^n \times \mathcal{S}_+^n), \\ \mathcal{L}_{++} &\equiv \mathcal{L} \cap (\mathcal{S}_{++}^n \times \mathcal{S}_{++}^n).\end{aligned}$$

In this paper, we deal with the semidefinite linear complementarity problem (SDLCP) of finding a pair (X, S) such that

$$(X, S) \in \mathcal{L}_+, \quad X \bullet S = 0. \quad (1)$$

Throughout our presentation, we assume that

[A1] $\mathcal{L}$ is monotone, that is $(X_1 - X_2) \bullet (S_1 - S_2) \geq 0$ for any $(X_1, S_1) \in \mathcal{L}$ and $(X_2, S_2) \in \mathcal{L}$.

[A2] $\mathcal{L}_{++}$ is nonempty.

This problem is a generalization of SDP which has numerous applications in systems and control theory and combinatorial optimization. Given $C \in \mathcal{S}^n$ and $(A_i, b_i) \in \mathcal{S}^n \times \mathbb{R}$ for $i = 1, \dots, m$, a primal-dual pair of SDP problems is defined as

$$(P) \quad \min\{C \bullet X : A_i \bullet X = b_i, i = 1, \dots, m, X \succeq 0\},$$

$$(D) \quad \max\{b^T y : \sum_{i=1}^m y_i A_i + S = C, S \succeq 0\},$$

where $b \equiv (b_1, \dots, b_m)^T$. Under the assumption that problems (P) and (D) have interior feasible solutions, that is feasible solutions X and (S, y) satisfying $X \succ 0$ and $S \succ 0$, it is known that (X, S) is a solution of (1) with

$$\mathcal{L} = \{(X, S) \in \mathcal{S}^n \times \mathcal{S}^n : A_i \bullet X = b_i, \sum_{i=1}^m y_i A_i + S = C \text{ for some } y \in \mathbb{R}^m\},$$

if and only if (X, S, y) is a solution of (P) and (D) for some $y \in \mathbb{R}^m$. In this case, it is easy to see that $\mathcal{L}$ is a monotone affine space satisfying $(X_1 - X_2) \bullet (S_1 - S_2) = 0$ for any $(X_1, S_1) \in \mathcal{L}$ and $(X_2, S_2) \in \mathcal{L}$.

Under assumptions [A1] and [A2], it is known that problem (1) has at least one solution. Since for $(X, S) \in \mathcal{S}_+^n \times \mathcal{S}_+^n$, we have $X \bullet S = 0$ if and only if $XS = 0$, problem (1) is equivalent to find a pair (X, S) such that

$$(X, S) \in \mathcal{L}_+, \quad XS = 0.$$

It has been shown by Kojima, Shindoh and Hara [10] that the perturbed system

$$(X, S) \in \mathcal{L}_+, \quad XS = \mu, \tag{2}$$

has a unique solution in $\mathcal{L}_{++}$, denoted by (X_μ, S_μ) , for every $\mu > 0$, and that $\lim_{\mu \rightarrow 0} (X_\mu, S_\mu)$ exists and is a solution of (1). The set $\{(X_\mu, S_\mu) : \mu > 0\}$ is called the central path associated with (1) and plays a fundamental role in the development of interior point algorithms for solving SDP and SDLCP. Another equivalent formulation of (2) is

$$(X, S) \in \mathcal{L}_+, \quad X^{1/2} S X^{1/2} = \mu I \quad (\text{or, } S^{1/2} X S^{1/2} = \mu I),$$

which motivates the following measure of closeness of $(X, S) \in \mathcal{S}_+^n \times \mathcal{S}_+^n$ to the point (X_μ, S_μ) of the central trajectory:

$$d_\mu(X, S) \equiv \|X^{1/2} S X^{1/2} - \mu I\|_F = \|S^{1/2} X S^{1/2} - \mu I\|_F,$$

and the following (feasible) neighborhood of (X_μ, S_μ) :

$$\mathcal{N}_F(\mu, \gamma) = \{(X, S) \in \mathcal{L}_+ : d_\mu(X, S) \leq \gamma\mu\},$$

where $\gamma > 0$ is a given constant. Both algorithms described in Section 4 generate their iterates in the neighborhood of the central path defined by

$$\mathcal{N}_F(\gamma) \equiv \cup_{\mu > 0} \mathcal{N}_F(\mu, \gamma).$$

Path-following algorithms for solving (1) are based on the idea of approximately tracing the central path. Application of Newton method for computing the solution of (2) with $\mu = \hat{\mu}$ leads to the Newton search direction $(\widehat{\Delta X}, \widehat{\Delta S})$ which solves the linear system

$$X\widehat{\Delta S} + \widehat{\Delta X}S = \hat{\mu}I - XS, \quad (X + \widehat{\Delta X}, S + \widehat{\Delta S}) \in \mathcal{L}. \quad (3)$$

Unfortunately, this system does not always have a solution. To overcome this difficulty, Kojima and Shindoh and Hara proposed the following modified Newton system of equations:

$$X(\Delta S + \widetilde{\Delta S}) + (\Delta X + \widetilde{\Delta X})S = \hat{\mu}I - XS, \quad (4a)$$

$$(X + \Delta X, S + \Delta S) \in \mathcal{L}, \quad (\widetilde{\Delta X}, \widetilde{\Delta S}) \in \mathcal{L}_\perp, \quad (4b)$$

where $\mathcal{L}_\perp$ is a linear subspace of $\mathfrak{R}^{n \times n} \times \mathfrak{R}^{n \times n}$ satisfying the following condition:

[A3] $\mathcal{L}_\perp \subseteq \mathcal{S}_\perp^n \times \mathcal{S}_\perp^n$, $\dim(\mathcal{L}_\perp) = n(n-1)/2$ and $\mathcal{L}_\perp$ is monotone, that is $U \bullet V \geq 0$ for every $(U, V) \in \mathcal{L}_\perp$.

It was shown in Corollary 4.3 of [10] that system (4) always has a unique solution. The symmetric component $(\Delta X, \Delta S)$ of this solution is then used as a search direction to generate the next point. In what follows we give another short proof of the existence and uniqueness of $(\Delta X, \widetilde{\Delta X}, \Delta S, \widetilde{\Delta S})$, which gives some intuition for the need to introduce the subspace $\mathcal{L}_\perp$.

Lemma 2.1 *Let $(X, S) \in \mathcal{S}_{++}^n \times \mathcal{S}_{++}^n$ and $\mathcal{W}$ be an n^2 dimensional affine subspace of $\mathfrak{R}^{n \times n} \times \mathfrak{R}^{n \times n}$ which is monotone, that is $(U_1 - U_2) \bullet (V_1 - V_2) \geq 0$ for every $(U_1, V_1), (U_2, V_2) \in \mathcal{W}$. Then, the system*

$$XV + US = H, \quad (U, V) \in \mathcal{W}, \quad (5)$$

has a unique solution for any $H \in \mathfrak{R}^{n \times n}$.

Proof. Consider the map $\Phi : \mathcal{W} \rightarrow \mathfrak{R}^{n \times n}$ defined by $\Phi(U, V) = XV + US$ for every $(U, V) \in \mathcal{W}$. Φ is an affine map between spaces of the same dimension since $\dim(\mathcal{W}) = n^2$ by assumption. Hence, it suffices to show that Φ is one-to-one. Indeed, assume that $\Phi(U_1, V_1) = \Phi(U_2, V_2)$ for some $(U_1, V_1), (U_2, V_2) \in \mathcal{W}$. Letting $\Delta U \equiv U_1 - U_2$ and $\Delta V \equiv V_1 - V_2$, and using the monotonicity of $\mathcal{W}$, we see that $\Delta U \bullet \Delta V \geq 0$ and $X\Delta V + \Delta U S = 0$. Multiplying the last relation on the left by $X^{-1/2}$ and on the right by $S^{-1/2}$, squaring both sides and using the fact that $\Delta U \bullet \Delta V \geq 0$, we obtain

$$0 = \|X^{1/2}\Delta U S^{-1/2} + X^{-1/2}\Delta V S^{1/2}\|_F^2 \geq \|X^{1/2}\Delta U S^{-1/2}\|_F^2 + \|X^{-1/2}\Delta V S^{1/2}\|_F^2.$$

Hence, $\Delta U = \Delta V = 0$, or equivalently $(U_1, V_1) = (U_2, V_2)$. ■

Lemma 2.1 provides the main reason for system (3) to not always have a solution, namely: the solution $(\widehat{\Delta X}, \widehat{\Delta S})$ is required to belong to the affine subspace $\mathcal{L} - (X, S)$, which only has dimension $n(n+1)/2 < n^2$. Adding the subspace $\mathcal{L}_\perp$ to $\mathcal{L}$ results in an affine subspace of dimension n^2 as required by Lemma 2.1. This fact is exploited in the proof of the following result which establishes the existence and uniqueness of the solution of (4).

Theorem 2.2 *System (4) has a unique solution.*

Proof. It is easy to see that $(\Delta X, \widetilde{\Delta X}, \Delta S, \widetilde{\Delta S})$ is a solution of (4) if and only if $(U, V) \equiv (\Delta X + \widetilde{\Delta X}, \Delta S + \widetilde{\Delta S})$ is a solution of (5) with $\mathcal{W} \equiv (\mathcal{L} - (X, Y)) + \mathcal{L}_\perp$ and $H \equiv \hat{\mu}I - XS$. Since $\mathcal{L}$ and $\mathcal{L}_\perp$ are monotone and orthogonal, $\dim(\mathcal{L}) = n(n+1)/2$ and $\dim(\mathcal{L}_\perp) = n(n-1)/2$, we easily see that $\mathcal{W}$ is a monotone affine subspace of $\Re^{n \times n} \times \Re^{n \times n}$ of dimension n^2 . The result now follows from Lemma 2.1. ■

3 Technical Results

In this section we provide some technical results which will be used to establish the polynomial convergence of the algorithms presented in Section 4.

We assume throughout this section that $(X, S) \in \mathcal{L}_{++}$ and that $(\Delta X, \widetilde{\Delta X}, \Delta S, \widetilde{\Delta S})$ is a solution of system (4) with $\hat{\mu} = \sigma\mu$ for some $\mu > 0$ and $\sigma \in [0, 1]$. Moreover, we let

$$W_x \equiv X^{-1/2} [\Delta XS + X\Delta S + XS - \sigma\mu I] X^{1/2}, \quad (6)$$

and, for every $\alpha \in \Re$,

$$X(\alpha) \equiv X + \alpha\Delta X, \quad S(\alpha) \equiv S + \alpha\Delta S, \quad (7)$$

$$\mu(\alpha) \equiv (1 - \alpha + \sigma\alpha)\mu. \quad (8)$$

Lemma 3.1 *We have*

$$W_x = -X^{-1/2} [\widetilde{\Delta X}S + X\widetilde{\Delta S}] X^{1/2}, \quad (9)$$

and, for every $\alpha \in \Re$,

$$\begin{aligned} X^{-1/2} [X(\alpha)S(\alpha) - \mu(\alpha)I] X^{1/2} &= (1 - \alpha) (X^{1/2}SX^{1/2} - \mu I) + \alpha W_x \\ &\quad + \alpha^2 X^{-1/2} \Delta X \Delta S X^{1/2}. \end{aligned} \quad (10)$$

Proof. Relation (9) follows immediately from (6) and (4a). Relation (10) follows from (7), (6) and (8) by simple algebraic manipulation. ■

For a nonsingular matrix $P \in \Re^{n \times n}$, consider the following operator $H_P : \Re^{n \times n} \rightarrow \mathcal{S}^n$ defined as

$$H_P(M) \equiv \frac{1}{2} [PMP^{-1} + (PMP^{-1})^T], \quad \forall M \in \Re^{n \times n}.$$

The operator H_P has been recently used by Zhang [31] to characterize the central path of SDP problems. The next two lemmas, due to Monteiro (see Lemma 2.1 and Lemma 3.5 of [15]), play a crucial role in our analysis.

Lemma 3.2 *Suppose that $(X, S) \in \mathcal{S}_{++}^n \times \mathcal{S}_{++}^n$ and $Q \in \Re^{n \times n}$ is a nonsingular matrix. Then, for every $\mu \in \Re$, we have*

$$\|X^{1/2}SX^{1/2} - \mu I\|_F \leq \|H_Q(XS - \mu I)\|_F,$$

with equality holding if $QXSQ^{-1} \in \mathcal{S}^n$.

Lemma 3.3 Let $W \in \mathfrak{R}^{n \times n}$ be such that $H_Q(W) = 0$ for some nonsingular $Q \in \mathfrak{R}^{n \times n}$. Then,

$$\|H_I(W)\| \leq \frac{1}{2} \|W - W^T\|_F, \quad (11)$$

$$\|W\|_F \leq \frac{\sqrt{2}}{2} \|W - W^T\|_F. \quad (12)$$

In particular, if $W = U_1 + U_2$ for some $U_1 \in \mathcal{S}^n$ and $U_2 \in \mathfrak{R}^{n \times n}$, then

$$\|W\|_F \leq \sqrt{2} \|U_2\|_F.$$

With the aid of the last lemma, we can now prove the following result.

Lemma 3.4 For every $\theta \in \mathfrak{R}$ and $\alpha \in [0, 1]$, we have

$$\begin{aligned} \|X^{-1/2} [X(\alpha)S(\alpha) - \mu(\alpha)I] X^{1/2}\|_F &\leq (1 - \alpha) \|X^{1/2}SX^{1/2} - \mu I\|_F + \alpha^2 \delta_x \delta_s \\ &\quad + \alpha(1 + 2\sqrt{2}) \max\{\delta_x, \tilde{\delta}_x\} \|X^{1/2}SX^{1/2} - \theta\mu I\|, \end{aligned} \quad (13)$$

where

$$\delta_x \equiv \|X^{-1/2}\Delta X X^{-1/2}\|_F, \quad \tilde{\delta}_x \equiv \|X^{-1/2}\widetilde{\Delta X} X^{-1/2}\|_F, \quad \delta_s \equiv \|X^{1/2}\Delta S X^{1/2}\|_F. \quad (14)$$

Proof. By (10) and (14), we have for every $\alpha \in [0, 1]$ that

$$\|X^{-1/2} [X(\alpha)S(\alpha) - \mu(\alpha)I] X^{1/2}\|_F \leq (1 - \alpha) \|X^{1/2}SX^{1/2} - \mu I\|_F + \alpha \|W_x\|_F + \alpha^2 \delta_x \delta_s. \quad (15)$$

Given $\theta \in \mathfrak{R}$, define

$$W \equiv W_x + X^{-1/2}\widetilde{\Delta X} X^{-1/2} (X^{1/2}SX^{1/2} - \theta\mu I). \quad (16)$$

Using (9), we easily see that

$$W = -X^{1/2}\widetilde{\Delta S} X^{1/2} - \theta\mu X^{-1/2}\widetilde{\Delta X} X^{-1/2},$$

and hence that $W \in \mathcal{S}^\perp$, due to the fact that $\widetilde{\Delta X}, \widetilde{\Delta S} \in \mathcal{S}^\perp$. Moreover, using (6) and (16), we easily see that $W = U_1 + U_2$, where

$$\begin{aligned} U_1 &\equiv X^{1/2}\Delta S X^{1/2} + \theta\mu X^{-1/2}\Delta X X^{-1/2} + X^{1/2}SX^{1/2} - \sigma\mu I, \\ U_2 &\equiv X^{-1/2}(\Delta X + \widetilde{\Delta X}) X^{-1/2} (X^{1/2}SX^{1/2} - \theta\mu I). \end{aligned} \quad (17)$$

Clearly, $U_1 \in \mathcal{S}^n$ since $\Delta X, \Delta S \in \mathcal{S}^n$. Noting that $W \in \mathcal{S}^\perp$ is equivalent to $H_I(W) = 0$, it follows that W , U_1 and U_2 satisfy the assumptions of Lemma 3.3 with $Q = I$. Hence, by Lemma (3.3), (17) and (14), we obtain

$$\begin{aligned} \|W\|_F &\leq \sqrt{2} \|U_2\|_F \leq \sqrt{2} \|X^{-1/2}(\Delta X + \widetilde{\Delta X}) X^{-1/2}\|_F \|X^{1/2}SX^{1/2} - \theta\mu I\| \\ &\leq \sqrt{2}(\delta_x + \tilde{\delta}_x) \|X^{1/2}SX^{1/2} - \theta\mu I\|, \end{aligned}$$

which together with (16) and (14) imply

$$\begin{aligned}\|W_x\| &\leq \|W\|_F + \|X^{-1/2}\widetilde{\Delta X}X^{-1/2}\|_F \|X^{1/2}SX^{1/2} - \theta\mu I\| \\ &\leq [\tilde{\delta}_x + \sqrt{2}(\delta_x + \tilde{\delta}_x)] \|X^{1/2}SX^{1/2} - \theta\mu I\| \\ &\leq (1 + 2\sqrt{2}) \max\{\delta_x, \tilde{\delta}_x\} \|X^{1/2}SX^{1/2} - \theta\mu I\|.\end{aligned}$$

This inequality together with (15) now yield (13). $\blacksquare$

The proof of next lemma is straightforward and therefore we omit the details.

Lemma 3.5 *If $(X, S) \in \mathcal{N}_F(\mu, \gamma)$ for some $\gamma \in (0, 1)$, then*

$$\|X^{1/2}S^{1/2}\|^2 \leq (1 + \gamma)\mu, \quad (18)$$

$$\|X^{-1/2}S^{-1/2}\|^2 \leq [(1 - \gamma)\mu]^{-1}, \quad (19)$$

$$\|X^{1/2}SX^{1/2} - \theta\mu I\|_F \leq (\gamma + (1 - \theta)\sqrt{n})\mu, \quad \text{for any } \theta \in [0, 1], \quad (20)$$

$$(1 - \gamma)n\mu \leq X \bullet S \leq (1 + \gamma)n\mu. \quad (21)$$

The next result gives bounds on the quantities δ_x , $\tilde{\delta}_x$ and δ_s defined in (14).

Lemma 3.6 *If $(X, S) \in \mathcal{N}_F(\mu, \gamma)$ for some $\gamma \in (0, 1)$, then*

$$\begin{aligned}\max\{\delta_x, \tilde{\delta}_x\} &\leq \frac{\gamma + (1 - \sigma)\sqrt{n}}{1 - \gamma}, \\ \delta_s &\leq \frac{\gamma + (1 - \sigma)\sqrt{n}}{1 - \gamma}\mu,\end{aligned}$$

where δ_x , $\tilde{\delta}_x$ and δ_s are defined in (14).

Proof. Multiplying (4a) on the left by $X^{-1/2}$ and on the right by $S^{-1/2}$, squaring both sides of the resulting equation and noting the fact that $(\Delta X + \widetilde{\Delta X}) \bullet (\Delta S + \widetilde{\Delta S}) \geq 0$, we obtain

$$\|X^{-1/2}(\Delta X + \widetilde{\Delta X})S^{1/2}\|_F^2 + \|X^{1/2}(\Delta S + \widetilde{\Delta S})S^{-1/2}\|_F^2 \leq \|X^{1/2}S^{1/2} - \sigma\mu X^{-1/2}S^{-1/2}\|_F^2, \quad (22)$$

Using the fact that for any $M \in \mathbb{R}^{n \times n}$,

$$\frac{\|M + M^T\|_F}{2} \leq \|M\|_F, \quad \frac{\|M - M^T\|_F}{2} \leq \|M\|_F,$$

relations (14) and (22), and Lemma 3.5, we obtain

$$\begin{aligned}\delta_s &= \|X^{1/2}\Delta SX^{1/2}\|_F \leq \|X^{1/2}(\Delta S + \widetilde{\Delta S})X^{1/2}\|_F \\ &\leq \|X^{1/2}(\Delta S + \widetilde{\Delta S})S^{-1/2}\|_F \|S^{1/2}X^{1/2}\| \\ &\leq \|X^{1/2}S^{1/2} - \sigma\mu X^{-1/2}S^{-1/2}\|_F \|S^{1/2}X^{1/2}\| \\ &\leq \|X^{1/2}SX^{1/2} - \sigma\mu I\|_F \|X^{-1/2}S^{-1/2}\| \|X^{1/2}S^{1/2}\| \\ &\leq \left(\frac{1 + \gamma}{1 - \gamma}\right)^{1/2} (\gamma + (1 - \sigma)\sqrt{n})\mu \leq \frac{\gamma + (1 - \sigma)\sqrt{n}}{1 - \gamma}\mu,\end{aligned}$$

and

$$\begin{aligned}
\max\{\delta_x, \tilde{\delta}_x\} &\leq \max\left\{\left\|X^{-1/2}\Delta X X^{-1/2}\right\|_F, \left\|X^{-1/2}\widetilde{\Delta X} X^{-1/2}\right\|_F\right\} \\
&\leq \left\|X^{-1/2}(\Delta X + \widetilde{\Delta X})X^{-1/2}\right\|_F \\
&\leq \left\|X^{-1/2}(\Delta X + \widetilde{\Delta X})S^{1/2}\right\|_F \left\|S^{-1/2}X^{-1/2}\right\| \\
&\leq \left\|X^{1/2}S^{1/2} - \sigma\mu X^{-1/2}S^{-1/2}\right\|_F \left\|X^{-1/2}S^{-1/2}\right\| \\
&\leq \left\|X^{1/2}SX^{1/2} - \sigma\mu I\right\|_F \left\|X^{-1/2}S^{-1/2}\right\|^2 \\
&\leq \frac{\gamma + (1 - \sigma)\sqrt{n}}{1 - \gamma}.
\end{aligned}$$

■

Now we are ready to state the main result of this section.

Lemma 3.7 *Suppose that $(X, S) \in \mathcal{N}_F(\mu, \gamma)$ for some $\gamma \in (0, 1)$ and let $(\Delta X, \widetilde{\Delta X}, \Delta S, \widetilde{\Delta S})$ be the solution of (4). Then,*

$$\begin{aligned}
&\left\|X^{-1/2}[X(\alpha)S(\alpha) - \mu(\alpha)]X^{1/2}\right\|_F \\
&\leq \left\{ (1 - \alpha)\gamma + \alpha(1 + 2\sqrt{2})\gamma \frac{\gamma + (1 - \sigma)\sqrt{n}}{1 - \gamma} + \alpha^2 \left(\frac{\gamma + (1 - \sigma)\sqrt{n}}{1 - \gamma} \right)^2 \right\} \mu.
\end{aligned}$$

Proof. Follows immediately from (13) with $\theta = 1$, the assumption that $(X, S) \in \mathcal{N}_F(\mu, \gamma)$ and Lemma 3.6. ■

4 Algorithms

In this section, we establish polynomial iteration-complexity bounds for two primal-dual feasible interior-point algorithms for SDLCP based on the KSH family of search directions given by (4). Both algorithms are extensions of well-known algorithms for linear programming: the first one is a short-step path-following method which generalizes the algorithms presented in Kojima, Mizuno and Yoshise [6] and Monteiro and Adler [16, 17]; the second one is a predictor-corrector algorithm similar to the predictor-corrector LP method of Mizuno, Todd and Ye [13].

4.1 Short-step path following algorithm

In this subsection, we analyze the polynomial convergence of a short-step path following algorithm based on the KSH family of search directions.

We start by stating the algorithm that will be considered in this subsection.

Algorithm-I:

Choose constants γ and δ in $(0, 1)$ satisfying the conditions of Theorem 4.2 below and let $\sigma \equiv 1 - \delta/\sqrt{n}$. Let $\mu_0 > 0$ and $(X^0, S^0) \in \mathcal{L}_{++}$ be such that $(X^0, S^0) \in \mathcal{N}_F(\mu_0, \gamma)$. Let $L > 1$.

Repeat until $\mu_k \leq 2^{-L}\mu_0$, **do**

- (1) Choose a linear subspace $\mathcal{L}_\perp^k$ satisfying [A3].
- (2) Compute the solution $(\Delta X^k, \widetilde{\Delta X}^k, \Delta S^k, \widetilde{\Delta S}^k)$ of system (4) with $(X, S) = (X^k, S^k)$, $\mathcal{L}_\perp = \mathcal{L}_\perp^k$ and $\hat{\mu} = \sigma\mu_k$;
- (3) Set $(X^{k+1}, S^{k+1}) \equiv (X^k, S^k) + (\Delta X^k, \Delta S^k)$ and $\mu_{k+1} = \sigma\mu_k$;
- (4) Increment k by 1.

End

When the constant Γ defined in (23) is such that $\Gamma \leq \gamma$, the lemma below implies that the sequence $\{(X^k, S^k)\}$ generated by Algorithm-I is contained in the neighborhood $\mathcal{N}_F(\gamma)$. This lemma is also used in the analysis of the corrector (or centering) steps of the predictor-corrector algorithm presented in the next subsection.

Lemma 4.1 *Let $\gamma \in (0, 1)$ and $\delta \in [0, n^{1/2})$ be constants satisfying*

$$\Gamma \equiv 5 \left(\frac{\gamma + \delta}{1 - \gamma} \right)^2 \left(1 - \frac{\delta}{\sqrt{n}} \right)^{-1} < 1. \quad (23)$$

Suppose that $(X, S) \in \mathcal{N}_F(\mu, \gamma)$ for some $\mu > 0$, and $(\Delta X, \widetilde{\Delta X}, \Delta S, \widetilde{\Delta S})$ is the solution of system (4) with $\hat{\mu} = \sigma\mu$ and $\sigma = 1 - \delta/\sqrt{n}$. Then, $(X + \Delta X, S + \Delta S) \in \mathcal{N}_F(\sigma\mu, \Gamma)$.

Proof. It follows from Lemma 3.7, the definition of σ and (23) that for every $\alpha \in [0, 1]$,

$$\begin{aligned} & \left\| X^{-1/2} [X(\alpha)S(\alpha) - \mu(\alpha)] X^{1/2} \right\|_F \\ & \leq \left\{ (1 - \alpha)\gamma + \alpha(1 + 2\sqrt{2})\gamma \frac{\gamma + (1 - \sigma)\sqrt{n}}{1 - \gamma} + \alpha^2 \left(\frac{\gamma + (1 - \sigma)\sqrt{n}}{1 - \gamma} \right)^2 \right\} \mu \\ & = (1 - \alpha)\gamma\mu + \left\{ \alpha(1 + 2\sqrt{2})\gamma \frac{\gamma + \delta}{1 - \gamma} + \alpha^2 \left(\frac{\gamma + \delta}{1 - \gamma} \right)^2 \right\} \mu \\ & \leq (1 - \alpha)\gamma\mu + 5\alpha \left(\frac{\gamma + \delta}{1 - \gamma} \right)^2 \mu \leq (1 - \alpha)\gamma\mu + 5\alpha\Gamma \left(1 - \frac{\delta}{\sqrt{n}} \right) \mu \\ & = \{(1 - \alpha)\gamma + \sigma\Gamma\alpha\} \mu, \end{aligned}$$

and hence, in view of (8) and (23), we have

$$\left\| \frac{X^{-1/2} X(\alpha) S(\alpha) X^{1/2}}{\mu(\alpha)} - I \right\|_F \leq \frac{(1 - \alpha)\gamma + \sigma\Gamma\alpha}{1 - \alpha + \sigma\alpha} \leq \max\{\gamma, \Gamma\} < 1.$$

This implies that $X^{-1/2} X(\alpha) S(\alpha) X^{1/2}$ is invertible for every $\alpha \in (0, 1]$. Hence, $X(\alpha)$ and $S(\alpha)$ are also invertible for every $\alpha \in (0, 1]$. Using the fact that $(X, S) \in \mathcal{L}_{++}$, $(X + \Delta X, S + \Delta S) \in \mathcal{L}$ and

a simple continuity argument, we see $(X(\alpha), S(\alpha)) \in \mathcal{L}_{++}$ for every $\alpha \in (0, 1]$. Applying Lemma 3.2 with $(X, S) = (X(\alpha), S(\alpha))$ and $Q = X^{-1/2}$, we then obtain

$$\begin{aligned} \|X(\alpha)^{1/2}S(\alpha)X(\alpha)^{1/2} - \mu(\alpha)I\|_F &\leq \|H_{X^{-1/2}}(X(\alpha)S(\alpha) - \mu(\alpha)I)\|_F \\ &\leq \|X^{-1/2}X(\alpha)S(\alpha)X^{1/2} - \mu(\alpha)I\|_F \leq \{(1-\alpha)\gamma + \sigma\Gamma\alpha\}\mu. \end{aligned}$$

Setting $\alpha = 1$ in the last relation and using the fact that $(X(1), S(1)) \in \mathcal{L}_{++}$ together with (7) and (8), we conclude that $(X(1), S(1)) \equiv (X + \Delta X, S + \Delta S) \in \mathcal{N}_F(\sigma\mu, \Gamma)$. ■

As an immediate consequence of Lemma 4.1, we have the following convergence result for Algorithm-I.

Theorem 4.2 Suppose that γ and δ are constants in $(0, 1)$ such that Γ defined by (23) satisfies $\Gamma \leq \gamma$. Then, every iterate (X^k, S^k) generated by Algorithm-I is in $\mathcal{N}_F(\mu_k, \gamma) \subseteq \mathcal{N}_F(\gamma)$ and satisfies

$$X^k \bullet S^k \leq \frac{1+\gamma}{1-\gamma} \left(1 - \frac{\delta}{\sqrt{n}}\right)^k (X^0 \bullet S^0). \quad (24)$$

Moreover, Algorithm-I terminates in at most $\mathcal{O}(\sqrt{n}L)$ iterations.

Proof. The proof that every iterate (X^k, S^k) is in $\mathcal{N}_F(\mu_k, \gamma)$ follows immediately from Lemma 4.1 and a simple induction argument. Relation (24) follows from the fact that $\mu_k = \sigma^k \mu_0$ and relation (21). ■

Examples of constants γ and δ satisfying the conditions of Theorem 4.2 are $\gamma = \delta = 1/10$.

4.2 Predictor-corrector algorithm

In this subsection, we give the polynomial convergence analysis of a predictor-corrector algorithm which is a direct extension of the LP predictor-corrector algorithm studied by Mizuno, Todd and Ye [13].

The algorithm considered in this subsection is as follows.

Algorithm-II:

Choose a constant $0 < \tau < 1$ satisfying the conditions of Theorem 4.3 below.

Let $L > 1$ and $(X^0, S^0) \in \mathcal{L}_{++}$ be such that $(X^0, S^0) \in \mathcal{N}_F(\mu_0, \tau)$,

Repeat until $\mu_k \leq 2^{-L}\mu_0$, **do**

- (1) Choose a linear subspace $\mathcal{L}_\perp^k$ satisfying [A3];
- (2) Compute the solution $(\Delta X_P^k, \widetilde{\Delta X}_P^k, \Delta S_P^k, \widetilde{\Delta S}_P^k)$ of system (4) with $(X, S) = (X^k, S^k)$, $\mathcal{L}_\perp = \mathcal{L}_\perp^k$ and $\hat{\mu} = 0$;
- (3) Let $\alpha_k \equiv \max\{\alpha \in [0, 1] : (X^k(\alpha), S^k(\alpha)) \in \mathcal{N}_F((1-\alpha')\mu_k, 2\tau), \forall \alpha' \in [0, \alpha]\}$, where $(X^k(\alpha), S^k(\alpha)) \equiv (X^k + \alpha\Delta X_P^k, S^k + \alpha\Delta S_P^k)$;
- (4) Let $(\hat{X}^k, \hat{S}^k) \equiv (X^k, S^k) + \alpha_k(\Delta X_P^k, \Delta S_P^k)$ and $\mu_{k+1} \equiv (1 - \alpha_k)\mu_k$;
- (5) Choose a linear subspace $\hat{\mathcal{L}}_\perp^k$ satisfying [A3];
- (6) Compute the solution $(\Delta X_C^k, \widetilde{\Delta X}_C^k, \Delta S_C^k, \widetilde{\Delta S}_C^k)$ of system (4) with $(X, S) = (\hat{X}^k, \hat{S}^k)$, $\hat{\mu} = \mu_{k+1}$ and $\mathcal{L}_\perp = \hat{\mathcal{L}}_\perp^k$;
- (7) Set $(X^{k+1}, S^{k+1}) \equiv (\hat{X}^k, \hat{S}^k) + (\Delta X_C^k, \Delta S_C^k)$;
- (8) Increment k by 1.

End

The following result provides the polynomial convergence analysis of the above algorithm.

Theorem 4.3 Assume that $\tau \in (0, 1/30]$. Then, Algorithm-II satisfies the following statements:

- a) for every $k \geq 0$, $(X^k, S^k) \in \mathcal{N}_F(\tau)$ and $(\hat{X}^k, \hat{S}^k) \in \mathcal{N}_F(2\tau)$;
- b) for every $k \geq 0$, $X^k \bullet S^k \leq \frac{1+\tau}{1-\tau}(1-\bar{\alpha})^k X^0 \bullet S^0$, where $\bar{\alpha} = 1/\mathcal{O}(\sqrt{n})$;
- c) the algorithm terminates in at most $\mathcal{O}(\sqrt{n}L)$ iterations.

Proof. Statement (c) and the well-definedness of Algorithm-II follow directly from (a) and (b). In turn, these two statements follow by a simple induction argument, the two lemmas below and relation (21). ■

The following lemma analyzes the predictor step of Algorithm-II, namely the step described in items (1)-(4) of Algorithm-II.

Lemma 4.4 Suppose that $(X, S) \in \mathcal{N}_F(\mu, \tau)$ for some $\tau \in (0, 1/2)$. For some subspace $\mathcal{L}_\perp$ satisfying [A3], let $(\Delta X, \Delta \hat{X}, \Delta S, \Delta \hat{S})$ denote the solution of (4) with $\hat{\mu} = 0$. Let $\bar{\alpha}$ denote the unique positive root of the second-order polynomial $p(\alpha)$ defined as

$$p(\alpha) = \left(\frac{\tau + \sqrt{n}}{1 - \tau} \right)^2 \alpha^2 + \tau \left[(1 + 2\sqrt{2}) \left(\frac{\tau + \sqrt{n}}{1 - \tau} \right) + 1 \right] \alpha - \tau \quad (25)$$

Then, for any $\alpha \in [0, \bar{\alpha}]$, we have:

$$(X(\alpha), S(\alpha)) \in \mathcal{N}_F((1 - \alpha)\mu, 2\tau). \quad (26)$$

Moreover, $\bar{\alpha} = 1/\mathcal{O}(n^{1/2})$.

Proof. Using Lemma 3.7 with $\gamma = \tau$ and $\sigma = 0$, the fact that $p(\alpha) \leq 0$ for $\alpha \in [0, \bar{\alpha}]$, $\tau \leq 1/2$ and (25), we obtain

$$\begin{aligned} \|X^{-1/2}[X(\alpha)S(\alpha) - \mu(\alpha)]X^{1/2}\|_F &\leq \left\{ (1 - \alpha)\tau + (1 + 2\sqrt{2})\tau \left(\frac{\tau + \sqrt{n}}{1 - \tau} \right) \alpha + \left(\frac{\tau + \sqrt{n}}{1 - \tau} \right)^2 \alpha^2 \right\} \mu \\ &= 2\tau\mu(\alpha) + p(\alpha)\mu \leq 2\tau\mu(\alpha). \end{aligned}$$

An argument similar to the one used in Lemma 4.1 together with (8) and the fact that $2\tau < 1$ and $\hat{\mu} = 0$ (or equivalently, $\sigma = 0$) can be used to show that (26) holds. The assertion that $\bar{\alpha} = 1/\mathcal{O}(n^{1/2})$ follows by a straightforward verification. ■

The following lemma analyzes the corrector step of Algorithm-II, namely the step described in items (5)-(7) of Algorithm-II.

Lemma 4.5 Suppose $(\hat{X}, \hat{S})$ is in $\mathcal{N}_F(\mu, 2\tau)$ for some $\tau \in (0, 1/30]$. Let $(\Delta \hat{X}_C, \Delta \hat{S}_C)$ denote the solution of (4) with $(X, S) = (\hat{X}, \hat{S})$, $\hat{\mu} = \mu$ and $\mathcal{L}_\perp$ satisfying [A3]. Then,

$$(\hat{X}, \hat{S}) + (\Delta \hat{X}_C, \Delta \hat{S}_C) \in \mathcal{N}_F(\mu, \tau).$$

Proof. Follows immediately from Lemma 4.1 with $\sigma = 1$ (or equivalently, $\delta = 0$), $(X, S) = (\hat{X}, \hat{S})$ and $\gamma = 2\tau$, and noting that Γ defined by (23) satisfies $\Gamma \leq \tau$ when $\tau \leq 1/30$. ■

5 Concluding remarks

For simplicity, we have analyzed two algorithms whose sequence $\{\mu_k\}$ in general differs from the sequence of normalized complementarity gaps $\{(X^k \bullet S^k)/n\}$. At the expense of a slightly more complicated analysis, it is possible to develop algorithms similar to the ones presented here in which $\mu_k = (X^k \bullet S^k)/n$ for every k .

The algorithms of this paper are based on the Frobenius neighborhood $\mathcal{N}_F(\gamma)$ of the central path. An interesting topic for future research would be to establish polynomial convergence of algorithms based on the KSH family of search directions which use one of the following two wider neighborhoods of the central path:

$$\begin{aligned} & \left\{ (X, S) \in \mathcal{L}_+ : \|X^{1/2} S X^{1/2} - \mu I\| \leq \gamma \mu, \text{ for some } \mu > 0 \right\}, \\ & \left\{ (X, S) \in \mathcal{L}_+ : \lambda_{\min}(XS - \mu I) \geq -\gamma \mu, \text{ for some } \mu > 0 \right\}. \end{aligned}$$

Acknowledgement

This work was done while the first author was visiting the Institute of Statistical Mathematics and the Tokyo Institute of Technology in Japan. This author is grateful to the first institute for the financial support and, to both institutes, for the congenial scientific atmosphere provided during his stay. The first author would also like to thank the second author and Prof. Masakazu Kojima of the Tokyo Institute of Technology for making his visit to Japan a very enjoyable experience.

References

- [1] F. Alizadeh. Interior point methods in semidefinite programming with applications to combinatorial optimization. *SIAM Journal on Optimization*, 5(1):13–51, 1995.
- [2] F. Alizadeh, J.-P.A. Haeberly, and M.L. Overton. Primal-dual interior-point methods for semidefinite programming. Technical Report 659, Computer Science Department, Courant Institute of Mathematical Sciences, New York University, 1994.
- [3] R. M. Freund. Complexity of an algorithm for finding an approximate solution of a semidefinite program with no regularity condition. Working paper OR 302-94, Operations Research Center, Massachusetts Institute of Technology, Cambridge, December 1994.
- [4] C. Helmberg, F. Rendl, R. J. Vanderbei, and H. Wolkowicz. An interior-point method for semidefinite programming. Manuscript, Program in Statistics and Operations Research, Princeton University, 1994. To appear in *SIAM Journal on Optimization*.
- [5] F. Jarre. An interior-point method for minimizing the maximum eigenvalue of a linear combination of matrices. *SIAM Journal on Control and Optimization*, 31:1360–1377, 1993.
- [6] M. Kojima, S. Mizuno, and A. Yoshise. A polynomial-time algorithm for a class of linear complementarity problems. *Mathematical Programming*, 44:1–26, 1989.

- [7] M. Kojima, S. Mizuno, and A. Yoshise. A primal-dual interior point algorithm for linear programming. In N. Megiddo, editor, *Progress in Mathematical Programming : Interior Point and Related Methods*, pages 29–47. Springer Verlag, New York, 1989.
- [8] M. Kojima, M. Shida, and S. Shindoh. Local convergence of predictor-corrector infeasible-interior-point algorithms for SDPs and SDLCPs. Research Report #B-306, Dept. of Mathematical and Computing Sciences, Tokyo Institute of Technology, 2-12-1 Oh-Okayama, Meguro-ku, Tokyo 152, December 1996.
- [9] M. Kojima, M. Shida, and S. Shindoh. A note on the Nesterov-Todd and the Kojima-Shindoh-Hara search directions in semidefinite programming. Research report #B-313, Dept. of Mathematical and Computing Sciences, Tokyo Institute of Technology, 2-12-1 Oh-Okayama, Meguro-ku, Tokyo 152, Japan, April 1996.
- [10] M. Kojima, S. Shindoh, and S. Hara. Interior-point methods for the monotone semidefinite linear complementarity problem in symmetric matrices. Research Reports on Information Sciences, Ser. B : Operations Research B-282, Dept. of Information Sciences, Tokyo Institute of Technology, 2-12-1 Oh-Okayama, Meguro-ku, Tokyo 152, Japan, April 1994. To appear in *SIAM Journal on Optimization*.
- [11] C-J. Lin and R. Saigal. A predictor-corrector method for semi-definite programming. Working paper, Dept. of Industrial and Operations Engineering, The University of Michigan, Ann Arbor, Michigan 48109-2177, 1995.
- [12] Z-Q. Luo, J. F. Sturm, and S. Zhang. Superlinear convergence of a symmetric primal-dual path-following algorithm for semidefinite programming. Report 9607/A, Econometric Institute, Erasmus University, Rotterdam, The Netherlands, January 1996.
- [13] S. Mizuno, M. J. Todd, and Y. Ye. On adaptive step primal-dual interior-point algorithms for linear programming. *Mathematics of Operations Research*, 18:945–981, 1993.
- [14] R. D. C. Monteiro. Primal-dual path following algorithms for semidefinite programming. manuscript, School of ISyE, Georgia Institute of Technology, Atlanta, GA 30332, USA, September 1995. To appear in *SIAM J. on Optimization*.
- [15] R. D. C. Monteiro. Polynomial convergence of primal-dual algorithms for semidefinite programming based on Monteiro and Zhang family of directions. manuscript, School of ISyE, Georgia Institute of Technology, Atlanta, GA 30332, USA, July 1996.
- [16] R. D. C. Monteiro and I. Adler. Interior path-following primal-dual algorithms. Part I: Linear programming. *Mathematical Programming*, 44:27–41, 1989.
- [17] R. D. C. Monteiro and I. Adler. Interior path-following primal-dual algorithms. Part II: Convex quadratic programming. *Mathematical Programming*, 44:43–66, 1989.
- [18] R. D. C. Monteiro and Y. Zhang. A unified analysis for a class of path-following primal-dual interior-point algorithms for semidefinite programming. manuscript, School of ISyE, Georgia Institute of Technology, Atlanta, GA 30332, USA, June 1996.

- [19] Y. E. Nesterov and A. S. Nemirovskii. A general approach to the design of optimal methods for smooth convex functions minimization. *Ekonomika i Matem. Metody*, 24:509–517, 1988. (In Russian; English transl. Matekon: Translations of Russian and East European Math. Economics.).
- [20] Y. E. Nesterov and A. S. Nemirovskii. Self-concordant functions and polynomial time methods in convex programming. preprint, Central Economic & Mathematical Institute, USSR Acad. Sci. Moscow, USSR, 1989.
- [21] Y. E. Nesterov and A. S. Nemirovskii. Optimization over positive semidefinite matrices: Mathematical background and user's manual. Technical report, Central Economic & Mathematical Institute, USSR Acad. Sci. Moscow, USSR, 1990.
- [22] Y. E. Nesterov and A. S. Nemirovskii. *Interior Point Methods in Convex Programming: Theory and Applications*. Society for Industrial and Applied Mathematics, Philadelphia, 1994.
- [23] Y. E. Nesterov and M. Todd. Primal-dual interior-point methods for self-scaled cones. Technical Report 1125, School of Operations Research and Industrial Engineering, Cornell University, Ithaca, New York, 14853-3801, 1995.
- [24] Y. E. Nesterov and M. Todd. Self-scaled barriers and interior-point methods for convex programming. Technical Report 1091, School of Operations Research and Industrial Engineering, Cornell University, Ithaca, New York, 14853-3801, 1995.
- [25] F. A. Potra and R. Sheng. A superlinearly convergent primal-dual infeasible-interior-point algorithm for semidefinite programming. Reports on Computational Mathematics 78, Department of Mathematics, The University of Iowa, Iowa City, Iowa, October 1995.
- [26] J. F. Sturm and S. Zhang. Symmetric primal-dual path-following algorithms for semidefinite programming. Report 9554/A, Econometric Institute, Erasmus University, Rotterdam, The Netherlands, November 1995.
- [27] M. J. Todd, K. C. Toh, and R. H. Tütüncü. On the Nesterov-Todd direction in semidefinite programming. Technical Report, School of Operations Research and Industrial Engineering, Cornell University, Ithaca, NY 14853, USA, 1996.
- [28] P. Tseng. Search directions and convergence analysis of some infeasible path-following methods for the monotone semi-definite LCP. manuscript, Department of Mathematics, University of Washington, Seattle, Washington, 98195, USA, June 1996.
- [29] L. Vandenberghe and S. Boyd. A primal-dual potential reduction method for problems involving matrix inequalities. *Mathematical Programming*, 69:205–236, 1995.
- [30] Y. Ye. A class of projective transformations for linear programming. *SIAM Journal on Computing*, 19:457–466, 1990.
- [31] Y. Zhang. On extending primal-dual interior-point algorithms from linear programming to semidefinite programming. Technical Report TR 95-20, Dept. of Math/Stat, University of Maryland Baltimore County, Baltimore, MD 21228, USA, 1995.