

# Structures of Lyapunov Regular Sets with Non-Zero Exponents

Masahiro KURATA  
 Nagoya Institute of Technology

## 1 Introduction

Let  $M$  be a closed manifold with a Riemannian metric and  $f; M \rightarrow M$  be a  $C^2$ -diffeomorphism. For  $\lambda, \mu > 0$  and  $1 \leq k \leq \dim M - 1$ , we denote by  $\Lambda = \Lambda(-\mu, \lambda, k)$  the set consisting of Lyapunov regular points  $x$  such that

1. Lyapunov exponents of  $x$  are less than  $-\mu$  or greater than  $\lambda$ ,
2. the dimension of the fiber of stable bundle  $E^s(x)$  equals  $k$ .

Without loss of generality, we may assume that  $\Lambda$  has a dense orbit. In this note we construct a "symbolic dynamics" that represents  $\Lambda$ , and study the structure of  $\Lambda$  in the case when  $\Lambda$  is a fractal set.

## 2 symbolic dynamics of $\Lambda$

The set  $\Lambda$  is represented by a "symbolic dynamics" as follows.

**I** Let  $\mathcal{W} = \{W_n\}_{n \geq 0}$  be a family of sets  $W_n$  of words of length  $n + 1$  such that

1. There are symbols  $\{B_1, \dots, B_p, C_1, \dots, C_q\}$ , and for any  $n \geq 0$

$$W_n \subset \{B_1, \dots, B_p, C_1, \dots, C_q\}^{n+1},$$

2.  $(\alpha_0, \dots, \alpha_n) \in W_n$  implies  $\alpha_0, \alpha_n \in \{B_1, \dots, B_p\}$ ,
3. If  $(\alpha_0, \dots, \alpha_n) \in W_n, (\beta_0, \dots, \beta_m) \in W_m$  and  $\alpha_n = \beta_0$ , then  $(\alpha_0, \dots, \alpha_n, \beta_1, \dots, \beta_m) \in W_{n+m}$ ,
4. for any  $B_i, B_j$  ( $1 \leq i, j \leq p$ ), there are  $n \geq 1$  and  $(\alpha_0, \dots, \alpha_n) \in W_n$  with  $\alpha_0 = B_i, \alpha_n = B_j$ .

**II** For a family of sets of words  $\mathcal{W} = \{W_n\}_{n \geq 0}$  as above, we define a subset of shift (not necessarily subshift)  $\Sigma = \Sigma(\mathcal{W})$  as follows:

1.  $\Sigma = \Sigma(\mathcal{W}) \subset \{B_1, \dots, B_p, C_1, \dots, C_q\}^{\mathbb{Z}}$
2.  $\Sigma$  is generated by  $\mathcal{W} = \{W_n\}_{n \geq 0}$ , that is,
  - (a)  $\underline{\alpha} = (\alpha_n)_{n \in \mathbb{Z}} \in \Sigma$  if and only if for any  $N > 0$  there are  $m, n \geq N$  such that  $(\alpha_{-m}, \alpha_{-m+1}, \dots, \alpha_0, \dots, \alpha_n) \in W_{m+n}$

(b)  $\alpha_0 \in \{B_0, \dots, B_p\}$

**III** We denote by  $\Sigma$  a collection of  $\Sigma = \Sigma(\mathcal{W})$ , where  $\mathcal{W} = \{W_n\}_{n \geq 0}$ , is defined in **I** and **II**. And an equivalence relation  $\sim$  is defined in the disjoint union  $\cup \Sigma(\mathcal{W})$  as follows:

if  $\underline{a} \sim \underline{b}$  for  $\underline{a} \in \Sigma, \underline{b} \in \Sigma'$  and  $\sigma^n(\underline{a}) \in \Sigma, \sigma^n(\underline{b}) \in \Sigma'$   
then  $\sigma^n(\underline{a}) \sim \sigma^n(\underline{b})$ , where  $\sigma$  is the shift map.

**IV** For the quotient space  $\tilde{\Sigma} = \cup \Sigma(\mathcal{W}) / \sim$ , a shift map  $\tilde{\sigma} : \tilde{\Sigma} \rightarrow \tilde{\Sigma}$  is defined as follows:

for  $\underline{\alpha} = (\alpha_n)_n \in \Sigma$  with  $\alpha_0, \alpha_k \in \{B_1, \dots, B_p\}$ , we have  $\tilde{\alpha}^k[\underline{\alpha}] = [\underline{\beta}]$ , where  $\underline{\beta} = (\beta_n)_n \in \Sigma$  is given by  $\beta_n = \alpha_{n+k}$ .

With the notations as above, we have the following.

**Theorem 2.1.** For a Lyapunov regular set  $\Lambda = \Lambda(-\mu, \lambda, k)$ , there are

1. a collection  $\Sigma = \{\Sigma(\mathcal{W})\}$  of countable subsets of shifts  $\Sigma(\mathcal{W})$ ,
2. an equivalence relation  $\sim$  on  $\cup \Sigma(\mathcal{W})$  and the shift map  $\tilde{\sigma} : \tilde{\Sigma} = \cup \Sigma(\mathcal{W}) / \sim \rightarrow \tilde{\Sigma}$ ,
3. a collection of maps  $\Psi = \Psi_\Sigma : \Sigma \rightarrow \Lambda$  (for  $\Sigma \in \Sigma$ ) which is compatible with the equivalence relation  $\sim$ ,

such that the map

$$\tilde{\Psi} : \tilde{\Sigma} \longrightarrow \Lambda$$

induced from  $\{\Psi = \Psi_\Sigma \mid \Sigma \in \Sigma\}$  is surjective and the diagram

$$\begin{array}{ccc} \tilde{\Sigma} & \xrightarrow{\tilde{\sigma}} & \tilde{\Sigma} \\ \tilde{\Psi} \downarrow & & \downarrow \tilde{\Psi} \\ \Lambda & \xrightarrow{f|_\Lambda} & \Lambda \end{array}$$

is commutative.

**Remark 2.1.** Let  $\varepsilon$  be an arbitrary positive number. In Theorem 2.1. we may choose the symbols  $\{B_1, \dots, B_p, C_1, \dots, C_q\}$  of any  $\mathcal{W}$  and maps  $\Psi = \Psi_\Sigma$  such that

1.  $p = 1$ ,
2.  $\text{diam } \Psi(\Sigma) < \varepsilon$  for  $\Sigma = \Sigma(\mathcal{W}) \in \Sigma$ ,
3. for  $x = \Psi((\alpha_n)_n) \in \Psi(\mathcal{W})$   
 $\|Tf^n | E^s(x)\| < \exp(-\mu n)$  if  $\alpha_0 = \alpha_n = B_1$ ,  
 $\|Tf^{-n} | E^u(x)\| < \exp(-\lambda n)$  if  $\alpha_{-n} = \alpha_0 = B_1$ .

### 3 Locally self-similarity with countable contractions

Let  $\Lambda = \Lambda(-\mu, \lambda, k)$  be a Lyapunov regular set. In the sequel we assume that  $\Sigma = \{\Sigma\}$ , where  $\Sigma = \Sigma(\mathcal{W})$ , and maps  $\Psi = \Psi_\Sigma : \Sigma \rightarrow \Lambda$  ( $\Sigma \in \Sigma$ ) are given as in section 2 and satisfy Remark 2.1.

Then  $\Lambda$  is a countable union of closed sets:

$$\Lambda = \bigcup_{\Sigma \in \Sigma} \Psi(\Sigma).$$

In this section we consider the structure of  $\Psi(\Sigma)$ .

Let  $G_k(\mathcal{W})$  be the set of generators of  $\mathcal{W}$ , that is,

$$G_k(\mathcal{W}) = \{(a_0, \dots, a_k) \in W_k \mid a_0 = a_k = B_1, a_i \in \{C_1, \dots, C_q\} \ 1 \leq i \leq k-1\},$$

$$G(\mathcal{W}) = \bigcup_k G_k(\mathcal{W}).$$

For  $\mathbf{a} = (a_0, \dots, a_k) \in G(\mathcal{W})$ , the right contraction

$$R(\mathbf{a}) : \Psi(\Sigma) \longrightarrow \Psi(\Sigma)$$

is defined by

$$R(\mathbf{a})(\Psi((\alpha_n)_n)) = \Psi((\beta_n)_n) \quad \text{for } (\alpha_n)_n \in \Sigma,$$

where

$$\beta_n = \begin{cases} \alpha_{n-k}, & k \leq n, \\ a_n, & 0 \leq n \leq k, \\ \alpha_n, & n \leq 0. \end{cases}$$

Similarly the left contraction

$$L(\mathbf{a}) : \Psi(\Sigma) \longrightarrow \Psi(\Sigma)$$

is defined by

$$L(\mathbf{a})(\Psi((\alpha_n)_n)) = \Psi((\beta_n)_n) \quad \text{for } (\alpha_n)_n \in \Sigma,$$

where

$$\beta_n = \begin{cases} \alpha_n, & 0 \leq n, \\ a_{n+k}, & -k \leq n \leq 0, \\ \alpha_{n+k}, & n \leq -k. \end{cases}$$

Then we have the following.

**Proposition 3.1.** *The set  $\Psi(\Sigma)$  is a countable union of images of  $\Psi(\Sigma)$  by maps  $R(\mathbf{a})L(\mathbf{b})$ ;*

$$\Psi(\Sigma) = \bigcup_{\mathbf{a}, \mathbf{b} \in G(\mathcal{W})} R(\mathbf{a})L(\mathbf{b})(\Psi(\Sigma)).$$

*If the map  $\Psi = \Psi_\Sigma : \Sigma \rightarrow \Lambda$  is injective, then for any  $\mathbf{a}, \mathbf{b} \in G(\mathcal{W})$  the map*

$$L(\mathbf{b})R(\mathbf{a}) = R(\mathbf{a})L(\mathbf{b}) : \Psi(\Sigma) \longrightarrow \Sigma(\Sigma)$$

*is a contraction. And  $\Psi(\Sigma)$  is self-similar by countable contractions.*

## 4 Hausdorff dimension of local stable manifolds

Form the propositions in the revious section,we have

$$\begin{aligned}
 \Psi(\Sigma) &= \bigcup_{\mathbf{a}_1, \mathbf{b}_1 \in G(\mathcal{W})} R(\mathbf{a}_1)L(\mathbf{b}_1)(\Psi(\Sigma)) \\
 &= \bigcup_{\mathbf{a}_2, \mathbf{b}_2 \in G(\mathcal{W})} \bigcup_{\mathbf{a}_1, \mathbf{b}_1 \in G(\mathcal{W})} R(\mathbf{a}_2)L(\mathbf{b}_2)R(\mathbf{a}_1)L(\mathbf{b}_1)(\Psi(\Sigma)) \\
 &= \bigcup_{\mathbf{a}_1, \mathbf{a}_2 \in G(\mathcal{W})} \bigcup_{\mathbf{b}_1, \mathbf{b}_2 \in G(\mathcal{W})} R(\mathbf{a}_2)R(\mathbf{a}_1)L(\mathbf{b}_2)L(\mathbf{b}_1)(\Psi(\Sigma)) \\
 &= \dots \\
 &= \bigcup_{\mathbf{a}, \mathbf{b} \in G(\mathcal{W})^{\mathbb{N}}} R(\mathbf{a})L(\mathbf{b})(\Psi(\Sigma)),
 \end{aligned}$$

where

$$\mathbf{a} = (\mathbf{a}_1, \mathbf{a}_2, \dots), \quad \mathbf{b} = (\mathbf{b}_1, \mathbf{b}_2, \dots) \in G(\mathcal{W})^{\mathbb{N}},$$

and

$$L(\mathbf{b})(\Psi(\Sigma)) = \bigcap_{N \geq 1} L(\mathbf{b}_N) \cdots L(\mathbf{b}_1)(\Psi(\Sigma)).$$

Besides the set  $L(\mathbf{b})(\Psi(\Sigma))$  coincides with an intersection of a local unstable manifold and  $\Psi(\Sigma)$ .

Let  $\text{Lip}(R(\mathbf{a}) | L(\mathbf{b})(\Psi(\Sigma)))$  be the Lipschitz constant of the map

$$R(\mathbf{a}): L(\mathbf{b})(\Psi(\Sigma)) \longrightarrow L(\mathbf{b})(\Psi(\Sigma)).$$

By choosing  $\varepsilon > 0$  in Remark 2.1 sufficiently small, we have

$$\text{Lip}(R(\mathbf{a}) | L(\mathbf{b})(\Psi(\Sigma))) < \exp(-\lambda n).$$

Because the number of the elements of  $G_k(\mathcal{W})$  is less than or equals  $q^{k-1}$ , this implies the following.

**Proposition 4.1.** For  $\mathbf{b} \in G(\mathcal{W})^{\mathbb{N}}$ , there is  $c(\mathbf{b}) > 0$  such that

$$\sum_{\mathbf{a} \in G(\mathcal{W})} \text{Lip}(R(\mathbf{a}) | L(\mathbf{b})(\Psi(\Sigma)))^{c(\mathbf{b})} = 1.$$

The number  $c(\mathbf{b})$  dominates the Hausdorff dimension of the intersection of the local unstable manifold and  $\Psi(\Sigma)$ :

**Proposition 4.2.** The Hausdorff dimension of  $L(\mathbf{b})(\Psi(\Sigma))$  is less than or equals  $c(\mathbf{b})$ .

## References

- [1] V.I.Oseledec, *A multiplicative ergodic theorem. Lyapunov characteristic numbers for dynamical systems*, Trans.Moscow Math.soc.**19** (1968),197-231.
- [2] Y.B.Pesin, *Lyapunov characteristic exponent and smooth ergodic theory*, Russian Math.Surveys **32** (1977),no.4,55-114.