

大内 忠 (上智大)

**Growth property of singular solutions of linear partial differential equations in the complex domain in  $\mathbf{C}^{d+1}$**

By Sunao ŌUCHI (Sophia Univ.)

ABSTRACT. Let  $P(z, \partial)$  be a linear partial differential operator with coefficients holomorphic in a neighbourhood  $\Omega$  of  $z = 0$  in  $\mathbf{C}^{d+1}$ . Consider equation  $P(z, \partial)u(z) = f(z)$ , where  $u(z)$  and  $f(z)$  admit singularities on the surface  $\{z_0 = 0\}$ . We assume that  $|f(z)| \leq A|z_0|^c$  in a region  $\Omega(\theta)$  which is sectorial with respect to  $z_0$ . The main result of this paper is the following:

There is an exponent  $\gamma^*$  such that for some class of operators if  $\forall \varepsilon > 0 \exists C_\varepsilon$  such that  $|u(z)| \leq C_\varepsilon \exp(\varepsilon|z_0|^{-\gamma^*})$  in  $\Omega(\theta)$ , then  $|u(z)| \leq C|z_0|^{c'}$  for some constants  $c'$  and  $C$ .

First we give the notations briefly. The coordinates of  $\mathbf{C}^{d+1}$  are denoted by  $z = (z_0, z_1, \dots, z_d) = (z_0, z') \in \mathbf{C} \times \mathbf{C}^d$ .  $|z| = \max\{|z_i|; 0 \leq i \leq d\}$  and  $|z'| = \max\{|z_i|; 1 \leq i \leq d\}$ . Its dual variables are  $\xi = (\xi_0, \xi') = (\xi_0, \xi_1, \dots, \xi_d)$ .  $\mathbf{N}$  is the set of all nonnegative integers  $\mathbf{N} = \{0, 1, 2, \dots\}$ . The differentiation is denoted by  $\partial_i = \partial/\partial z_i$ , and  $\partial = (\partial_0, \partial_1, \dots, \partial_d) = (\partial_0, \partial')$ . For a multi-index  $\alpha = (\alpha_0, \alpha') \in \mathbf{N} \times \mathbf{N}^d$ ,  $|\alpha| = \alpha_0 + |\alpha'| = \sum_{i=0}^d \alpha_i$ . Define  $\partial^\alpha = \prod_{i=0}^d \partial_i^{\alpha_i}$ . We denote  $\partial^{\alpha'} = \prod_{i=1}^d \partial_i^{\alpha_i}$  by  $\partial^{\alpha'}$ .

We define spaces of holomorphic functions in some regions to state the results. Let  $\Omega = \Omega_0 \times \Omega'$  be a polydisk with  $\Omega_0 = \{z_0 \in \mathbf{C}^1; |z_0| < R\}$  and  $\Omega' = \{z' \in \mathbf{C}^d; |z'| < R\}$  for some positive constant  $R$ . Put  $\Omega_0(\theta) = \{z_0 \in \Omega_0 - \{0\}; |\arg z_0| < \theta\}$  and  $\Omega(\theta) = \Omega_0(\theta) \times \Omega'$ .  $\mathcal{O}(\Omega)$  ( $\mathcal{O}(\Omega')$ ,  $\mathcal{O}(\Omega(\theta))$ ) is the set of all holomorphic functions on  $\Omega$  (resp.  $\Omega'$ ,  $\Omega(\theta)$ ).  $\mathcal{O}(\Omega(\theta))$  contains multi-valued functions, if  $\theta > \pi$ .

We introduce  $\mathcal{O}_{(\kappa)}(\Omega(\theta))$  and  $Asy_{\{\kappa\}}(\Omega(\theta))$ , which are subspaces of  $\mathcal{O}(\Omega(\theta))$  and fundamental function spaces in this paper.

**Definition 1.**  $\mathcal{O}_{(\kappa)}(\Omega(\theta))$  ( $0 < \kappa < +\infty$ ) is the set of all  $u(z) \in \mathcal{O}(\Omega(\theta))$  such that for any  $\varepsilon > 0$  and any  $\theta'$  with  $0 < \theta' < \theta$

$$(1) \quad |u(z)| \leq C \exp(\varepsilon|z_0|^{-\kappa}) \quad z \in \Omega(\theta')$$

holds for a constant  $C = C(\varepsilon, \theta')$ . We put  $\mathcal{O}_{(+\infty)}(\Omega(\theta)) = \mathcal{O}(\Omega(\theta))$  for  $\kappa = +\infty$ .

**Definition 2.**  $\mathcal{O}_{reg,c}(\Omega(\theta))$  is the set of all  $u(z) \in \mathcal{O}(\Omega(\theta))$  such that any  $\theta'$  with  $0 < \theta' < \theta$

$$(2) \quad |u(z)| \leq C|z_0|^c \quad z \in \Omega(\theta')$$

holds for a constant  $C = C(\theta')$ .

We say that  $u(z) \in \mathcal{O}(\Omega(\theta))$  is slowly increasing in  $\Omega(\theta)$ , if  $u(z) \in \bigcup_{|c| < +\infty} \mathcal{O}_{reg,c}(\Omega(\theta))$ .

Now let  $P(z, \partial)$  be an  $m$ -th order linear partial differential equation with coefficients in  $\mathcal{O}(\Omega)$

$$(3) \quad P(z, \partial) = \sum_{|\alpha| \leq m} a_\alpha(z) \partial^\alpha = \sum_{|\alpha| \leq m} z_0^{j_\alpha} b_\alpha(z) \partial^\alpha,$$

where  $j_\alpha \in \mathbf{N}$  is the valuation of  $a_\alpha(z)$  with respect to  $z_0$ ,  $a_\alpha(z) = z_0^{j_\alpha} b_\alpha(z)$ . Let us define some quantities for  $P(z, \partial)$ :

$$(4) \quad \begin{cases} e_* := \min\{j_\alpha - \alpha_0; \alpha \in \mathbf{N}^{d+1}\}, \Delta = \{\alpha \in \mathbf{N}^{d+1}; j_\alpha - \alpha_0 = e_*\} \\ k^* := \max\{|\alpha|; \alpha \in \Delta\}. \end{cases}$$

Put

$$(5) \quad \mathfrak{P}(z, \partial) = \sum_{\alpha \in \Delta} z_0^{j_\alpha} b_{\alpha,0}(z') \partial^\alpha.$$

Let us introduce an index which plays an important role in this paper.

**Definition 3.** (*Minimal irregularity*)

$$(6) \quad \begin{cases} \gamma^* := \min\left\{\frac{j_\alpha - \alpha_0 - e_*}{|\alpha| - k^*}; \alpha \in \mathbf{N}^{d+1}, |\alpha| > k^*\right\}, & \text{if } k^* < m, \\ \gamma^* := \infty, & \text{if } k^* = m. \end{cases}$$

Let us introduce conditions on  $P(z, \partial)$ .

**Condition 0.** If  $\alpha = (\alpha_0, \alpha') \in \Delta$ , then  $\alpha' = (0, 0, \dots, 0)$ .

The following condition is more strict than Condition 0.

**Condition 1.**  $P(z, \partial)$  satisfies Condition 0 and  $b_{(k^*, 0, 0, \dots, 0)}(0) \neq 0$ .

Suppose that  $P(z, \partial)$  satisfies Condition 0. Then  $\mathfrak{P}(z, \partial)$  is an ordinary differential operator,

$$(7) \quad \mathfrak{P}(z, \partial) = \sum_{\alpha \in \Delta} z_0^{e_*} b_{\alpha,0}(z') z_0^{\alpha_0} \partial_0^{\alpha_0},$$

and  $\{z_0 = 0\}$  is regular singular. Define the indicial polynomial  $\chi_P(z', \lambda)$  of  $\mathfrak{P}(z, \partial)$ ,

$$(8) \quad \chi_P(z', \lambda) := \sum_{\alpha \in \Delta} b_{\alpha,0}(z') \lambda(\lambda - 1) \cdots (\lambda - \alpha_0 + 1).$$

Further suppose that  $P(z, \partial)$  satisfies Condition 1. Then  $\chi_P(z', \lambda)$  is a polynomial of  $\lambda$  with degree  $k^*$  in  $\{z; |z| \leq R\}$ . Hence there exist real constants  $a_0, a_1$  and  $b_0$  such that all the roots of  $\chi_P(z', \lambda) = 0$  for  $|z| \leq R$  are contained in  $\{\lambda; a_0 \leq \Re \lambda \leq a_1, |\Im \lambda| \leq b_0\}$ .

Now let us consider

$$(Eq) \quad P(z, \partial)u(z) = f(z).$$

We have results concerning the growth properties of solutions of (Eq).

**Theorem 4.** *Suppose that  $P(z, \partial)$  satisfies Condition 1. Let  $u(z) \in \mathcal{O}_{(\gamma^*)}(\Omega(\theta))$  be a solution of (Eq). Suppose that  $f(z) \in \mathcal{O}_{reg,c}(\Omega(\theta))$ . Then there is a polydisk  $U$  centered at  $z = 0$  such that  $u(z) \in \mathcal{O}_{reg,c'}(U(\theta))$  for any  $c' < \min\{c - e_*, a_0\}$ .*

**Theorem 5.** *Suppose that  $P(z, \partial)$  satisfies Condition 0. Let  $u(z) \in \mathcal{O}_{(\gamma^*)}(\Omega(\theta))$  be a solution of (Eq). Suppose that  $f(z) \in \mathcal{O}_{reg,c}(\Omega(\theta))$ . Then there is a polydisk  $U$  centered at  $z = 0$  and a constant  $c''$  such that  $u(z) \in \mathcal{O}_{reg,c''}(U(\theta))$ .*

We show Theorem 4 by constructing a parametrix and Theorem 5 follows from Theorem 4. The proof theorems and the details of this paper will be appeared in the forthcoming paper.

We give some examples satisfying *Condition 1*:

(a). Operators of normal type with respect to  $\partial_0$ ,

$$\partial_0^{k^*} + \sum_{\alpha_0 < k^*} a_\alpha(z) \partial^\alpha.$$

(b). Operators of Fuchsian type.

(c). Other concrete examples are

$$I_d + z_0^2 \partial_0 + z_0 \partial_1^2, \quad z_0 \partial_0^2 + a(z) \partial_0 + \partial_1^3.$$

The present paper follows Ōuchi [4]. The class of operators considered in [4] was more strict than that of this paper. The main Theorem in [4] was the following:

*If  $u(z)$  grows at most some exponential order near  $z_0 = 0$ , that is, for any  $\varepsilon > 0$   $|u(z)| \leq C_\varepsilon \exp(\varepsilon|z_0|^{-\gamma^*})$  near  $z_0 = 0$ , and if  $f(z)$  behaves asymptotically  $f(z) \sim \sum_{n=0}^{+\infty} f_n(z') z_0^n$  as  $z_0 \rightarrow 0$  in a sectorial region  $\Omega(\theta)$ , where  $|f_n(z')| \leq AB^n \Gamma(n/\gamma^* + 1)$ , then  $u(z)$  has also the asymptotic expansion like  $f(z)$  as  $z_0$  tends to 0.*

It was an extension of the main result of [1] and [2]. But in the present paper we treat a wider class of operators which contains that of [4]. So even if  $f(z)$  has a Gevrey type asymptotic expansion,  $u(z)$  does not always have. Hence, Theorem 4 in this paper is somewhat different. Roughly speaking,

*if  $u(z)$  grows at most some exponential order near  $\{z_0 = 0\}$ , and if  $f(z)$  has the slowly increasing singularities on  $\{z_0 = 0\}$ , then the growth order of singularities of  $u(z)$  are also slowly increasing.*

We can show the results in [4], by using Theorem 4.

#### REFERENCES

- [1] Ōuchi, S., *An integral representation of singular solutions and removable singularities to linear partial differential equations*, Publ. RIMS Kyoto Univ. 26, 735-783 (1990).

- [2] Ōuchi, S., *The behaviour of solutions with singularities on a characteristic surface to linear partial differential equations in the complex domains*, Publ. RIMS Kyoto Univ. 29, 63-120 (1993).
- [3] Ōuchi, S., *Genuine solutions and formal solutions with Gevrey type estimates of nonlinear partial differential equations*, J. Math. Sci. Univ. Tokyo 2 (1995), 375-417.
- [4] Ōuchi, S., *Singular solutions with asymptotic expansion of linear partial differential equations in the complex domain*

Sunao ŌUCHI

Department of Mathematics, Sophia University  
Kioicho Chiyoda-ku, Tokyo 102, JAPAN