

Title	Fat solenoidal attractors (New developments in dynamical systems)
Author(s)	Tsujii, Masato
Citation	数理解析研究所講究録 (2000), 1179: 99-102
Issue Date	2000-12
URL	http://hdl.handle.net/2433/64532
Right	
Type	Departmental Bulletin Paper
Textversion	publisher

Fat solenoidal attractors

Masato TSUJII
Department of Mathematics
Hokkaido University

November 8, 2000

Abstract

We study dynamical systems generated by skew products

$$T : S^1 \times \mathbb{R} \rightarrow S^1 \times \mathbb{R}, \quad T(x, y) = (\ell x, \lambda y + f(x))$$

where $\ell \geq 2$, $1/\ell < \lambda < 1$ and f is a C^2 function on S^1 . We show that the SBR measure for T is absolutely continuous for almost every f .

1 Introduction

In this paper, we study a class of dynamical systems that stably admit an absolutely continuous ergodic measure (*acem*) with a *negative Lyapunov exponent*. It is well-known that expanding dynamical systems generally admit *acem*'s whose Lyapunov exponents are all positive. The aim of this paper is to study another kind of *acem*'s which is produced by a quite different mechanism: *overlap and sliding* in short.

We can find a typical example of such *acem*'s in a paper of Alexander and Yorke[1], where the so-called generalized baker's transformation is considered:

$$B : [-1, 1] \times [-1, 1] \circlearrowleft, \quad B(x, y) = \begin{cases} (2x - 1, \beta y + (1 - \beta)) & x \geq 0 \\ (2x + 1, \beta y - (1 - \beta)) & x < 0. \end{cases}$$

When $\beta = 1/2$, this map B is nothing but the ordinary baker's transformation. Alexander and Yorke studied the case $1/2 < \beta \leq 1$. In such case, the images of left and right halves of the domain, *i.e.*, $B([-1, 0] \times [-1, 1])$ and $B([0, 1] \times [-1, 1])$ overlap with some sliding. This makes the dynamical nature of the map B more complicated and interesting. They observed that the map B admits an *acem* if and only if the number β satisfies a delicate numerical condition: absolute continuity of the corresponding infinitely convoluted Bernoulli measure. As they noted, there are infinitely many numbers in $(1/2, 1]$ (*e.g.* $(\sqrt{5} - 1)/2$) for which B admits no *acem*'s, according to a result of Erdős[2]. On the other hand, B admits an *acem* for Lebesgue almost every β in $(1/2, 1]$ according to a more recent result of Solomyak[3].

In this paper, we consider a class of dynamical systems generated by maps

$$T : S^1 \times \mathbb{R} \rightarrow S^1 \times \mathbb{R}, \quad T(x, y) = (\ell x, \lambda y + f(x)) \quad (1)$$

where $\ell \geq 2$ is an integer, $0 < \lambda < 1$ is a real number, and f is a C^2 function on $S^1 = \mathbb{R}/\mathbb{Z}$. We may regard this class of maps as a conceptual generalization of the generalized baker's transformations B in the sense that the translation in vertical direction depends smoothly on x .

The map T is a skew product on the expanding map $\tau : x \mapsto \ell x$ and it is uniformly contracting in the fiber direction. So T is an Anosov endomorphism. The ergodic property of T is rather simple: there exists an ergodic probability measure μ on $S^1 \times \mathbb{R}$, for which Lebesgue almost every point $\mathbf{x} \in S^1 \times \mathbb{R}$ is generic, that is,

$$\lim_{n \rightarrow \infty} \frac{1}{n} \sum_{i=0}^{n-1} \delta_{T^i(\mathbf{x})} = \mu \quad \text{weakly.}$$

We will call this measure μ the *SBR measure* for T .

The question is smoothness of the SBR measure μ with respect to the Lebesgue measure on $S^1 \times \mathbb{R}$. In the case $\lambda \ell < 1$, the SBR measure is totally singular because T contracts area. The case $\lambda \ell > 1$, which corresponds to the case $\beta > 1/2$ for the generalized baker's transformations, is more interesting. We will focus on this case. First we give two examples in opposite directions.

Example 1 Let $\ell = 2$, $0.5 < \lambda \leq 0.51$ and $f(x) = \sin 2\pi x$. Then the SBR measure μ for T is absolutely continuous with respect to the Lebesgue measure of $S^1 \times \mathbb{R}$. (See figure 1.)

Example 2 If $f(x) = \varphi(\tau(x)) - \lambda\varphi(x)$ for some measurable function φ on S^1 , the SBR measure for T is supported on the graph of φ and totally singular.

We claim that the SBR measure is absolutely continuous for almost every T and, moreover, that the absolute continuity is robust. Fix an integer $\ell \geq 2$. Let $\mathcal{D} \subset (0, 1) \times C^2(S^1, \mathbb{R})$ be the set of combinations (λ, f) for which the SBR measure is absolutely continuous w.r.t. the Lebesgue measure on $S^1 \times \mathbb{R}$. We consider the interior \mathcal{D}° of \mathcal{D} with respect to the topology that is defined as the product of the canonical topology on $(0, 1)$ and C^2 -topology on $C^2(S^1, \mathbb{R})$. The main result of this paper is the following.

Theorem 1 Let $\ell^{-1} < \lambda < 1$. There exists a finite collection of C^∞ functions $\varphi_i : S^1 \rightarrow \mathbb{R}$, $i = 1, 2, \dots, m$, such that, for any C^2 function $g \in C^2(S^1, \mathbb{R})$, the subset of \mathbb{R}^m ,

$$\left\{ (t_1, t_2, \dots, t_m) \in \mathbb{R}^m \mid \left(\lambda, g(x) + \sum_{i=1}^m t_i \varphi_i(x) \right) \notin \mathcal{D}^\circ \right\},$$

is a null set with respect to the Lebesgue measure on \mathbb{R}^m .




Figure 1: The orbit of the point $(0.1, 0)$ up to time 100000 when $\ell = 2$, $\lambda = 0.51$ and $f(x) = \sin(x)$.

As simple consequences, we obtain

Corollary 2 \mathcal{D} contains an open and dense subset of $(1/\ell, 1) \times C^2(S^1, \mathbb{R})$.

Corollary 3 For $\ell^{-1} < \lambda < 1$ and $2 \leq r \leq \infty$, the set of functions

$$\mathcal{D}_\lambda^r = \{f \in C^r(S^1, \mathbb{R}) \mid (\lambda, f) \in \mathcal{D}^\circ\}$$

is an open and dense subset of $C^r(S^1, \mathbb{R})$.

Moreover, the claim of theorem 1 implies that the subset \mathcal{D}_λ^r above occupies *almost everywhere* in $C^r(S^1, \mathbb{R})$. In fact, if $C^r(S^1, \mathbb{R})$ were a finite dimensional Euclidean space, the claim would imply that the subset \mathcal{D}_λ^r had full measure with respect to the 'Lebesgue measure' on $C^r(S^1, \mathbb{R})$. See [5] and [6] for discussions about measure-theoretical conditions that imply "almost everywhere" for subsets in infinite dimensional spaces.

The proof of theorem 1 is based on an idea that transversality of the unstable manifolds leads to absolute continuity of the SBR measure. We took this idea from a paper of Solomyak and Peres[4] where the authors gave a simplified proof of the above mentioned result of Solomyak.

One can download the full paper at

<http://www.math.sci.hokudai.ac.jp/~tsujii/index.html>

References

- [1] J.C.Alexander & J.A. Yorke, *Fat Baker's transformations*, Ergodic theory and dynamical systems, **4**, 1–23 (1984)
- [2] P. Erdős, *On a family of symmetric Bernoulli convolutions*, Amer. J. Math. **61**, 974–976, (1939)
- [3] B. Solomyak, *On the random series $\sum \pm \lambda^n$ (an Erdős problem)*, Ann. of Math. (2) **142**, no. 3, 611–625, (1995)
- [4] B.Solomyak & Y. Peres, *Absolute continuity of Bernoulli convolutions, a simple proof*, Math. Research Letters **3** , 231-239 (1996)
- [5] M. Tsujii, *A measure on the space of smooth mappings and dynamical system theory.*, J. Math. Soc. Japan **44**, no. 3, 415–425, (1992)
- [6] B. R.Hunt, T. Sauer & J. A. Yorke, *Prevalence: a translation-invariant "almost every" on infinite-dimensional spaces*, Bull. Amer. Math. Soc. , **27**, no.2, 217–238(1992); Addendum *ibid.* **28** , no. 2, 306–307, (1993)