TITLE:

Preliminary Surveys of the Commensal Amphipod, *Leucothoe spinicarpa* (Abildgaard, 1789), in the colonial tunicate, *Ecteinascidia thurstoni* Herdman, 1891, in the Andaman Sea, Thailand

AUTHOR(S):

CHAVANICH, SUCHANA; KETDECHA,
NIMMANORADEE; VIYAKARN, VORANOP;
BUSSARAWIT, SOMCHAI

CITATION:

CHAVANICH, SUCHANA ...[et al]. Preliminary Surveys of the Commensal Amphipod, *Leucothoe spinicarpa* (Abildgaard, 1789), in the colonial tunicate, *Ecteinascidia thurstoni* Herdman, 1891, in the Andaman Sea, Thailand. Publications of the Seto Marine Biological Laboratory. Special Publication Series 2007, 8: 97-101

ISSUE DATE:

2007

URL:

<http://hdl.handle.net/2433/70906>

RIGHT:

Preliminary Surveys of the Commensal Amphipod, *Leucothoe spinicarpa* (Abildgaard, 1789), in the colonial tunicate, *Ecteinascidia thurstoni* Herdman, 1891, in the Andaman Sea, Thailand

SUCHANA CHAVANICH¹*, NIMMANORADEE KETDECHA¹, VORANOP VIYAKARN¹
and SOMCHAI BUSSARAWIT²

¹Department of Marine Science, Faculty of Science, Chulalongkorn University, Bangkok, Thailand

²Phuket Marine Biological Center, Phuket, Thailand

Corresponding author's e-mail: suchana.c@chula.ac.th

Abstract Amphipods identified as, *Leucothoe spinicarpa* (Abildgaard, 1789), were found in the colonial tunicate, *Ecteinascidia thurstoni* Herdman, 1891, at 1-3 m depth, within a single coral reef area on the coast of the Andaman Sea of Phuket Province in southern Thailand. This represents the first record of commensalism between this amphipod and tunicate in Thai waters. Host-symbiont occurrences were low, with only 2.2% of all tunicate zooids harboring *L. spinicarpa*, and always with a single amphipod per zooid. All *L. spinicarpa* occurred in the branchial chamber of the tunicate and included female and male specimens. Amphipods found in the tunicates ranged between 0.4-2.1 mm in length.

Key words: amphipod, tunicate, *Leucothoe spinicarpa*, *Ecteinascidia thurstoni*, commensal

Introduction

Amphipods can be found in association with macroalgae and marine invertebrates, using them as habitat, food sources, and as protection from predators (Buschmann, 1990; Chavanich, 2006; Chavanich and Wilson, 2000; Duffy and Hay, 1991; Hacker and Steneck, 1990). Some amphipods living inside host organisms such as tunicates, sponges, and sea anemones are considered commensal symbionts (Thomas, 1979; Vader, 1970; Vader, 1984; Vader and Krapp-Schickel, 1996). They receive benefits from their hosts through the provision of microhabitats and food sources, while not harming their hosts (Jaramillo *et al.*, 1981; Thiel, 1999; Vader, 1985; Vader and Beehler, 1983).

As a group, leucothoids are commensal amphipods usually found inside sponges and ascidians (Barnard and Karaman, 1991; Ortiz, 1975). To date, *Leucothoe spinicarpa* (Abildgaard, 1789) is the only species found inside the Thai colonial tunicate, *Ecteinascidia thurstoni* Herdman, 1891. This tunicate has attracted attention in Thailand recently as a new potential source of anti-cancer compounds (Chavanich *et al.*, 2005). A group of alkaloids, the Ecteinascidins (Et), including Et 770 and Et 786, have been isolated with very high yields from *E. thurstoni* pretreated with potassium cyanide. These extracts exhibit potent cytotoxic activity against cancer cells of breast, lung, colorectal, and nasopharyngeal tissues (Suwanborirux *et al.*, 2002).

The purpose of the present study was to investigate the host-symbiont association of *Leucothoe spinicarpa* and *Ecteinascidia thurstoni* in Thai waters, to discover preliminary details of the commensal amphipod populations and ascertain specifics such as the positions inside the tunicates where the amphipods were found.

Materials and Methods

During the 2nd Annual JSPS-NaGISA workshop on the taxonomy of marine amphipods in Nha Trang, Vietnam from September 30-October 3, 2004, the presence of an amphipod inside the tunicate,

Ecteinascidia thurstoni was noted. Subsequent identification during the workshop lead by Dr. Manuel Ortiz Touzet showed this amphipod to be *Leucothoe spinicarpa* (Abildgaard, 1789).

Specimens of the tunicate, *E. thurstoni*, were collected at 1-3 m depth along the coast of Phuket Province, southern Thailand. Tunicates were dissected, and amphipods found inside host specimens were collected and fixed in 5% buffered formalin for later identification. A total of 1500 tunicate zooids were collected between the months of September 2004 and April 2006. In addition, notes on the sex and position of amphipods in the tunicates collected, were recorded. Amphipod specimens were also measured from the anterior edge of the first pereion segment to the posterior edge of the fifth pereion segment.

Results

Leucothoe spinicarpa was the only amphipod found inside the tunicate *Ecteinascidia thurstoni* (Figures 1, 2). 2.2% of the tunicate zooids acted as hosts, with most zooids containing no amphipods

Fig. 1. *Leucothoe spinicarpa* (Abildgaard, 1789).

Fig. 2. The commensal amphipod *Leucothoe spinicarpa* (Abildgaard, 1789) inside the branchial chamber (arrow) of tunicate *Ecteinascidia thurstoni* Herdman, 1891.

Table 1. Percentage of amphipods found inside the tunicate *Ecteinnascidia thurstoni* Herdman, 1891.

Tunicates	Percent of amphipods
without amphipods	97.8%
with amphipods	2.2%
with female amphipods	0.67%
with male amphipods	1.06%

Fig. 3. Size classes of the amphipod *Leucothoe spinicarpa*.

(Table 1). All zooids hosted only a single individual *L. spinicarpa* amphipod. Both female and male amphipods were found, and all *L. spinicarpa* occurred in the branchial chamber part of the tunicate (Table 1). Amphipods found in the tunicates ranged between 0.4-2.1 mm in length (Figure 3).

Discussion

A previous study found three commensal amphipod species in four sponges and one solitary tunicate in Phuket, Thailand (Bahrndorff and Lofstedt, 2004). However, *L. spinicarpa* was not among the amphipods recorded. The present study is the first record of the commensal amphipod *L. spinicarpa* found associated with the colonial tunicate *E. thurstoni* in Thai water. *L. spinicarpa* has also been found inhabiting other tunicate species: *Ecteinnascidia turbinata*, *Styela plicata*, *Ascidia nigra*, *Microcosmos exasperatus*, *Clavelina oblonga* (Thiel, 1999).

The amphipod specimens obtained from inside the tunicates spanned a large size and age range (both juveniles and adults were present). This finding is contrary to Thiel's (1999) report that only small specimens of *L. spinicarpa* live in colonial tunicates while larger ones live in solitary tunicates. He suggested that juveniles first lived with their parents in solitary tunicates, and when mature, sought colonial tunicates as their host. Our results do not match the suggested pattern. Another study by Bahrndorff and Lofstedt (2004) also seems to contradict Thiel (1999) in that it found no *L. spinicarpa* in solitary tunicates. In agreement with other studies (Thiel, 1999; Thomas and Klebba, 2006), we found *L. spinicarpa* occurring in the branchial chamber of the tunicate. This tunicate organ seems to facilitate commensal amphipods in finding food among the materials filtered by the host (Thiel, 1999).

Results on the population of commensal amphipods showed that only 2.2% of all tunicate zooids were host to *L. spinicarpa* (Table 1). Previous studies have reported that high percentages of *L. spinicarpa* were found inside solitary tunicates (77-95% of their hosts) and colonial tunicates (9.4% of their hosts) (Thiel, 1999). Several factors such as choice of preferred hosts and abundance of hosts and other commensal amphipod species might influence the numbers and reproduction of *L. spinicarpa* populations. It appears that more studies are needed to further elucidate the factors involved and to determine the relationship between *L. spinicarpa* and its hosts.

Acknowledgements

We would like to thank Dr. Manuel Ortiz Touzet for helping with the amphipod identification and Piya Koeyin for field assistance. We also thank Dr. Yoshihisa Shirayama for encouragement in the initial project. JSPS and NaGISA provided travel funding to SC and SB for attending the 2nd Annual JSPS-NaGISA Workshop Taxonomy of Marine Amphipods in Nha Trang, Vietnam. We thank the anonymous reviewers for providing useful comments on the manuscript.

References

- Bahrndorff, S. and Lofstedt, M. 2004. Commensal amphipods in ascidians and sponges off Phuket Island. Phuket Marine Biological Center Research Bulletin, 65, 45-54.
- Barnard, J. L. and Karaman, G. S. 1991. Records of the Australian Museum: The Families and Genera of Marine Gammaridean Amphipoda (Except Marine Gammaroids), Part 1 and 2. Australian Museum. 866 pp.
- Buschmann, A. H. 1990. Intertidal macroalgae as refuge and food for Amphipoda in central Chile. Aquatic Botany, 36, 237-245.
- Chavanich, S. 2006. The occurrence of *Hyale nilssonii* in the rocky intertidal zone in New Hampshire, U.S.A. Crustaceana, 79, 1005-1010.
- Chavanich, S., Koeyin, P., Viyakarn, V., Piyatiratitvorakul, S., Menasveta, P., Suwanborirux, K., and Poovachiranon, S. 2005. A tunicate from a Thai coral reef: a potential source of new anticancer compounds. Coral Reefs, 24 pp. 621.
- Chavanich, S. and Wilson, K. A. 2000. Rocky intertidal zonation of gammaridean amphipods in Long Island Sound, Connecticut. Crustaceana, 73, 835-846.
- Duffy, J. E. and Hay, M. E. 1991. Food and shelter as determinants of food choice by a herbivorous marine amphipod. Ecology, 72, 1286-1298.
- Hacker, S. D. and Steneck, R. S. 1990. Habitat architecture and the abundance and body-size-dependent habitat selection of a phytal amphipod. Ecology, 71, 2269-2285.
- Jaramillo, E., Navarro, J., and Winter, J. 1981. The association between *Mytilus chilensis* Hupe (Bivalvia, Mytilidae) and *Edotea megellanica* Cunningham (Isopoda, Valvifera) in southern Chile. Biological Bulletin, 160, 107-113.
- Ortiz, L. M. 1975. Algunos datos ecologicos de *Leucothoe spinicarpa* Abildgaard (Amphipoda, Gammaridea) en aguas Cubanas. Ciencias, Series 8, Investigaciones Marinas, 16, 1-12.
- Suwanborirux, K., Charupant, K., Amnuoypol, S., Pumangura, S., Kubo, A., and Saito, N. 2002. Ecteinascidins 770 and 786 from the Thai tunicate *Ecteinascidia thurstoni*. Journal of Natural Products, 65, 935-937.
- Thiel, M. 1999. Host-use and population demographics of the ascidian-dwelling amphipod *Leucothoe spinicarpa*: indication for extended parental care and advanced social behavior. Journal of Natural History, 33, 193-206.
- Thomas, J. D. 1979. Occurrence of the amphipod *Leucothoides pottsi* Shoemaker in the tunicate *Ecteinascidia turbinata* Herdman from Big Pine Key, Florida, USA. Crustaceana, 37, 107-109.
- Thomas, J. D. and Klebba, K. N. 2006. Studies of commensal Leucothoid amphipods: two new sponge-inhabiting species from south Florida and the western Caribbean. Journal of Crustacean Biology, 26, 13-22.
- Vader, W. 1970. The amphipods associated with the sea anemone, *Bolocera tuediae*, in western Norway. Sarsia, 43, 87-98.
- Vader, W. 1984. Notes on Norwegian marine Amphipoda 8. Amphipods found in association with sponges and tunicates. Fauna norvegica, Seria A, 5, 14-16.
- Vader, W. 1985. On the road towards parasitism? *Orchomene recondite* (Stasek, 1958), an amphipod that lives

- inside a sea-anemone. Information Institute of Parasitology Abo Akademi, 18, pp. 49.
- Vader, W. and Beehler, C. L. 1983. *Metopa glacialis* (Amphipoda, Stenothoidea) in the Barents and Beaufort Seas, and its association with the lamellibranches *Musculus niger* and *M. discors* s. 1. *Astarte*, 12, 57-61.
- Vader, W. and Krapp-Schickel, G. 1996. Redescription and biology of *Stenothoe brevicornis* Sars (Amphipoda: Crustacea), an obligate associate of the sea anemone *Actinostola callosa* (Verrill). *Journal of Natural History*, 30, 51-66.