

## NOTES

### Reactions of Ketene with Phenol, Resorcine, Phloroglucine and Dimedone

Ryuzaburo NODZU, Toshizo ISOSHIMA\*






(Nodzu Laboratory)

*Received June 25, 1954*

The reactions of ketene with phenol, resorcine, phloroglucine and dimedone have been studied.

- (1) Without catalyst, ketene gave always O-acetyl derivatives of them.
- (2) With sulfuric acid, pyridine or sodium acetate as catalyst, O-acetyl de-

Table 1. Reactions of ketene with phenol, resorcine, phloroglucine and dimedone.

Catalyst	<sup>(1)</sup> React. Temp.	Phenols Products	OH	OH	OH	
						
None	Low	Name <sup>(2)(3)</sup>	O	—	×	O*
		Yield <sup>(4)</sup>	80			55
	High	Name	—	O(mono), O(di)	O(tri)	O*
		Yield		67, 98	2	72
Na <sup>(5)</sup> Salt	Low	Name	O*	—	O(tri)	O*
		Yield	68		29	61
	High	Name	O*	O*(mono, di)	resine	O*
		Yield	74	78		61
H <sub>2</sub> SO <sub>4</sub>	Low	Name	O	—	resine	×
		Yield	85			
	High	Name	—	O(di)	resine	resine
		Yield		77		
	Low	Name	O	—	O(tri)	×
		Yield	87		14	
	High	Name	—	O(di)	O(tri)	O*
		Yield		88	26	61
AcONa	Low	Name	—	—	—	×
		Yield				
	High	Name	—	—	—	O*
		Yield				44

- (1) React. Temp.: Low—Cooling with ice or room temp. (solvent: ether).  
High—Warming on a steam bath (solvent: benzene).
- (2) O : O-acetyl derivative.  
O(mono) : O-mono-acetyl derivative.  
O(di) : O-di-acetyl derivative.  
O(tri) : O-tri-acetyl derivative.  
O\* : O-acetyl derivative with a small amount of C-acetyl derivative, which seems to be present although not confirmed.
- (3) × : No reaction.
- (4) Yield (%) : Theoretical yield.
- (5) Na-Salt : Treated with 20% H<sub>2</sub>SO<sub>4</sub> after the reaction.

\* 野津 竜三郎・磯 島 敏 三






NOTES

rivatives were produced.

(3) In the reactions with their sodium salts, O-acetyl derivatives were produced, containing a small amount of C-acetyl derivatives.

Some experimental results were cited in Tables 1 and 2.

Table 2. Some physical constants of O-acetyl derivatives obtained from phenol, resorcine, phloroglucine and dimedone.

O-acetyl derivative	Physical properties		
	m.p. (°C)	b.p. (°C)	$n_D^{20}$
	—	111 (60 mm.)	1.5200
	—	135-7 (7 mm.)	1.5328
	—	130-1 (7 mm.)	1.5034
	105-7	—	—
	—	128-132 (15 mm.)	1.4814

Reaction of Ketene with Ethyl Acetoacetate  
in the Presence of Pyridine

Toshizo ISOSHIMA\*  
(Nodzu Laboratory)

Received June 25, 1954

In the presence of a very small amount of pyridine, ketene was reacted with ethyl acetoacetate above  $-20^{\circ}\text{C}$ , and a reaction product rich in O-acetyl- (II), poor in C-acetyl ethyl acetoacetate (I), was obtained.

\* 磯 島 敏 三