

ABSTRACTS

**On the Transglycosidation relating to Riboflavin by  
*Escherichia coli*. (I)**

**Formation of Riboflaviny Glucoside**

Hideo KATAGIRI, Hideaki YAMADA and Kazutami IMAI

(Katagiri Laboratory)

*The Journal of Vitaminology*, 3, 264 (1957)

Riboflaviny glucoside has been isolated from reaction mixture composed of riboflavin, maltose and the cell suspension of *Escherichia coli*.

The suitable conditions and the effects of several substances on the formation of riboflaviny glucoside were studied, and it is ascertained that riboflavin plays the role of specific acceptor of transglycosidation, and concluded that the interaction of the enzyme on isoalloxazine ring of riboflavin molecule would take place in the first step of the reaction.

Thus the authors suggest that riboflavin reveals a new biochemical significance in carbohydrate metabolism.

---

**Microbiological Assay of Vitamins and Amino  
Acids by One-dimensional Diffusion Method**

Tomoji SUZUKI and Fumihiko TANAKA

(Suzuki Laboratory)

*Journal of the Pharmaceutical Society of Japan*

(*Yakugaku Zasshi*), 77, 1159 (1957)

We designed the one-dimensional diffusion method for microbiological assay of vitamins and amino acids. The agar synthetic medium containing triphenyl-tetrazoliumchloride (TTC) (50mg/100cc. of medium) and test organisms for example, *Lactobacillus arabinosus* 17-5 is poured into the test tube which is 75 mm in high and 5 mm in diameter. After the medium solidified, about 0.1 ml of sample were over it. Following incubation about 12 hours, red zone of reduced TTC were observed. This red zone is proportional to logarithms of concentration of amino acids and vitamins. Using *Lactobacillus arabinosus* 17-5, the assay range is as following : biotin 1~100 m $\gamma$ /cc., pantothenic acid and nicotinic acid 0.1~100 $\gamma$ /cc., DL-phenylalanine and DL-methionine 15~1000 $\gamma$ /cc., L-leucine 60~1000 $\gamma$ /cc. This method is distinguished with large range of determination and small amount of medium, still more convenient for routine assay