TITLE:

Inductive Effect of Substituents on the Symmetrical Methyl Deformation Frequencies of Aliphatic Hydrocarbons

AUTHOR(S):

Takenaka, Tohru

CITATION:

Takenaka, Tohru. Inductive Effect of Substituents on the Symmetrical Methyl Deformation Frequencies of Aliphatic Hydrocarbons. Bulletin of the Institute for Chemical Research, Kyoto University 1962, 39(6): 407-407

ISSUE DATE:

1962-03-25

URL:

<http://hdl.handle.net/2433/75871>

RIGHT:

ABSTRACTS

Inductive Effect of Substituents on the Symmetrical Methyl Deformation Frequencies of Aliphatic Hydrocarbons

Tohru TAKENAKA

*Nippon Kagaku Zasshi (Journal of the Chemical Society of
Japan, Pure Chemistry Section), 82, 1309 (1961)*

In order to discuss the inductive effect of polar group upon methyl group in a molecule, the symmetrical methyl deformation frequencies δ_{CH_3} of $\text{CH}_3(\text{CH}_2)_{n-1}\text{XH}_{m-1}$ molecules were measured, where X represents an atom in groups IV~VII of the periodic table, m the valence of X atom and n the carbon number of the molecule.

For $\text{CH}_3\text{XH}_{m-1}$ molecules ($n=1$), a linear relationship was found to exist between the frequency and logarithm of the electronegativity x_X of X atom for each row and column of the periodic table. These frequency variations are ascribed to change in the deformation force constants, and can be expressed as

$$\delta_{\text{CH}_3} = 375 \log \left(\frac{x_X}{r_{\text{CX}}^2} \right) + 1366$$

where r_{CX} is the C-X bond length. This equation holds not only for $\text{CH}_3\text{XH}_{m-1}$ molecules but also for the series $(\text{CH}_3)_2\text{XH}_{m-2}, \dots, (\text{CH}_3)_m\text{X}$ within the error of $\pm 1\%$.

Frequency variations due to the change of the carbon number of $\text{CH}_3(\text{CH}_2)_{n-1}\text{XH}_{m-1}$ molecule ($n \geq 2$) can be interpreted by a modified equation in which the factor $\{(x_{\text{C}} + 0.40\varepsilon_X\sigma^{n-2})/r_{\text{CC}}^2\}$ is used instead of (x_X/r_{CX}^2) in the above equation. Here, r_{CC} is the C-C bond length, x_{C} the electronegativity of carbon atom, ε_X the induced charge on the carbon atom adjacent to X atom, and σ the ratio of the induced charges at any two adjacent carbon atoms in the alkyl chain.

Dielectric Properties of Emulsions. (III)

Dielectric Behavior of W/O Emulsions

Tetsuya HANAI

Kolloid Zeitschrift, 177, 57 (1961)

Dielectric constants and electrical conductivities of W/O emulsions at rest and under shear were measured over a wide range of concentration and at frequencies ranging from 20 cps. to 5 mc.

Striking dielectric dispersions due to the interfacial polarization were observed at high frequency range above 100 kc., while the electrode polarization was found below 1 kc.

It was found that the dielectric dispersions due to the interfacial polarization