

The Papers Published by the Staff Members of the Institute from July 1985 to June 1986

Nuclear Chemistry

Relaxation of Quasiparticles Excited by a Strong Perturbation —Excitation by Pulse Laser—, H. Mazaki, *Ionizing Radiation*, **11**, 38 (1985).

Dimensional Crossover and Commensurability Effect in V/Ag Superconducting Multilayers, K. Kanoda, H. Mazaki, T. Yamada, N. Hosoi and T. Shinjo, *Phys. Rev. B*, **33**, 2052 (1986).

Measurements of Positron Annihilation Rate in Low-Temperature Gaseous Nitrogen, T. Kawaratani, Y. Nakayama and T. Mizogawa, *Bull. Inst. Chem. Res., Kyoto Univ.*, **63**, 177 (1985).

Operation of a Proportional Counter with He/CH₄ and Ar/CH₄ Mixtures at Low Temperatures, S. Kishimoto, Y. Isozumi, R. Katano and H. Takekoshi, *Nucl. Instr. & Methods*, **A249**, 349 (1986).

Vacancy Cascade Following Inner-Shell Ionization, T. Mukoyama, *Bull. Inst. Chem. Res., Kyoto Univ.*, **63**, 373 (1985).

Statistical Exchange Parameter and X-Ray Emission Rates, T. Mukoyama and H. Adachi, *Bull. Inst. Chem. Res., Kyoto Univ.*, **63**, 173 (1985).

Relativistic Calculations of the Excitation Cross Sections of Hydrogen-Like Ions by Heavy Charged-Particle Impact, T. Mukoyama, *Bull. Inst. Chem. Res., Kyoto Univ.*, **64**, 12 (1986).

Electron Shakeoff Accompanying Photoionization, T. Mukoyama, *Bull. Inst. Chem. Res., Kyoto Univ.*, **64**, 1 (1986).

Nuclear Excitation by Synchrotron Radiation and Its Applications, T. Nakajima, T. Mukoyama, H. Kaji, and H. Kobayakawa, *Prospects in Synchrotron Radiation Science in the X-Ray Region with the Next Generation Storage Ring*, ed. by S. Kikuta, O. Shimomura, and T. Matsushita, *KEK Report No. 85-16*, 106 (1986).

Calculations of the Energy Distribution of Electrons Ejected in Ion-Atom Collisions using Pseudostatse, T. Mukoyama, C.-D. Lin, and W. Fritsch, *Phys. Rev. A*, **32**, 2490 (1985).

Measurement of the Mass of the Electron Neutrino using Electron Capture in ¹⁶³Ho, S. Yasumi, F. Ochiai, M. Ando, H. Maezawa, H. Kitamura, M. Maruyama, M. Fujioka, K. Ishii, T. Shinozuka, K. Sera, T. Omori, G. Izawa, M. Yagi, K.

Masumoto, K. Shima, T. Mukoyama, Y. Inagaki, H. Taketani, I. Sugai, A. Masuda, and O. Kawakami, Neutrino Mass and Low Energy Weak Interactions, Telemark, 1984, ed. by V. Barger and D. Cline (World Scientific, Singapore, 1985) p. 398.

Calculations of Electron Energy Distribution Ejected in Ion-Atom Collisions by Psuedostate Method, T. Mukoyama, C.-D. Lin, and W. Fritsch, *Abst. Intern. Conf. On the Physics of Electronic and Atomic Collisions*, Stanford, Aug. 23–30, 1985 (North-Holland, Amsterdam, 1986) p. 411.

Measurement of the Mass of the Electron Neutrino using Electron Capture in ^{163}Ho , S. Yasumi, M. Ando, H. Maezawa, H. Kitamura, T. Ohta, F. Ochiai, A. Mikuni, M. Maruyama, M. Fujioka, K. Ishii, T. Shinozuka, K. Sera, T. Omori, G. Izawa, M. Yagi, K. Masumoto, K. Shima, T. Mukoyama, Y. Inagaki, I. Sugai, A. Masuda, and O. Kawakami, *KEK Preprint*, 85–45 (1985).

Breakup of ^4He with Polarized Protons at 65 MeV, K. Fukunaga, S. Kakigi, T. Ohsawa, A. Okihana, T. Sekioka, H. Nakamura-Yokota, S. Tanaka, *Nuclear Physics*, **A456**, 48 (1986).

Break Up of ^4He with Polarized Proton at 65 MeV, K. Fukunaga, S. Kakigi, T. Ohsawa, A. Okihana, T. Sekioka, H. Nakamura-Yokota, and S. Tanaka, *Proceedings of the Sixth International Symposium on Polarization Phenomena in Nuclear Physics*, Osaka 1985, 902 (1985).

Vector Polarization of ^6Li Elastically Scattered by ^{12}C at $E=150$ MeV, M. Tanaka, T. Yamagata, K. Yuasa, S. Nakayama, M. Inoue, Y. Sakuragi, M. Kamimura, H. Goto, K. Katori, M. Yanaga and H. Ogata, *Proceedings of the Sixth international Symposium on Polarization Phenomena in Nuclear Physics*, Osaka, 1985, 764 (1985).

Study of Permanent Magnet Quadrupole Lens for Proton Linac, H. Okamoto and Y. Iwashita, *Bull. Inst. Chem. Res., Kyoto Univ.*, **64**, 20 (1986).

Proton Inelastic Scattering to the 1^+ States in the M1-Resonance Energy Region in ^{58}Ni and ^{60}Ni , K. Hosono, M. Fujiwara, H. Ikegami, M. Kondo, N. Matsuoka, T. Saito, K. Ogino, S. Matsuki, and S. Kato, *J. Phys. Soc. Japan*, **55**, 606 (1986).

Hexadecapole Moments of the Medium-Weight Transitional Nuclei by 65 MeV Polarized Proton Inelastic Scattering, S. Matsuki, T. Ohsawa, H. Okamura, K. Ogino, K. Haga, N. Sakamoto, M. Matoba, and T. Komatuzaki, *J. Phys. Soc. Japan*, **55**, 608 (1986).

Excitation of M1 States in ^{58}Ni and ^{60}Ni by 65 MeV Polarized Proton Inelastic Scattering, K. Hosono, M. Fujiwara, H. Ikegami, M. Kondo, N. Matsuoka, T. Saito, T. Yamazaki, S. Matsuki, K. Ogino, S. Kato, and T. Yanagihara, *Nuclear Physics*, **A454**, 237 (1986).

The First Three 2^+ States of ^{34}S by the Inelastic Scattering of Polarized Protons at 65 MeV, N. Sakamoto, S. Matsuki, T. Ohsawa, H. Okamura, and T. Komatuzaki,

J. Phys. Soc. Japan, **55**, 604 (1986).

The (p , 3He) Reactions on p-f Shell Nuclei at 65 MeV, M. Matoba, H. Ijiri, H. Kametani, H. Ohgaki, S. Imabeppu, I. Kumabe, M. Hyakutake, N. Koori, T. Sakae, T. Maki, and S. Matsuki, *J. Phys. Soc. Japan*, **55**, 642 (1986).

Physical Chemistry

Improved Evaluation of Electrical Phase Parameters in a Dielectric Theory for Concentrated Suspensions of Spherical Particles, T. Hanai, K. Sekine and N. Koizumi, *Bull. Inst. Chem. Res., Kyoto Univ.*, **63**, 227 (1985).

Being Intimate with Books. —My Wanderings over the Fields in Natural Science—, T. Hanai, *Seibutubuturi (Biophysics)*, **25**, 133 (1985).

Theory of Dielectric Relaxations due to the Interfacial Polarization for Two-Component Suspensions of Spheres, T. Hanai and K. Sekine, *Colloid and Polymer Sci.*, **264**, 888 (1986).

What does It Mean by Microcapsules? —What Kind of Information can be Obtained by Measuring the Permittivity?—, T. Hanai, H. Zhang, K. Sekine and K. Asami, *Hyomen (Surface)*, **7**, 393 (1986).

Does “One-Sphere” System Conform to the Wagner Equation for Simple Binary Dispersions? An Experimental Assessment by Use of a Ball of Glass, Steel or Hydrated Gelatin Suspended in Aqueous Solutions, K. Asami and A. Irimajiri, *Bull. Inst. Chem. Res., Kyoto Univ.*, **63**, 259 (1985).

Dielectric Properties of Cultured Epithelial-Cell (MDCK)-Monolayer that Encompasses a Gelatin Droplet, K. Asami and A. Irimajiri, *J. Physiol. Soc. Jpn.*, **47**, 379 (1985).

Dielectric Dispersion of Cultured Epithelial Monolayers Formed on a Tubular Support, K. Asami and A. Irimajiri, *J. Physiol. Soc. Jpn.*, **48**, 170 (1985).

Effect of Cell Surface Irregularities on the ‘Apparent’ Membrane Capacity as Deduced from the Impedance of Spherical Cell Suspensions, A. Irimajiri, K. Asami, Y. Kinoshita and M. Ogawa, *J. Physiol. Soc. Jpn.*, **48**, 165 (1985).

Dielectric Cell for Deformable Solid, Y. Kita and N. Koizumi, *Bull. Inst. Chem. Res., Kyoto Univ.*, **63**, 249 (1985).

Magnetic and Dielectric Properties of $\text{Ba}_2\text{Ti}_4\text{Fe}_2\text{O}_{13}\text{-KT}_{2\text{i}6}\text{O}_{13}$, K. Iwauchi and Y. Ikeda, *Phys. Stat. Sol. (a)*, **92**, 623 (1985).

Dielectric and Magnetic Relaxation in $\text{BaO}\cdot2\text{TiO}_2\cdot2\text{Fe}_2\text{O}_3$ and $\text{BaO}\cdot2\text{SnO}_2\cdot2\text{Fe}_2\text{O}_3$, K. Iwauchi, Y. Ikeda, N. Koizumi and Y. Bando, *Bull. Inst. Chem. Res., Kyoto Univ.*, **63**, 241 (1985).

Dielectric Properties of Hexagonal Ferrites, K. Iwauchi and Y. Ikeda, *Phys. Stat. Sol. (a)*, **93**, 309 (1986).

Reentrant Spin Glass Behavior of $\text{BaTi}_2\text{Fe}_4\text{O}_{11}$, K. Iwauchi and Y. Ikeda, *J. Magn. Magn. Matr.*, **59**, 73 (1986).

Observation of Magnetic Clusters and Crystal Structures of $\text{BaTi}_{2-x}\text{Sn}_x\text{Fe}_4\text{O}_{11}$, M. Shiojiri, T. Maeda, S. Sekimoto, K. Iwauchi and Y. Ikeda, *Proc. 7th Inst. Summer School Detets in Crystals, Szczyrk, Poland*, (1985).

High-Resolution Electron Microscopy of $\text{Ba}_x\text{K}_{2-x}\text{Fe}_x\text{Ti}_{6-x}\text{O}_{13}$, M. Shiojiri, T. Maeda, Y. Hirota, T. Isshiki, S. Sekimoto, K. Iwauchi and Y. Ikeda, *J. Microscopy*, **142**, part II 223 (1986).

Pressure Dependence of the Ferroelectric Transition in Copolymers of Vinylidene Fluoride and Trifluoroethylene, N. Koizumi, Y. Murata and N. Haikawa, *Jpn. J. Appl. Phys.*, **24**, suppl. 24-2, 862 (1985).

Polarization Reversal and Double Hysteresis Loop in Copolymers of Vinylidene Fluoride and Trifluoroethylene, N. Koizumi, Y. Murata and H. Tsunashima, *IEEE Trans., EI-21*, 543 (1986).

Formation of Unoriented Form I Poly(vinylidene fluoride) by High-Rate Quenching and its Electrical Properties, Y. Oka and N. Koizumi, *Bull. Inst. Chem. Res., Kyoto Univ.*, **63**, 186 (1985).

Structure and Spontaneous Polarization in Fast-Quenched Copolymers of Vinylidene Fluoride and Trifluoroethylene, Y. Oka, Y. Murata and N. Koizumi, *Polym. J.*, **18**, 417 (1986).

An Application of Electron Energy Loss Spectroscopy to the Study of Residual Aluminum in Raney Nickel Catalyst, H. Kasano, H. Kurata, T. Kobayashi and N. Uyeda, *Bull. Inst. Chem. Res., Kyoto Univ.*, **63**, 216 (1985).

Direct Observation of Disorders of Molecular Stacking in Pt-Phthalocyanine Epitaxil Film, K. Yase, T. Kobayashi and N. Uyeda, *Bull. Inst. Chem. Res., Kyoto Univ.*, **63**, 383 (1985).

Structure and Growth Mechanism of $(\text{CrO}_3)_3\text{CH}_3\text{COOH}$ Intercalated Graphite, T. Kobayashi, H. Kurata and N. Uyeda, *J. Phys. Chem.*, **90**, 2231 (1986).

Comments on the Computation of Electron Wave-Propagation in the Slice Methods, K. Ishizuka, *J. of Microscopy*, **137**, 233 (1985).

Image Formation Theory of High-Resolution Electron Microscopy, K. Ishizuka, *J. Cryst. Soc. Japan*, **28**, 1 (1986), in Japanese.

Electron Microscopic Study of 5S rRNA Crystals from *thermus thermophilus* HB8, K. Morikaaw, Y. Fujiyoshi, K. Ishizuka, J. Sugiyama, M. Kawakami and S. Take-mura, *J. of Microscopy*, **142**, 247-258 (1986).

Surface and Interface Researches at the Institute for Chemical Research, Kyoto University, Y. Fujiyoshi, *J. of The Surface Science Society of Japan*, **6**, 376-377 (1985).

Adsorption of H₂O and Ar on Pulverized CaF₂, Y. Kuroda, T. Takenaka, J. Umemura, S. Kittaka, K. Morishige and T. Morimoto, *Langmuir*, **1**, 679 (1985).

FTIR-ATR Studies on Langmuir-Blodgett Films of Stearic Acid with 1-9 Monolayers, F. Kimura, J. Umemura and T. Takenaka, *Langmuir*, **2**, 96 (1986).

FT-IR Studies on Molecular Structure and Orientation of a Liquid Crystal, 4'-N-Octyl-4-Cyanobiphenyl, S. Hayashi, K. Kurita, N. Kimura, J. Umemura and T. Takenaka, *Bull. Inst. Chem. Res., Kyoto Univ.*, **63**, 276 (1985).

Orientation of Gramicidin D Incorporated into Phospholipid Multibilayers: a Fourier Transform Infrared-Attenuated Total Reflection Spectroscopic Study, E. Okamura, J. Umemura and T. Takenaka, *Biochim. Biophys. Acta*, **856**, 68 (1986).

A Fourier Transform Infrared Study of Bilayer Membranes of Double-Chain Ammonium Amphiphiles, N. Nakashima, N. Yamada, T. Kunitake, J. Umemura and T. Takenaka, *J. Phys. Chem.*, **90**, 3374 (1986).

Polarized UV Resonance Raman and Absorption Spectra of Monolayer Adsorbed at the Aqueous Solution-Carbon Tetrachloride Interface, T. Takenaka, N. Isono, J. Umemura, M. Shimomura and T. Kunitake, *Chem. Phys. Letters*, **128**, 551 (1986).

Structure of Dodecylpropyldimethylammonium Bromide Monohydrate, T. Taga, K. Machida, N. Kimura, S. Hayashi, J. Umemura and T. Takenaka, *Acta Cryst.*, **C42**, 2608 (1986).

Raman Spectral Studies of Dynamics of Ions in Molten LiNO₃-RbNO₃ Mixture. I. Rotational Relaxation, T. Kato and T. Takenaka, *J. Chem. Phys.*, **84**, 3405 (1986).

Infrared and Raman Spectra, T. Takenaka, *LB Films and Electronics*, Ed. by K. Fukuda et al., CMC, Tokyo, 1986, 72.

Studies on Interfacial Molecular Membranes by Infrared and Raman Spectroscopy, T. Takenaka, *Oyo Buturi*, **55**, 903 (1986).

FTIR Spectra of Langmuir-Blodgett Films, T. Takenaka, *Bunko Kenkyu*, **35**, 323 (1986).

Effect of the 295K Phase Transition in the Carbazole-Trinitrobenzene Complex on Vibrational Bands Measured by Fourier Transform Infrared Spectroscopy, A. Miniewicz, J. Sworakowski, D. F. Williams, D. G. Cameron and J. Memmura, *Spectrochim. Acta*, **41A**, 1305 (1985).

Structure Determinations of Thin Films by Vibrationla Spectroscopy, J. Umemura, *Yukagaku*, **34**, 767 (1985).

Tris(triphenylphosphine)rhodium (I) Chloride Catalyzed Synthesis of Methyl Formate from Methanol, Carbon Dioxide and Hydrogen in the Presence of a Tertiary Amine, H. Phala, K. Kudo, S. Mori and N. Sugita, *Bull. Inst. Chem. Res., Kyoto Univ.*, **63**, 63 (1985).

Kinetics and Mechanistic Study of the Methanol Homologation with Cobalt-Ruthenium Mixed Catalyst, K. Watanabe, K. Kudo and N. Sugita, *Bull. Chem. Soc. Jpn.*, **58**, 2029 (1985).

Double Bond Geometry of the Alkenes produced by Oxidative Elimination of Alkyl Phenyl Selenides and Tellurides, S. Umemura, Y. Hirai, K. Ohe and N. Sugita, *J. Chem. Soc., Chem. Commun.*, 1037 (1985).

Photoabsorption Cross Section of H/S, T. Ibuki, H. Koizumi, T. Yoshimi, M. Morita, S. Arai, K. Hironaka, K. Shinsaka, Y. Hatano, Y. Yagishita and K. Ito, *Chem. Phys. Letters*, **119**, 327 (1985).

Solubility of Carbon Monoxide in Hydrogen Fluoride and Hydrogen Fluoride-Boron Trifluoride, K. Kudo and N. Sugita, *Chemistry Express*, **1**, 65 (1986).

Carbonylation of Organometallic Compounds with Carbon Monoxide via Trans-metallation, S. Uemura, K. Ohe and N. Sugita, *Bull. Inst. Chem. Res., Kyoto Univ.*, **63**, 156 (1985).

Cobalt Carbonyl-Catalyzed Polycarbonylation of Aryl Halides in NaOMe/MeOH under Photostimulation, T. Kashimura, K. Kudo, S. Mori and N. Sugita, *Chem. Lett.*, 851 (1986).

Synthesis of a Novel Endogenous Opioid Peptide, Adrenorphin-Metorphamide using a New Deprotecting Procedure, Y. Kiso, T. Fujisaki, M. Shimokura, K. Okamoto and S. Uemura, *Peptide Chemistry 1984*: N. Izumiya (Ed.), 289 (1985).

Cobalt Carbonyl Catalyzed Polycarbonylation of Polyhalogenated Aromatics under Photostimulation, T. Kashimura, K. Kudo, S. Mori and N. Sugita, *Chem. Lett.*, 299 (1986).

Cobalt Carbonyl-Catalyzed Double-Carbonylation of α -Halogneated Benzoic Acids under Photostimulation, T. Kashimura, K. Kudo, S. Mori and N. Sugita, *Chem. Lett.*, 483 (1986).

Emission Spectra of $\text{SiH}(\text{A}^2\Delta \rightarrow \text{X}^2\Pi)$ and $\text{SiCl}_2(\tilde{\text{A}}^1\text{B}_1 \rightarrow \tilde{\text{X}}^1\text{A}_1)$ in the VUV Photolyses of Silane and Chlorinated Silanes, N. Washida, Y. Matsumi, T. Hayashi, T. Ibuki, A. Hiraya and K. Shobatake, *J. Chem. Phys.*, **83**, 2769 (1985).

Liquid-phase 1,4-Diacetoxylation of Conjugated Dienes with Tellurium (IV) Oxide and Alkali Metal Halides, S. Uemura and M. Okano, *Reports of the Asahi Glass Foundation for Industrial Technology*, **46**, 137 (1985).

Aromatic Thallation and Nitration with Thallium (III) Nitrate, S. Uemura and A. Toshimitsu, *Chemistry Express*, **1**, 99 (1986).

Analytical and Inorganic Chemistry

Anodic Stripping Voltammetry at a Nickel-Based Mercury Film Electrode, Z. Yoshida and S. Kihara, *Analytica Chimica Acta*, **172**, 39 (1985).

Determination of Rare Earth Elements in Phosphate Minerals by Inductively Coupled Plasma Atomic Emissoin Spectrometry, O. Fujino, K. Araki, Y. Kometani, M. Sugiyama, and M. Matsui, *Bunseki Kagaku (Analytical Chemistry)*, **34**, 386 (1985).

Charge Transfer at Liquid/liquid Interface and Oscillation of Interfacial Potential —Voltammetric Approach—, S. Kihara and M. Matsui, *Hyomen (Surface)*, **23**, 694 (1985).

Chemical Forms of Cobalt (0) Reated to Polarographic and Voltanmetric Catalytic Hydrogen Currents, S. Kihara, M. Matsui and Z. Yoshida, *J. Electroanal. Chem.*, **197**, 331 (1986).

Determination of Oxygen/Uranium Ratio in Irradiated Uranium Dioxide Based on Dissolution with Strong Phosphoric Acid, H. Takeishi, H. Muto, H. Aoyagi, T. Adachi, K. Izawa, Z. Yoshida, H. Kawamura, and S. Kihara, *Analytical Chemistry*, **58**, 458 (1986).

Preconcentration by Dithiocarbamate Extraction for Determination of Trace Elements in Natural Waters by Inductively —Coupled Plasma Atomic Emission Spectrometry, M. Sugiyama, O. Fujino, S. Kihara and M. Matui, *Analytica Chimica Acta*, **181**, 159 (1986).

Electrical Properties of Copper-Containing Aluminosilicate Glasses and Glass-Cermics, T. Yoko, K. Kamiya, Y. Kiyagi and S. Sakka, *Res. Rep. Fac. Eng., Mie Univ.*, **10**, 49 (1985).

Chemical and Physical Properties of Na-Si-O-N Oxynitride Glasses, S. Sakka, T. Kokubo and N. Sugimoto, *Collected Papers of XIV Intl. Congr. on Glass*, **II**, 218 (1986).

Glass-Formation and Optical Properties of Glasses in the systems (R_2O or $R'O$)- Ta_2O_5 - Ga_2O_3 , T. Kokubo, Y. Inaka and S. Sakka, *J. Non-Crystal. Solids*, **80**, 518 (1986).

Formation and Optical Properties of (R_2O or $R'O$)- Nb_2O_5 - Ga_2O_3 Glasses, T. Kokubo, Y. Inaka and S. Sakka, *J. Non-Crystal. Solids*, **81**, 337 (1986).

Photoconductivity of Bi_2O_3 -Containing Glasses, T. Kokubo and J. Ikeda, *Bull. Inst. Chem. Res., Kyoto Univ.*, **63**, 361 (1985).

Migration of Cu^+ ions in Cu_2O - Al_2O_3 - SiO_2 Glasses on Heating in Air, K. Kamiya, T. Yoko nad S. Sakka, *J. Non-Crystal. Solids*, **80**, 405 (1986).

Structural Change of Glass-Ceramics of the $\text{Cu}_2\text{O}-\text{Al}_2\text{O}_3-\text{SiO}_2$ System on Heating in Air, K. Kamiya, T. Yoko, K. Moroishi, K. Matsushita and S. Sakka, *J. Mater. Sci.*, **21**, 131 (1986).

X-ray Diffraction Study of $\text{Na}_2\text{O}-\text{GeO}_2$ Melts, K. Kamiya, T. Yoko, Y. Itoh and S. Sakka, *J. Non-Crystal. Solids*, **79**, 825 (1986).

Preparation of Porous TiO_2 Fibers by Unidirectional Freezing of Gel, T. Maki, Y. Teranishi, T. Kokubo and S. Sakka, *Yogyo-Kyokai-Shi (J. Ceram. Soc. Japan)*, **93**, 387 (1985), in Japanese.

Preparation of Porous Alumina Fibers by Unidirectional Freezing of Gel, T. Maki and S. Sakka, *J. Mater. Let.*, **5**, 28 (1986).

Hydrolysis and Polycondensation of Dimethyldiethoxysilane and Methyltriethoxysilane as Materials for the Sol-Gel Process, S. Sakka, Y. Tanaka and T. Kokubo, *J. Non-Crystal. Solids*, **82**, 24 (1986).

Formation of Alumina Fibers by Unidirectional Freezing of Gel, T. Maki and S. Sakka, *J. Non-Crystal. Solids*, **82**, 239 (1986).

Preparation of TiO_2 Film by the Sol-Gel Method and its Application to Photoelectrochemical Electrode, T. Yoko, K. Kamiya and S. Sakka, *Denki Kagaku (Electrochemistry)*, **54**, 284 (1986), in Japanese.

Properties of $\text{ZrO}_2-\text{SiO}_2$ Glasses Prepared by the Sol-Gel Method, K. Kamiya, T. Yoko, M. Iwanaka, A. Sakai and S. Sakka, *Zirconia Ceramics*, **5**, 39 (1986), in Japanese.

A New Glass-Ceramic for Bone Replacement: Evaluation of its Bonding to Bone Tissue, T. Nakamura, T. Yamamoto, S. Higashi, T. Kokubo and S. Ito, *J. Biomed. Mater. Res.*, **19**, 685 (1985).

Structure and Properties of a Load-Bearable Bioactive Glass-Ceramic, T. Kokubo, S. Ito, T. Hayashi, S. Sakka, T. Kitsugi, T. Yamamoto, M. Takagi and T. Shibuya, *Collected Papers of XIV Intl. Congr. on Glass*, **II**, 408 (1986).

SEM-EPMA Observation between Bone and Glass-Ceramics, T. Kitsugi, T. Yamamoto, T. Nakamura, S. Higashi, K. Kakutani, K. Hyakuna, T. Saito, M. Kitaoji, T. Hayashi, T. Kokubo, M. Takagi and T. Shibuya, *Chubunihon-Seikeigeka-Saigaigeka-Gakkaishi*, **28**, 2159 (1985), in Japanese.

Fatigue of Apatite- and Wollastonite-Containing Glass-Ceramic and Hydroxyapatite Ceramic in Vivo, T. Kitsugi, T. Yamamoto, T. Nakamura, S. Higashi, K. Kakutani, K. Hyakuna, T. Saito, T. Hayashi, S. Ito, T. Kokubo, S. Sakka, T. Shibuya, M. Takagi, K. Takeuchi and M. Ono, *Seikeigeka-Kisokagaku*, **13**, 648 (1986), in Japanese.

Formation of a High-Strength Bioactive Glass-Ceramic in the System MgO-CaO-SiO₂-P₂O₅, T. Kokubo, S. Ito, S. Sakka and T. Yamamuro, *J. Mater. Sci.*, **21**, 536, (1986).

Directionally Solidified Bi₂O₃-6Bi₂O₃·SiO₂ Eutectic, T. Kokubo and H. Kim, *J. Mater. Sci.*, **21**, 1441 (1986).

Medical Application of Ceramic Materials, S. Sakka, *Byotai-Seiri*, **4**, 553 (1985), Review in Japanese.

Glasses and Glass-ceramics from Gels, S. Sakka, *J. Non-Crystal. Solids*, **73**, 651 (1985), Review.

Functional Glasses and Glass-Ceramics, S. Sakka, *J. Soc. Mat. Sci. Japan*, **34**, 763 (1985), Review in Japanese.

Materials Design of Functional Glasses, S. Sakka, *Kinzoku (Metala)*, **55**(11), 34 (1985), Review in Japanese.

High Technology Glasses and Methods of their Materials Design, S. Sakka, *J. Korean Ceram. Soc.*, **22**(5), 85 (1985).

Sol-Gel Synthesis of Glasses: Present and Future, S. Sakka, *Am-Ceram. Soc. Bull.*, **64**, 1463 (1985).

Low Temperature Synthesis of Glass Fibers and Films Using Metal A;oxides. Formation of Gel Fibers and Films from Solution, S. Sakka and K. Kamiya, *Nihon Reoroji Gakkaishi (J. Soc. Rheol. Japan)*, **14**, 9 (1986), Review in Japanese.

What are NEW GLASSES?, S. Sakka, *Mol*, 65 (1986).

Glasses for Biomedical Applications, T. Kokubo, *Boundary*, **2**, 18 (1986), Review in Japanese.

Bioactive Artificial Bone, T. Kokubo, *Kobushi (Polymers)*, **35**, 247 (1986), Review in Japanese.

Magnetism of Ultrathin Fe₃O₄ Films, T. Terashima and Y. Bando, *J. Phys. Soc. Jpn.*, **54** (10), 3920 (1985).

Dielectric and Magnetic Relaxations in BaO·2TiO₂·2Fe₂O₃ and BaO·2SnO₂·2Fe₂O₃, K. Iwauchi, Y. Ikeda, N. Koizumi and Y. Bando, *Bull. Inst. Chem. Res., Kyoto Univ.*, **63**, 241 (1985).

High-Resolution Electron Microscopy of Ba_xK_{2-x}Fe_xTi_{6-x}O₁₃ Crystals in Polar-Glass State, M. Shiojiri, T. Maeda, Y. Hirota, T. Isshiki, S. Sekimoto, K. Iwauchi and Y. Ikeda, *J. Microscopy*, **142**, 223 (1985).

Magnetic and Dielectric Properties of Ba₂Ti₄Fe₂O₁₃-K₃Ti₆O₁₃, K. Iwauchi and Y. Ikeda, *Phys. Stat. Sol. (a)*, **92**, 623 (1985).

Preparation and Coercivity of Cobalt Ultrafine Particles by Reduction of Multi-layered CoO-SiO Films, Y. Bando and T. Terashima, *Bull. Chem. Soc. Jpn.*, **59**, 607 (1986).

Reentrant Spin Glass Behavior of BaTi₂Fe₄O₁₁, K. Iwauchi and Y. Ikeda, *J. Mag. Mag. Mat.*, **59**, 73 (1986).

Dielectric Properties of Hexagonal Ferrites, K. Iwauchi and Y. Ikeda, *Phys. Stat. Sol. (a)*, **93**, 309 (1986).

Formation of Hexagonal, Platelike Particles of BaFe_{12-2x}Ti(IV)_xCo(II)_xO₁₉ with 0≤x<1.2, M. Kiyama and T. Takada, *Adv. Cer.* (ed. Am. Cer. Soc.), **15**, 51 (1986).

Phase Relation in the Oxygen Nonstoichiometric System, SrDeO_x (2.5≤x≤3.0), T. Takeda, K. Kanno, T. Takada and O. Yamamoto, *J. Solid State Chem.*, **63**, 237 (1986).

Preparation and Characterization of Oxygen-Deficient Perovskite Oxides: SrFe_{1-x}V_xO_{2.5+x}, M. Takano, N. Nakayama and Y. Bando, *Adv. Cer.* (ed. Am. Cer. Soc.), **15**, 547 (1986).

Structures and Magnetic Properties of Artificial Superlattices Composed of NiO-CoO, NiO-Fe₃O₄, and CoO-Fe₃O₄, Y. Bando and T. Terashima, *Adv. Cer.* (ed. Am. Cer. Soc.), **15**, 607 (1986).

Study of the Mossbauer Effect on the Surface of BaO·6Fe₂O₃, H. Yamada, M. Takano, M. Kiyama, T. Takada and T. Shinjo, *Adv. Cer.* (ed. Am. Cer. Soc.), **16**, 169 (1986).

Formation of γ -(Fe_{1-x}Cr_x)₂O₃ (x=0.43) by Wet Method, K. Watanabe, T. Shigematsu, M. Kiyama, T. Shinjo and T. Takada, *Adv. Cer.* (ed. Am. Cer. Soc.), **16**, 331 (1986).

Ceramics with Superlattice Structures (in Japanese), Y. Bando, *Electronic Ceramics* **17**, 62 (1986).

Ultrathin Metal Film, T. Shinjo, *Kagaku to Kogyo (Chemistry and Industry)*, **38**, 749-751 (1985), in Japanese.

Structural Aspects of Fe-Mg Artificial Superstructure Films Studied by X-Ray Diffraction, Y. Fujii, T. Ohnishi, T. Ishihara, Y. Yamada, K. Kawaguchi, N. Nakayama, and T. Shinjo, *J. Phys. Soc. Jpn.*, **55**, 251-262 (1986).

Magnetic Properties of Artificial Metallic Superlattices, T. Shinjo, N. Hosoi, K. Kawaguchi, N. Nakayama, T. Takada and Y. Endoh, *J. Magn. & Magn. Mater.*, **54-57**, 737-742 (1986).

Metallic Superlattices, T. Shinjo, *Hyperfine Interactions*, **27**, 193-202 (1986).

NMR investigations for New-Type Alloys Formation at the interface in Artificial Superstructure Film, K. Takanashi, H. Yasuoka and T. Shinjo, *J. Magn. & Magn. Mater.*, **54-57**, 783-784 (1986).

A Structural Study of Co-Sb Multilayered Film by X-Ray Diffraction, N. Nakayama, K. Takahashi, T. Shinjo, T. Takada and H. Ichinose, *Jpn. J. Appl. Phys.*, **25**, 552-557 (1986).

Structure and Magnetic Properties of Mn-Sb Multilayered Films with Artificial Superstructure, N. Nakayama, T. Katamoto, T. Shinjo and T. Takada, *J. Magn. Soc. Jpn.*, **10**, 283-286 (1986).

Observation of Optical Activity of Some Biologically Related Substances, N. Baba, *Kagaku Kyoiku*, **33**, 70-71 (1985).

A Mechanistic Aspect of the Remote Asymmetric Transmission Observed during Substrate Reduction with Chiral NADH Model Compounds, N. Baba, F. Hoshide, J. Oda and Y. Inouye, *Israel Journal of Chemistry*, **25**, 236-240 (1985).

Facile Synthesis of Morpholines and Azacrown Ethers by Ozonolysis of Cyclic Olefins and Reductive *N*-Alkylation, M. Kawaguchi, J. Ohashi, Y. Kawakami, Y. Yamamoto and J. Oda, *Synthesis*, 701-703 (1985).

Optical Resolution and Asymmetric Synthesis of Optically Active Compounds by Enzymatic and Non-Enzymatic Methods, Y. Yamamoto and J. Oda, *Kagaku to Seibutsu*, **23**, 446-454 (1985), Review in Japanese.

Kinetic Treatment of Kinetic Resolution of Racemic Acid Anhydrides with Chiral Amines, Y. Kawakami, J. Hiratake, Y. Yamamoto, and J. Oda, *Bull. Inst. Chem. Res., Kyoto Univ.*, **63**, 100-103 (1985).

Catalytic Asymmetric Induction from Prochiral Cyclic Acid Anhydrides Using Cinchona Alkaloids, J. Hiratake, Y. Yamamoto, and J. Oda, *J. Chem. Soc. Chem. Commun.*, 1717-1719 (1985).

NADH Model Reaction: Asymmetric Reduction of Non-Activated Carbonyl Compounds with *N*-Anionized Hantzsch Ester, M. Watanabe, M. Fushimi, N. Baba, J. Oda and Y. Inouye, *Agric. Biol. Chem.*, **49**, 3533-3538 (1985).

Quantitative Structure-Reactivity Analysis of the Inclusion Mechanism by Cyclodextrins, Y. Matsui, T. Nishioka and T. Fujita, *Topics in current chemistry*, **128**, 61-89 (1985).

An Asymmetric Synthesis of Lactones from Cyclic Acid Anhydrides with Chiral Binaphthylidiamines, Y. Kawakami, J. Hiratake, Y. Yamamoto and J. Oda, *Agric. Biol. Chem.*, **50**, 693-698 (1986).

Quantitative Analysis of Effects of Substituted Phenols on Membrane Characteristics of Lecithin Liposomes, H. Miyoshi, T. Nishioka and T. Fujita, *Bull. Chem. Soc. Japan.*, **59**, 1099-1107 (1986).

Organic Chemistry

Evaluation of the Affinity of New N Atom-Containing Tumor Inhibitors for Nucleic Acids, Y. Nagao, S. Sano, T. Miyasaka, M. Ochiai, K. Fuji, E. Fujita, and H. Ishii, *Nucleic Acid Res. (Symposium Series No. 16)*, 37 (1985).

Hard Acid and Soft Nucleophile Systems. 10. The Mechanistic Study for Debromination of *para*-Bromoanisole with Aluminum Chloride and Ethanedithiol, M. Node, T. Kawabata, E. Fujita, and K. Fuji, *Bull. Inst. Chem. Res. Kyoto Univ.*, **63**, 47–51 (1985).

Design and Study of Tumor Inhibitors, Y. Nagao and M. Ochiai, *Bull. Inst. Chem. Res., Kyoto Univ.*, **63**, 132 (1985).

Dehydrosilylation of Alkenylsilanes Utilizing Polyvalent Organoiodine Compounds, M. Ochiai, K. Sumi, Y. Nagao, E. Fujita, M. Arimoto and H. Yamaguchi, *J. Chem. Soc., Chem. Commun.*, 697 (1985).

Investigation of New Chiral 1,3-Oxazolidine-2-thiones: Analytical Separation and Optical Resolution of Racemic Carboxylic Acids and Aminoacids, Y. Nagao, T. Kumagai, S. Yamada, E. Fujita, Y. Inoue, Y. Nagase, S. Aoyagi and T. Abe, *J. Chem. Soc., Perkin Trans. I*, 2361 (1985).

Use of Chiral 1,3-Oxazolidine-2-thiones in the Diastereo-selective Synthesis of Aldols, Y. Nagao, S. Yamada, T. Kumagai, M. Ochiai, and E. Fujita, *J. Chem. Soc., Chem. Commun.*, 1418 (1985).

A Facile Chiral Synthesis of (+)-Prelog-Djerassi Lactonic Acid Methyl Ester Using Five-membered Heterocyclic Chiral Reagents, Y. Nagao, T. Inoue, K. Hashimoto, Y. Hagiwara, M. Ochiai, and E. Fujita, *J. Chem. Soc., Chem. Commun.*, 1419 (1985).

Reaction of Alkynyltrimethylsilanes with a Hypervalent Organoiodine Compound: A New General Synthesis of Alkynylodonium Salts, M. Ochiai, M. Kunishima, K. Sumi, Y. Nagao, E. Fujita, M. Arimoto, and H. Yamaguchi, *Tetrahedron Lett.*, **26**, 4501 (1985).

Highly Selective Nonenzymatic Chiral Induction into 3-Methylglutaric Acid and *cis*-4-Cyclohexen-1,2-ylenebis (acetic acid) Utilizing a Functional Five-Membered Heterocycle 4(*R*)-MCTT, Y. Nagao, T. Ikeda, M. Yagi, M. Shiro, and E. Fujita, *J. Org. Chem.*, **50**, 4072 (1985).

Steroidogenic Effect of *Ent*-Kaur-16-en-15 β -ol(Kaurenol) on Isolated Rat Adrenal Cells, K. Moriwaki, M. Gomi, Y. Itoh, S. Iida, M. Tsugawa, S. Tarui, K. Fuji, M. Node, and T. Kajimoto, *Life Sciences*, **38**, 453–458 (1986).

New C4-Chiral 1,3-Thiazolidine-2-thiones: Excellent Chiral Auxiliaries for Highly Diasterecontrolled Aldol-Type Reactions of Acetic Acid and α , β -Unsatu-

rated Aldehydes, Y. Nagao, Y. Hagiwara, T. Kumagai, M. Ochiai, T. Inoue, K. Hashimoto, and E. Fujita, *J. Org. Chem.*, **51**, 2391 (1986).

Highly Selective Asymmetric Induction onto Prochiral σ -Symmetric Molecules, Y. Nagao and E. Fujita, "Fine Synthetic Chemistry" *Kagaku Zokan*-108, Eds. by H. Nozaki and K. Oshima; Kagakudojin, Kyoto, 189 (1986).

New Synthetic Reaction Utilizing Organosilicon and Tin Compounds, M. Ochiai, *Advances in Pharmaceutical Sciences*, **2**, 185 (1986).

Enantioselective Reactions, K. Fuji, *J. Synth. Org. Chem. Jpn.*, **44**, 623-632 (1986).

Hypervalent Organoiodine Compounds in Organic Syntheses; Reaction with Organosilicon and Tin, M. Ochiai and Y. Nagao, *J. Synth. Org. Chem. Japan*, **44**, 660 (1986).

The reaction of 2,2-Dimethyl-4-lithio-1,3-oxathiane 3,3-Dioxide. General Synthesis of γ -Hydroxyketones, K. Fuji, M. Node, and Y. Usami, *Chem. Lett.*, 961 (1986).

Asymmetric Induction via Addition-Elimination Process: Nitroolefination of α -Substituted Lactones, K. Fuji, M. Node, H. Nagasawa, Y. Naniwa, and S. Terada, *J. Am. Chem. Soc.*, **108**, 3855-3856 (1986).

NAD(P) $^+$ -NAD(P)H Model. 58. Desulfonylation of β -Keto Sulfones with the Hantzsch Ester, an NAD(P)H Model., K. Nakamura, M. Fujii, H. Mekata, S. Oka and A. Ohno, *Chemistry Lett.*, 87 (1986).

NAD(P) $^+$ -NAD(P)H Models. 57. Stereochemistry in (net) Hydride Transfer from and to NAD(P) $^+$ -NAD(P)H Models: Chirality Shink., A. Ohno, M. Kashiwagi, Y. Ishihara, S. Ushida, and S. Oka, *Tetrahedron*, **42**, 961 (1986).

Bacterial Degradation of Biotin and Dethiobiotin: Isolation and Identification of β -Hydroxy and Keto Metabolites., M. Osakai, Y. Izumi, K. Nakamura and H. Yamada, *Agric. Biol. Chem.*, **50**, 311 (1986).

Stereochemical Control in Microbial Reduction. 3. Diastereo- and Enantioselective Reduction of 2-Methyl-3-oxobutanoate by Bakers' Yeast., K. Nakamura, K. Miyai, K. Nozaki, K. Ushio, S. Oka and A. Ohno, *Tetrahedron Lett.*, **27**, 3155 (1986).

NAD(P) $^+$ -NAD(P)H Models. 55. Transition Metal Catalyzed Reduction of Organic Halides: High Selectivity for Reductive Dehalogenation., S. Yasui, K. Nakamura, M. Fujii and A. Ohno, *J. Org. Chem.*, **50**, 3283 (1985).

Stereochemical Control of Microbial Reduction. 2. Reduction of β -Keto Esters by Immobilized Bakers' Yeast., K. Nakamura, M. Higaki, K. Ushio, S. Oka and A. Ohno, *Tetrahedron Lett.*, **26**, 4213 (1985).

Stereochemical Control in Microbial Reduction 4. Effect of Cultivation Conditions on the Reduction of β -Keto Esters by Methylotrophic Yeasts., K. Ushio, K. Inoue, K. Nakamura, S. Oka, and A. Ohno, *Tetrahedron Lett.*, **27**, 2657 (1986).

Simulation of NAD(P)⁺-NAD(P)H Redox System., A. Ohno and S. Ushida, *Bull. Inst. Chem. Res., Kyoto Univ.*, **63**, 104 (1985).

Model Studies with Nicotinamide Derivatives., S. Yasui and A. Ohno, *Bioorg. Chem.*, **14**, 70 (1986).

Incorporation of Dioxygen into the Hydroxylated Product during the C-C Single Bond Cleavage of 1,2-Bis(p-methoxyphenyl)-propane-1,3-diol Catalyzed by Hemin. A Nobel Model System for the Hemoprotein Ligninase, T. Habe, M. Shimada, T. Okamoto, B. Panijpan, and T. Higuchi, *J. Chem. Soc., Chemical Communication*, 1323, (1985).

Phenylselenenyl Chloride in Acetonitrile-Water: A Highly Convenient Reagent for Hydroxyselenation of Olefins and Preparation of Cyclic Ethers from Dienes, A. Toshimitsu, T. Aoai, H. Owada, S. Uemura, and M. Okano, *Tetrahedron*, **41**, 5301 (1985).

Design of New Organoselenium Reagents and Their Utilization in Organic Synthesis., A. Toshimitsu, *Yuki Gosei Kagaku Kyoka Shi (J. of Synth. Org. Chem., Jpn)*, **43**, 921 (1985). Review in Japanese.

Amidoselenation of Olefins via Hydroxyselenation: Reactions using Nitriles in Reagent Quantity and the Synthesis of β -(Acrylamido)alkyl Phenyl Selenides, A. Toshimitsu, G. Hayashi, K. Terao, and S. Uemura, *J. Chem. Soc., Perkin 1*, 343 (1986).

Intramolecular Amidoselenation of Olefinic Imidates to Afford Lactams, A. Toshimitsu, K. Terao, and S. Uemura, *J. Chem. Soc., Chem. Commun.*, 530 (1986).

Intramolecular Amidoselenation of N-Alkenylamides: Formation of Nitrogen Heterocycles, A. Toshimitsu, K. Terao, and S. Uemura, *J. Org. Chem.*, **51**, 1724 (1986).

Polymer Chemistry

Dilute Solution Properties of Imogolite, N. Donkai, H. Inagaki, K. Kajiwara, H. Urakawa and M. Schmidt, *Makromol. Chem.*, **186**, 2623 (1985).

Some Remarks on Imogolite Mesophase, K. Kajiwara, N. Donkai, Y. Fujiyoshi, Y. Hiraki, H. Urakawa and H. Inagaki, *Bull. Inst. Chem. Res., Kyoto Univ.*, **63**, 320 (1985).

Fundamental Study on Thin Layer Chromatography I. Solvent Strength of Mixture for Polymeric Samples, N. Donkai and H. Inagaki, *Bull. Inst. Chem. Res., Kyoto Univ.*, **63**, 313 (1985).

On the Excluded Volume Perturbation, H. Suzuki, *Macromolecules*, **18**, 2082 (1985).

Heat Capacities of Liquid Linear Alipahtic Polyoxides, H. Suzuki and B. Wunderlich, *J. Polym. Sci.: Polym. Phys. Ed.*, **23**, 1671 (1985).

Artificial Crimping of Wool Fibres by Draft/Relaxation Process —Crimp Stabilization by Chemical Modification—, R. Umehara, Y. Shibata, H. Ito, T. Miyamoto and H. Inagaki, *Proc. 7th Int. Wool Textile Res. Conf.*, **4**, 402 (1985).

An Aspect for the Mechanism of Bilateral Staining of Merino Wool Fibers by Basic Dyes, T. Miyamoto, H. Sakabe, H. Ito and H. Inagaki, *Proc. 7th Wool Textile Res. Conf.*, **1**, 125 (1985).

Recent Developments in Vascular Prostheses, T. Miyamoto and Y. Noishiki, *Bio Industry*, **2**, 884 (1985).

Isolation and Characterization of Orthocortical and Paracortical Cells from Merino Wool Fibers, H. Ito, H. Sakabe, T. Miyamoto and H. Inagaki, *Proc. 7th Int. Wool Textile Res. Conf.*, **1**, 115 (1985).

Excluded Volume in An Isolated Chain, H. Suzuki, *Kaigai Kobunshi Kenkyu*, **31**, 205 (1985), Review in Japanese.

A New Method to Heparinize the Biological Materials, Y. Noishiki, T. Miyata, H. Ito and T. Miyamoto, *Jpn. J. Artif. Organs*, **15**, 154 (1986), Monograph in Japanese

Preparation of Cellulose Graft Copolymers Having Polypeptide Side Chains and Their Blood Compatibility, T. Miyamoto, S. Takahashi, S. Tsuji, H. Ito and H. Inagaki, *J. Appl. Polym. Sci.*, **31**, 2303 (1986).

Cellulose as Materials of Functional Polymers, T. Miyamoto, *Sen-i Gakkaishi (J. Soc. Fiber Sci. Tech., Japan)*, **42**, 25 (1986).

Light Scattering from Polymer Blends in Solution. 2. Nondilute Solutions of Polystyrene and Poly(methyl methacrylate), T. Fukuda, M. Nagata and H. Inagaki, *Macromolecules*, **19**, 1411 (1986).

Light Scattering from Polymer Blends in Solution. 3. Accurate Determination of Polymer-Polymer Interactions by Using Mixed Solvents, M. Nagata, T. Fukuda and H. Inagaki, *Makromol. Chem., Rapid Commun.*, **7**, 127 (1986).

Interpretation for Velocities of Free-Radical Copolymerization, T. Fukuda, *Kaigai Kobunshi Kenkyu*, **31**, 120 (1985), Review in Japanese.

Measurements of Molecular Weights and Molecular Weight Distributions, "Kobunshi-Bunseki Handbook", T. Fukuda, N. Donkai and S. Teramachi, *Society of Analytical Chemistry of Japan, ed., Asakura, Tokyo*, Chapter 4.

Formation of Polyelectrolyte Complexes between Cellulose Derivatives and Their Blood Compatibility, H. Ito, T. Shibata and T. Miyamoto, *J. Appl. Polym. Sci.*, **31**, 2491 (1986).

Dynamics of Polymer-Polymer-Solvent Ternary Systems. 2. Diffusion and Sedimentation of Poly(methyl methacrylate) in Semidilute Solutions of Polystyrene in Thiophenol, N. Nemoto, T. Inoue, Y. Makita, Y. Tsunashima, and M. Kurata, *Macromolecules*, **18**, 2516 (1985).

The Particle Scattering Function and the Distribution Functions in the Interior of Fully Swollen Polymer Molecules, Y. Tsunashima and M. Kurata, *J. Chem. Phys.* **84**, 6432 (1986).

Permeation of Gases in Poly(2-vinylpyridine)-Block-Polyisoprene, H. Odani, M. Uchikura, Y. Ogino, and M. Kurata, *Bull. Inst. Chem. Res., Kyoto Univ.*, **63**, 332 (1985).

Stress Relaxation of a Branched Polybutadiene in Double-Step Shear Deformations, K. Osaki, S. Kimura, and M. Kurata, *J. Rheology*, **29**, 533 (1985).

Rheology of Polymers with Various Shapes, K. Osaki, *Kaigai Kobunshi Kenkyu*, **31**, 33 (1985), Review in Japanese.

Recent Developments in Fundamental Membranology, H. Odani, *Kaigai Kobunshi Kenkyu*, **31**, 178 (1985).

Structural Formation During High-Speed Melt-Spinning, M. Tsuji, S. Murakami and K. Katayama, *Proc. USF-85, Hakone*, 223, (1985).

Fiber Structure of Poly(p-phenylene), A. Kawaguchi and K. Katayama, *Proc. ISF-85, Hakone*, 130, (1985).

Some Aspects of Doping Mechanism in Polyschweifelnitrid (SN)_x, A. Kawaguchi, S. Isoda, J. Petermann and K. Katayama, *Colloid Polym. Sci.*, **263**, 631, (1985).

Epitaxial Crystallization of Polyethylene and Paraffin on Oriented Polypropylene, G. Broza, U. Rieck, A. Kawaguchi and J. Petermann, *J. Polym. Sci.; Polym. Phys. Ed.*, **23**, 2623, (1985).

Paracrystalline Nature of Poly(p-phenylene), A. Kawaguchi and J. Petermann, *Mol. Cryst. Liq. Cryst.*, **133**, 189 (1986).

Image Analysis in the Electron Microscopy of Cellulose Protofibrils. II. Digital Correlation Method, M. Tsuji, J. Frank and R. St. J. Manley, *Colloid Polym. Sci.*, **264**, 89 (1986).

One-dimensional Image Simulation in the Electron Microscopy of Negatively Stained Cellulose Protofibrils, M. Tsuji and R. St. J. Manley, *Sen-i Gakkaishi*, **42**, T-323, (1986).

Polymer Crystallization in Melt Spinning: Mathematical Simulation, K. Katayama and M.-G. Yoon, "High-Speed Fiber Spinning", Edited by A. Ziabicki and H. Kawai, John Wiley & Sons, New York, 1985.

High Resolution Electron Microscopy of Fiber Structure, K. Katayama, *Proc. ISF-85, Hakone*, 24, (1985).

Solid-State High-Resolution NMR, R. Kitamaru and F. Horii, "Sen-i Kobunshi Sokuteiho no Gijutsu", the Society of Fiber Science and Technology, Japan, Ed., Asakura 1985, 210, in Japanese.

Carbon-13 Nuclear Spin Relaxation Study as an Aid to Analysis of Chain Dynamics and Conformation of Macromolecules, R. Kitamaru, "Applications of NMR Spectroscopy to Problems in Stereochemistry and Conformational Analysis", Y. Takeuchi and P. Marchand, Eds., Verlag Chemic International, Deerfield Beach, FL., 1986, 75.

Structure of Insoluble and Infusible Materials, F. Horii, *Koubunshi (High Polymers, Japan)*, 34, 550 (1985), in Japanese.

Cross-Polarization/Magic-Angle Spinning ^{13}C NMR Studies of Cotton and Cupra Rayon with Different Water, F. Horii, A. Hirai, R. Kitamaru and I. Sakurada, *Cellulose Chem. Technol.*, 19, 513 (1985).

Solid Structure Analyses of Cellulose-An Approach by CP/MAS ^{13}C NMR, F. Horii, A. Hirai and R. Kitamaru, *Nippon Kagaku Sen-i Kenkyusho Koenshu (Annual Report of the Res. Inst. for Chem. Fibers, Japan)*, 42, 41 (1985), in Japanese.

Solid-State High-Resolution Carbon-13 NMR Studies of Celluloses, A. Hirai, F. Horii and R. Kitamaru, *Bull. Inst. Chem. Res., Kyoto Univ.*, 63, 340 (1985).

Phase Structure of Lamellar Crystalline Polyethylene by Solid-State High-Resolution ^{13}C NMR. Detection of the Crystalline-Amorphous Interphase, R. Kitamaru, F. Horii and K. Murayama, *Macromolecules*, 19, 636 (1986).

CP/MAS ^{13}C NMR Spectroscopy of Hydrated Amyloses Using a Magic-Angle Spinning Rotor with an O-ring Seal, F. Horii, A. Hirai and R. Kitamaru, *Macromolecules*, 19, 930 (1986).

Neutron Inelastic Scattering Spectra of Native and Regenerated Celluloses, K. Kaji, T. Kanaya, K. Inoue, R. Kitamaru and I. Sakurada, *Cellulose Chem. Technol.*, 19, 321 (1985).

Effects of Added Salts on the Chain Conformation of Polyelectrolytes, K. Kaji, T. Kanaya, I. Ishida and H. Yamaoka, *KURRI-TR*, 280, 42 (1985), in Japanese.

Molecular weight-Concentration Phase Diagram of Polyelectrolyte Solution, K. Kaji, *Report of the Meeting on Kinetics of Ordering Process in Phase Transition*, 79 (1986), in Japanese.

Crystal Analyser Type Spectrometer Using Pulsed Neutrons and Polymer Spectroscopy, T. Kanaya, *Report of the First Meeting, KUR Sem-mon Kenkyukai*, **60**, 100 (1986), in Japanese.

Determination of Long Period in Fluorocarbon Polymers by Small-Angle Neutron Scattering, R. Kitamaru, T. Kanaya, K. Nishida, M. Ono and F. Yoshida, *KURRI-TR*, **280**, 39 (1985), in Japanese.

Application of a Wide Band Chopper for Neutron Small Angle Scattering from Polyelectrolyte Solutions without Added Salts, M. Ono, S. Okamoto, T. Kanaya, K. Nishida, H. Urakawa, K. Kaji and R. Kitamaru, *Physica*, **138B**, 49 (1986).

Neutron Inelastic Scattering Spectra of Native and Regenerated Celluloses, K. Kaji, T. Kanaya, K. Inoue, R. Kitamaru and I. Sakurada, *Cellulose Chem. Technol.*, **19**, 321 (1985).

Effects of Added Salts on the Chain Conformation of Polyelectrolytes, K. Kaji, T. Kanaya, I. Ishida and H. Yamaoka, *KURRI-TR*, **280**, 42 (1985), in Japanese.

Application of Neutron Scattering to Chemistry, K. Kaji, *Symposium on Scientific Boundary Research by High Intensity Proton Accerelator*, 100 (1986), in Japanese.

Neutron Scattering and Collective Oscillation under Thermal Fluctuation (I), T. Kanaya, *Bull. Inst. Chem. Res., Kyoto Univ.*, **64**, 49 (1986).

Neutron Scattering, K. Kaji, *Sen-i Gakkaishi (J. Soc. Fiber Science and Technology)*, **41**, 468 (1985), Review in Japanese.

Static and Dynamic Structure of Polymers Revealed by Neutron Scattering, K. Kaji, *Koubunshi (High Polymers, Japan)*, **35**, 240 (1986), Review in Japanese.

Biochemistry

Leucine Dehydrogenase, T. Ohshima and K. Soda, *Hakko to Kogyo*, **43**, 919, (1985), Review in Japanese.

Mechanism of Inactivation of α -Amino- ϵ -caprolactam Racemase by α -Amino- δ -valerolactam, S.A. Ahmed, N. Esaki, H. Tanaka and K. Soda, *Agric. Biol. Chem.*, **49**, 299 (1985).

Actions of *S*-(*N*-Methylthiocarbamoyl)-L-cysteine and Iys Oxygen Analog on Pyridoxal Enzymes, T. Kimura, N. Esaki, H. Tanaka, and K. Soda, *Bull. Inst. Chem. Res., Kyoto Univ.*, **63**, 95 (1985).

Enzyme Chemistry of Mammalian Selenocysteine Metabolism, H. Tanaka, N. Esaki, T. Nakamura, N. Karai and K. Soda, *Biryo-Eiyoso Kenkyu*, **2**, 35 (1985) in Japanese.

A Versatile Bacterial Enzyme: L-Methionine γ -Lyase, H. Tanaka, N. Esaki, and K. Soda, *Enzyme Microb. Technol.*, **7**, 530 (1985).

Selenocysteine Metabolism in Mammals, H. Tanaka, N. Esaki, and K. Soda, *Current Topics in Cellular Regulation*, **27**, 487 (1985).

Improvement for L-Leucine Production in a Continuously Operated Enzyme Membrane Reactor, T. Ohshima, C. Wandrey, M.-R. Kula, and K. Soda, *Bio-tech. Bioeng.*, **27**, 1616 (1985).

Screening of Thermostable Leucine and Alanine Dehydrogenases in Thermophilic *Bacillus* Strains, T. Ohshima, C. Wandrey, M. Sugiura, and K. Soda, *Bio-tech. Lett.*, **7**, 871 (1985).

Purification and Characterization of Selenocysteine β -Lyase from *Citrobacter freundii*, P. Chocat, N. Esaki, K. Tanizawa, K. Nakamura, H. Tanaka, and K. Soda, *J. Bacteriol.*, **163**, 669 (1985).

Catalytic Action on Sulfur Analog of α -Amino- ϵ -caprolactam, H. Tanaka, S.A. Ahmed, N. Esaki and K. Soda, *Ganryu-Aminosan (Sulfur Amino Acids)*, **8**, 49 (1985).

Catalytic Action and Mechanism of Selenocysteine β -Lyase, N. Esaki, T. Nakamura, P. Chocat, N. Karai, H. Tanaka and K. Soda, *Ganryu-Aminosan (Sulfur Amino Acids)* **8**, 63 (1985).

Glutamate-Aspartate Transaminase from Microorganisms, T. Yagi, H. Kagamiyama, M. Nozaki, and K. Soda, *Methods Enzymol.*, **113**, 83 (1985).

L-Kynurenine Transaminase from *Hansenula schneggii*, K. Tanizawa, Y. Asada, and K. Soda, *Methods Enzymol.*, **114**, 90 (1985).

L-Lysine Transaminase from *Flavobacterium lutescens*, K. Tanizawa, T. Yoshimura, and K. Soda, *Methods Enzymol.*, **113**, 96 (1985).

Inactivation of Ω -Amino Acid: Pyruvate Transaminase and Taurine Transaminase with Se-Hypotaurine, K. Yonaha, S. Toyama, T. Sadeh, H. Tanaka and K. Soda, *Ganryu-Aminosan (Sulfur Amino Acids)*, **8**, 67 (1985).

Sites of *dnaA* Protein-binding in the Replication Origin of the *Escherichia coli*, M. Matsui, A. Oka, M. Takanami, S. Yasuda and Y. Hirota, *J. of Molecular Biology*, **184**, 529-533 (1985).

Transcription and Translation of the Kanamycin Resistance Transposon Tn903, M. Matsui and A. Oka, *Bull. Inst. Chem. Res., Kyoto Univ.*, **63**, 52-62 (1985).

General Recombination Mechanisms in Extracts of Meiotic Cells, Y. Hotta, S. Tabata, R.A. Rouchard, R. Pionon and H. Stern, *Chromosoma (Berlin)*, **93**, 140-146 (1985).

Meiosis-specific Transcripts of a DNA Component Replicated during Chromosome Pairing: Homology Across the Phylogenetic Spectrum, Y. Hotta, S. Tabata, L. Stubbs and H. Stern, *Cell*, **40**, 785-790 (1985).

Structure of Viral DNA in a Rat Cell Line, GY1, Transformed by Ad12 *Hind* III Fragment-G, S. Kudo, H. Sugisaki, M. Takanami, T. Yamashita, M. Ishino, Y. Shimizu, Y. Iida and K. Fujinaga, *Virology*, **148**, 133–145, (1986).

Complete Structure of an Insecticidal Crystal Protein Gene from *Bacillus thuringiensis*, S. Shibano, A. Yamagata, T. Amachi and M. Takanami, *Bacillus Molecular Genetics*, 307–320 (1986).

The Folding Type of a Protein is Relevant to the Amino Acid Composition, H. Nakashima, K. Nishikawa and T. Ooi, *J. Biochem.*, **99**, 153–162 (1986).

Comparison of Amino Acid Sequences between Phosphoenolpyruvate Carboxylases from *Escherichia coli* (Allosteric) and *Anacystis nidulans* (Non-allosteric): Identification of Conserved and Variable Regions, S. Ishijima, F. Katagiri, T. Kodaki, K. Izui, H. Katsuki, K. Nishikawa, H. Nakashima and T. Ooi, *Biochem. Biophys. Res. Commun.*, **133**, 436–441 (1985).

Amino Acid Sequence Homology Applied to the Prediction of Protein Secondary Structures, and Joint Prediction with Existing Methods, K. Nishikawa and T. Ooi, *Biochim. Biophys. Acta*, **871**, 45–54 (1986).

Examination of Protein Sequence Homologies: II. Six *Escherichia coli* L7/L12-Type Ribosomal “A” Protein Sequences from Eukaryotes and Metabacteria, Contrasted with Those from Prokaryotes, E. Otaka, T. Ooi, T. Kumazaki and T. Itoh, *J. Mol. Evol.*, **22**, 342–350 (1985).

Relation between Sequence Similarity and Structural Similarity in Proteins. Role of Important Properties of Amino Acids, A. Kidera, Y. Konishi, T. Ooi and H.A. Scheraga, *J. Prot. Chem.*, **4**, 265–297 (1985).

Statistical Analysis of the Physical Properties of the 20 Naturally Occurring Amino Acids, A. Kidera, Y. Konishi, M. Oka, T. Ooi and H.A. Scheraga, *J. Prot. Chem.*, **4**, 23–55 (1985).

Information Analysis and Sequence Data Banks of DNA and Protein, T. Ooi, S. Tsuda and H. Nishikawa, *Proc. Int. Symp. Computer Anal. for Life Sci.*, 212–219 (1986).

Flow of Information in Biological Systems, T. Ooi, *J. Mol. Sci.*, **4**, 75–88 (1986).

Dynamic Study of Crystallization of Poly(ϵ -caprolactone) and Poly(ϵ -caprolactone)/Poly(vinyl chloride) Blend, S. Nojima, W. Kosaka, T. Ashida, Y. Muroga, T. Ueki, Y. Hiragi, M. Kataoka, Y. Izumi, H. Tagawa and Y. Amemiya, *Polymer Journal*, **17**, 1229–1231 (1985).

Some Remark on Imogolite Mesophase, K. Kajiwara, N. Donkai, Y. Fujiyoshi, Y. Hiragi, H. Urakawa and H. Inagaki, *Bull. Inst. Chem. Res., Kyoto Univ.*, **63**, 320–331 (1985).

Statistical Procedure for Deconvoluting Experimental Data, Y. Hiragi, H. Urakawa, and K. Tanabe, *J. Appl. Phys.*, **58**, 5–11 (1985).

X-Ray Scattering Study on Hemoglobin Solution with Synchrotron Radiation: A Simple Analysis of Scattering Profile at Moderate Angles in Terms of Arrangement of Subunits, T. Ueki, Y. Inoko, M. Kataoka, Y. Amemiya and Y. Hiragi, *J. Biochem.*, **99**, 1127–1136 (1986).

Some Remarks on Imogolite Mesophase, K. Kajiwara, N. Donkai, Y. Fujiyoshi, Y. Hiragi, H. Urakawa, and H. Inagaki, *Bull. Inst. Chem. Res., Kyoto Univ.*, **63**, 320–33 (1985).

Dilute Solution Properties of Imogolite, N. Donkai, H. Inagaki, K. Kajiwara, H. Urakawa and M. Schmidt, *Makromol. Chem.*, **186**, 2623–2638 (1985).

Molecular Dynamics Using Internal Coordinates I. Infrared Spectra of Sulfur Dioxide, S. Hayashi, M. Oobatake, T. Ooi, and K. Machida, *Bull. Chem. Soc. Japan*, **58**, 1105–1108 (1985).

Molecular Orientation of Plastocyanin on Spinach Thylakoid Membranes as determined by Acetylation of Lysine Residues, M. Takano, M. Takahashi, M. Oobatake, and K. Asada, *J. Biochemistry (Tokyo)*, **98**, 1333–1340 (1985).

Optimization of Amino Acid Parameters for Correspondence of Sequence to Tertiary Structures of Proteins, M. Oobatake, Y. Kubota, and T. Ooi, *Bull. Inst. Chem. Res., Kyoto Univ.*, **63**, 82–94 (1985).

Structure of the Human Interleukin-2 Receptor Gene, W. J. Leonard, J.M. Depper, M. Kanehisa, M. Krönke, N.J. Peffer, P.B. Svetlik, M. Sullivan, and W.C. Greene, *Science*, **230**, 633–639 (1985).

The GenBank Nucleic Acid Sequence Database, G. Burks, J.W. Fickett, W.B. Goad, M. Kanehisa, F.I. Lewitter, W.P. Rindone, C.D. Swindell, C.-S. Tung and H.S. Bilofsky, *Comp. Appl. Biosci.*, **1**, 225–233 (1985).

Design of DNAs and Proteins: Application of Computer Databases, M. Kanehisa, *Seibutubuturi no Tenkai (Development of Biophysics)*, 39–54 (1986).

IDEAS: Integrated Database and Extended Analysis System for Nucleic Acids and Proteins, M. Kanehisa, *Kyoto University Computer Center News*, **19**, 247–259 (1986).

Design and Application of DNA and Protein Data Banks, M. Kanehisa, *Tanpakusitu Kakusan Koso (Protein, Nucleic Acid and Enzyme)*, **29**, 187–194 (1986).

Genetic Information Databases, M. Kanehisa, *Parity*, **1**(7), 59–61 (1986), Communication in Japanese.

L-Aspartate Ammonia-lyase and Fumarate Hydratase Share Extensive Sequence Homology, J.S. Takagi, M. Tokushige, Y. Shimura, and M. Kanehisa,

Biochem. Biophys. Res. Commun., **138**, 568-572 (1986).

Two Independent Mutations are Required for Temperature-sensitive Cell Transformation by a Rous Sarcoma Virus Temperature-sensitive Mutant, M. Nishizawa, B.J. Mayer, T. Takeya, T. Yamamoto, K. Toyoshima, H. Hanafusa and S. Kawai, *J. of Virology*, **56**, 743-749 (1985).

Classification and Characterization of Oncogene Products Based on their Homology, J. Kato and T. Takeya, *Tanpakusitu, Kakusan, Koso (Protein, Nucleic Acid and Enzyme)*, 151-158 (1986).