

# On pair-splitting and pair-reaping pairs of $\omega$

Hiroaki Minami

## Abstract

In this paper we investigate variations of splitting number and reaping number, pair-splitting number  $\mathfrak{s}_{pair}$ , pair-reaping number  $\mathfrak{r}_{pair}$ . We prove that it is consistent that  $\mathfrak{s}_{pair} < \mathfrak{d}$ . We also prove it is consistent that  $\mathfrak{r}_{pair} > \mathfrak{b}$ .

## Introduction

The splitting number  $\mathfrak{s}$  and the reaping number  $\mathfrak{r}$  are cardinal invariants related to the structure  $\mathcal{P}(\omega)/fin$ .

For  $X, Y \in [\omega]^\omega$  we say  $X$  splits  $Y$  if  $X \cap Y$  and  $Y \setminus X$  are infinite. We call  $\mathcal{S} \subset [\omega]^\omega$  a splitting family if for each  $Y \in [\omega]^\omega$ , there exists  $X \in \mathcal{S}$  such that  $X$  splits  $Y$ . The splitting number  $\mathfrak{s}$  is the least size of a splitting family.

We call  $\mathcal{R}$  a reaping family if for each  $X \in [\omega]^\omega$ , there exists  $Y \in \mathcal{R}$  such that  $Y$  is not split by  $X$ , that is,  $X \cap Y$  is finite or  $Y \setminus X$  is finite. The reaping number  $\mathfrak{r}$  is the least size of a reaping family.

We shall study variations of splitting number and reaping number, pair-splitting number  $\mathfrak{s}_{pair}$  and pair-reaping number  $\mathfrak{r}_{pair}$ . They are introduced and investigated in [7] to analyze dual-reaping number  $\mathfrak{r}_d$  and dual-splitting number  $\mathfrak{s}_d$  which are reaping number and splitting number for the structure of all infinite partitions of  $\omega$  ordered by “almost coarser”  $(([\omega]^\omega, \leq^*))$  respectively.

We call  $A \subset [\omega]^2$  unbounded if for  $k \in \omega$ , there exists  $a \in A$  such that  $a \cap k = \emptyset$ . For  $X \in [\omega]^\omega$  and unbounded  $A \subset [\omega]^2$ ,  $X$  pair-splits  $A$  if there exist infinitely many  $a \in A$  such that  $a \cap X \neq \emptyset$  and  $a \setminus X \neq \emptyset$ . We call  $\mathcal{S} \subset [\omega]^\omega$  a pair-splitting family if for each unbounded  $A \subset [\omega]^2$ , there exists  $X \in \mathcal{S}$  such that  $X$  pair-splits  $A$ . The pair-splitting number  $\mathfrak{s}_{pair}$  is the least size of a pair-splitting family.

We call  $\mathcal{R} \subset \mathcal{P}([\omega]^2)$  a pair-reaping family if for each  $A \in \mathcal{R}$ ,  $A$  is unbounded and for  $X \in [\omega]^\omega$ , there exists  $A \in \mathcal{R}$  such that  $X$  doesn't pair-split  $A$ . The pair-reaping number  $\tau_{pair}$  is the least size of a pair-reaping family.

In [7] it is proved that there is the following relationship between  $\tau_{pair}$ ,  $\mathfrak{s}_{pair}$  and other cardinal invariants.

**Proposition 0.1** 1.  $\mathfrak{s}_{pair} \leq non(\mathcal{M}), non(\mathcal{N})$ .

2.  $\tau_{pair} \geq cov(\mathcal{M}), cov(\mathcal{N})$ .

3.  $\mathfrak{s}_{pair} \geq \mathfrak{s}$ .

4.  $\tau_{pair} \leq \mathfrak{r}, \mathfrak{s}_d$ .


It is not known that  $\mathfrak{r}_d \leq \mathfrak{s}_{pair}$  or not.

**Question 0.1**  $\mathfrak{r}_d \leq \mathfrak{s}_{pair}$ ?

$\mathfrak{s} \leq \mathfrak{d}$  and  $\mathfrak{r} \geq \mathfrak{b}$  hold (see in [2]). And Kamo proved the following statement in [7]:

**Theorem 0.1**  $\mathfrak{r}_d \leq \mathfrak{d}$  and  $\mathfrak{s}_d \geq \mathfrak{b}$ .

So we have the following diagram:



An arrow  $\kappa \rightarrow \lambda$  denotes the inequality  $\kappa \geq \lambda$ .

In [7] by using finite support iteration of Hechler forcing, the following consistency results are proved.

**Theorem 0.2** *It is consistent that  $s_{pair} < \text{add}(\mathcal{M})$ . Dually it is consistent that  $\tau_{pair} > \text{cof}(\mathcal{M})$ .*

$\tau_{pair}$  is a lower bound of  $\tau$  and  $\mathfrak{s}$  and  $s_{pair}$  is an upper bound of  $\mathfrak{s}$  (and maybe of  $\tau_d$ ). So it is natural to ask the following question.

**Question 0.2**  *$s_{pair} \leq \mathfrak{d}$ ? Dually  $\tau_{pair} \geq \mathfrak{b}$ ?*

In the present paper we shall investigate the relationship between  $\tau_{pair}$  and  $\mathfrak{b}$  and the relationship between  $s_{pair}$  and  $\mathfrak{d}$ . In section 1 we shall prove the consistency of  $s_{pair} > \mathfrak{d}$ . In section 2 we shall show the consistency of the consistency of  $\tau_{pair} < \mathfrak{b}$ . In section 3 we mention the development of results in section 1 and 2.

# 1 pair-splitting number and dominating number

**Notation and Definition** We present the related notions. We use standard set theoretical conventions and notation. For a set  $X$ ,  $X^\omega$  denotes the set of all functions from  $\omega$  to  $X$ . For  $f, g \in \omega^\omega$ ,  $f$  dominates  $g$ , written  $f \leq^* g$ , if for all but finitely many  $n \in \omega$   $g(n) \leq f(n)$ . We call  $\mathcal{F}$  a dominating family if for each  $g \in \omega^\omega$  there exists  $f \in \mathcal{F}$  such that  $g \leq^* f$ . The dominating number  $\mathfrak{d}$  is the least size of a dominating family.

We call  $\mathcal{G}$  an unbounded family if for each  $f \in \omega^\omega$  there exists  $g \in \mathcal{G}$  such that  $g \not\leq^* f$ , i.e., there exist infinitely many  $n \in \omega$  such that  $g(n) > f(n)$ . The unbounded number  $\mathfrak{b}$  is the least size of an unbounded family.

For a set  $X$ ,  $X^{<\omega}$  denote the set of all functions from natural numbers to  $X$ .

We call partial ordering  $(T, <)$  a tree if the set  $\{s \in T : s < t\}$  is well-ordered by  $<$ . We say  $T$  is a tree on  $X$  if  $T$  is a subtree of  $(X^{<\omega}, \subset)$ . For a tree  $T$  and  $t \in T$ ,  $\text{succ}_T(t)$  is the set of all immediate successors of  $t$  in  $T$ . For a tree  $T$ ,  $\text{stem}(T)$  is the first element of  $T$  which has at least 2-many immediate successors.

**Theorem 1.1** *It is consistent  $\mathfrak{s}_{\text{pair}} > \mathfrak{d}$ .*

To prove theorem 1.1, we shall construct a proper forcing notion which enlarges  $\mathfrak{s}_{\text{pair}}$  and is  $\omega^\omega$ -bounding to show  $\mathfrak{d}$  is preserved by the forcing notion.

**Definition 1.1** [4, pp340] *A forcing notion  $\mathbb{P}$  is  $\omega^\omega$ -bounding if*

$$\Vdash_{\mathbb{P}} \forall f \in \omega^\omega \cap V[G] \exists g \in \omega^\omega \cap V (f \leq^* g).$$

The  $\omega^\omega$ -boundingness has the following good property.

**Theorem 1.2** [4, pp341] *The countable support iteration of proper  $\omega^\omega$ -bounding forcing notions is  $\omega^\omega$ -bounding.*

To prove theorem 1.1 we shall construct a forcing notion which consists of finitely branching trees on  $[\omega]^2$  such that the set of successors of any node carries a norm as [8].

To present the desired forcing notion, we define “norm” for finite subsets of  $[\omega]^2$ . Let  $R(n)$  be a natural number such that if  $m \geq R(n)$ , then for any

function  $f : [m]^2 \rightarrow 2$  there exists  $H \in [m]^n$  such that  $|f([H]^2)| = 1$ . Then recursively define  $l_1 = 3$ ,  $l_{n+1} = \max\{2l_n, R(l_n)\}$ . Then for a finite subset  $A$  of  $[\omega]^2$   $\text{norm}(A) \geq n$  if  $A$  contains a complete graph with  $l_n$ -many vertices.

This norm has the following properties:

**Proposition 1.1** *For a finite subset  $A$  of  $[\omega]^2$ ,*

1.  *$\text{norm}(A) \geq 1$  implies for any  $X \in [\omega]^\omega$  there exists  $a \in A$  such that  $a \cap X = \emptyset$  or  $a \subset X$ .*
2. *Suppose  $\text{norm}(A) \geq n+1$ . For  $X \in [\omega]^\omega$  let  $A_X^0 = \{a \in A : a \cap X = \emptyset\}$  and  $A_X^1 = \{a \in A : a \subset X\}$ . Then  $\text{norm}(A_X^0) \geq n$  or  $\text{norm}(A_X^1) \geq n$ .*
3. *Suppose  $\text{norm}(A) \geq n+1$ . If  $A = A_0 \cup A_1$ , then  $\text{norm}(A_0) \geq n$  or  $\text{norm}(A_1) \geq n$ .*

**Proof of proposition 1.1**

1. Since  $\text{norm}(A) \geq 1$ ,  $A$  contains a complete graph  $A' \subset A$  with 3-many vertices. Then for any 2-coloring of the vertices of  $A'$ , there exists an edge whose vertices have the same color. So there exists  $a \in A' \subset A$  such that  $a \subset X$  or  $a \cap X = \emptyset$ .

2. Since  $\text{norm}(A) \geq n+1$ ,  $A$  contain a complete graph  $A'$  with  $l_{n+1}$ -many vertices. So for each  $X \subset \omega$ ,  $X$  contains  $l_n$ -many vertices of  $A'$  or  $X$  doesn't meet  $l_n$ -many vertices of  $A'$  because  $l_{n+1} \geq 2l_n$ . Anyway  $A_X^0 = \{a \in A : a \cap X = \emptyset\}$  or  $A_X^1 = \{a \in A : a \subset X\}$  contains a complete graph with  $l_n$ -many vertices. Therefore  $\text{norm}(A_X^0) \geq n$  or  $\text{norm}(A_X^1) \geq n$ .

3. Since  $\text{norm}(A) \geq n+1$ ,  $A$  contain a complete graph  $A'$  with  $l_{n+1}$ -many vertices. Define  $f : A' \rightarrow 2$  by  $f(a) = i$  if  $a \in A_i$  for  $i < 2$ . Since  $l_{n+1} \geq R(l_n)$ , there exists a complete graph  $A^* \subset A'$  which has  $l_n$ -many vertices of  $A'$  and  $|f[A^*]| = 1$ . So  $A^* \subset A_0$  or  $A^* \subset A_1$ . Hence  $\text{norm}(A_0) \geq n$  or  $\text{norm}(A_1) \geq n$ .  $\square$

Then let  $\mathbb{P}$  be the set of perfect trees such that

1.  $T$  is a finitely branching tree on  $[\omega]^2$ ,
2. for any branch of  $T$  and  $n \in \omega$  there exist  $m \geq n$  such that whenever  $t \in T$  with  $|t| \geq m$ ,  $\text{norm}(\text{succ}_T(t)) \geq n$ .

For  $T$  and  $S$  in  $\mathbb{P}$ ,  $T \leq S$  if  $T \subset S$ .

**Lemma 1.1** *Let  $G$  be a generic filter on  $\mathbb{P}$  and  $A_G = \bigcap \{T : T \in G\}$ . Then  $A_G \subset [\omega]^2$  and for any  $X \in [\omega]^\omega \cap V$ ,  $X$  doesn't pair-split  $A_G$ .*

**Proof** For  $X \in [\omega]^\omega$  define a subset  $D_X$  of  $\mathbb{P}$  by  $T \in D_X$  if for all  $t \in T \setminus \{s : s \subset \text{stem}(T)\}$  and  $a \in \text{succ}_T(t)$ ,  $a \subset X$  or  $a \cap X = \emptyset$ . Then for a given  $S \in \mathbb{P}$  we can find  $T \leq S$  such that for all  $t \in T \setminus \{s : s \subset \text{stem}(T)\}$  and  $a \in \text{succ}_T(t)$ ,  $a \subset X$  or  $a \cap X = \emptyset$  by 1 and 2 in Proposition 1.1. So  $D_X$  is dense. So  $X$  doesn't pair-split  $A_G$ . □

By this lemma,  $\mathbb{P}$  adds an infinite subset of  $[\omega]^2$  which is not pair-split by any infinite subset of  $\omega$  in ground model. Therefore  $\omega_2$ -stage countable support iteration of  $\mathbb{P}$  forces  $\mathfrak{s}_{\text{pair}} = \omega_2$ .

From now on we shall prove  $\mathbb{P}$  is  $\omega^\omega$ -bounding and proper.

For  $T \in \mathbb{P}$ , let  $\text{ess}(T) = \{t \in T : \text{stem}(T) \subset t\}$ . For  $T, S \in \mathbb{P}$ ,  $T \leq^* S$  if  $T \leq S$  and for all  $t \in \text{ess}(T)$ ,  $\text{norm}(\text{succ}_T(t)) \geq \text{norm}(\text{succ}_S(t)) - 1$ .  $T \leq_m S$  if  $T \leq S$  and for all  $t \in T$  with  $\text{norm}(\text{succ}_S(t)) \leq m$ , we have  $\text{succ}_S(t) \subset T$ .

As [8] we can prove the following lemmata.

**Lemma 1.2** *If  $S \in \mathbb{P}$  and  $W \subset S$ , then there is some  $T \leq^* S$  such that*

*I. every branch of  $T$  meets  $W$ , or else*

*II.  $T$  is disjoint from  $W$ .*

**Proof** Let  $S^W$  be the set of all  $s \in S$  such that there exists  $S' \leq^* S_s$  such that every branch of  $S'$  meets  $W$  where  $S_s$  is the set of  $t \in S$  comparable to  $s$ .

If  $\text{stem}(S) \in S^W$ , then (I) holds. Otherwise we will construct  $T \leq^* S$  which satisfies (II).

Suppose  $\text{stem}(S) \notin S^W$ . Recursively construct  $t \in T$  with  $|t| = n$ . If  $n \leq |\text{stem}(T)|$ ,  $t \in T$  with  $|t| = n$  if  $t \in S$  with  $|t| = n$ . If  $n \geq |\text{stem}(T)|$ , assume  $t \in T$  with  $|t| \leq n$  are given and  $t \notin S^W$  for  $t \in T$  with  $|t| \leq n$ . For  $t \in T$  with  $|t| = n$ , let  $A^t = \text{succ}_S(t)$ ,  $A_0^t = S^W \cap A^t$  and  $A_1^t = A^t \setminus A_0^t$ . By Proposition 1.1 (iii),  $\text{norm}(A_i^t) \geq \text{norm}(A^t) - 1$  for some  $i < 2$ . Since  $t \notin S^W$ , there is no  $S' \leq^* S_t$  such that  $S'$  holds I. So  $\text{norm}(A_0^t) < n$ . Hence  $\text{norm}(A_1^t) \geq \text{norm}(A^t) - 1$ . Define  $t \in T$  with  $|t| = n + 1$  if  $t \upharpoonright n \in T$  and  $t(n) \in A_1^{t \upharpoonright n}$ . Then for any  $t \in T$  with  $|t| = n + 1$ ,  $t \notin S^W$ .

By construction  $T \leq^* S$  and satisfies II. □

**Lemma 1.3** *Let  $\dot{\alpha}$  be a  $\mathbb{P}$ -name for an ordinal. Let  $S \in \mathbb{P}$  such that for  $t \in S \setminus \{s : s \subset \text{stem}(S)\}$ ,  $\text{norm}(\text{succ}_S(t)) > m + 1$ . Then there exists  $T \leq_m S$  and a finite subset  $w$  of ordinal such that  $T \Vdash \dot{\alpha} \in w$ .*

**Proof** Let  $W$  be the set of nodes  $s \in S$  such that there exists  $S^s \leq_m S_s$  which decides the value  $\dot{\alpha}$ .

We shall prove that there exists  $S_1 \leq^* S$  such that every branch of  $S_1$  meets  $W$ . Suppose  $S' \leq^* S$  and  $S'' \leq S'$  such that  $S'' \Vdash \dot{\alpha} = \beta$  for some  $\beta$ . Then for some  $t \in S''$  for each extension  $s$  of  $t$  in  $S''$  satisfies  $\text{norm}(\text{succ}_{S''}(s)) > m$ . Because  $S''_t \leq_m S_t$  and  $S''$  decides  $\dot{\alpha}$ ,  $t \in W$ . Hence by Lemma 1.2 there exists  $S_1 \leq^* S$  which satisfies I in Lemma 1.2.

Let  $S_1 \leq^* S$  such that every branch of  $S_1$  meets  $W$ . Let  $W_0$  be the set of minimal elements of  $W$  in  $S_1$ . Since  $S_1$  is finitely branching,  $W_0$  is finite. (Otherwise, by König's Lemma we can construct infinitely branch which doesn't meet  $W$ ). For  $v \in W_0$  choose  $T^v \leq_m S_v$  and  $\alpha_v$  such that  $T^v \Vdash \dot{\alpha} = \alpha_v$ . Put  $T = \bigcup_{v \in W_0} T^v$  and  $w = \{\alpha_v : v \in W_0\}$ . Then  $T \leq_m S$  and  $T \Vdash \dot{\alpha} \in w$ . □

**Lemma 1.4** *If  $S \in \mathbb{P}$ ,  $\dot{\alpha}$  be a  $\mathbb{P}$ -name for an ordinal and  $m < \omega$ . Then there exists  $T \leq_m S$  and a finite set of ordinals  $w$  such that  $T \Vdash \dot{\alpha} \in w$ .*

**Proof** Choose  $k \in \omega$  such that for any  $s \in S$  with  $|s| \geq k$   $\text{norm}(\text{succ}_S(s)) > m + 1$ . For each  $s \in S$  with  $|s| = k$ , apply Lemma 1.3 to  $S_s$ , pick  $T^s \leq_m S_s$  and a finite set of ordinals  $w_s$  so that  $T^s \Vdash \dot{\alpha} \in w_s$ . Put  $T = \bigcup_{s \in S, |s|=k} T^s$  and  $w = \bigcup_{s \in S \cap \omega^k} w_s$ . Then  $T \leq_m S$  and  $T \Vdash \dot{\alpha} \in w$ . Since  $S$  is finitely branching,  $w$  is a finite set. □

**Proof of theorem 1.1** Lemma 1.4 implies that  $\mathbb{P}$  is  $\omega^\omega$ -bounding. Given a  $\mathbb{P}$ -name for a function  $f$  from  $\omega$  to  $\omega$  and  $S \in \mathbb{P}$ , we can construct a sequence  $\langle T_n : n \in \omega \rangle$  of conditions of  $\mathbb{P}$  such that  $T_0 = S$ ,  $T_{n+1} \leq_n T_n$  and for each  $n \in \omega$ , there exists some finite  $w_n$  of natural numbers such that  $T_n \Vdash f(n) \in w_n$ . Then there exists  $T \in \mathbb{P}$  such that  $T \leq_n T_n$  and  $T \Vdash \forall n \in \omega (f(n) \in w_n)$ . Put  $g(n) = \max\{w_n\}$ . Then  $T \Vdash \forall n \in \omega (f(n) \leq g(n))$ . So  $\mathbb{P}$  is  $\omega^\omega$ -bounding. Also this claim say  $\mathbb{P}$  satisfies Baumgartner's Axiom A. Hence  $\mathbb{P}$  is proper.

Hence the  $\omega_2$ -stage countable support iteration of  $\mathbb{P}$  is  $\omega^\omega$ -bounding by theorem 1.2. Therefore if  $V \models CH$ , then the  $\omega_2$ -stage countable support iteration of  $\mathbb{P}$  forces  $\omega^\omega \cap V$  is a dominating family. So the  $\omega_2$ -stage countable support iteration of  $\mathbb{P}$  forces  $\mathfrak{d} = \omega_1$ . Hence it is consistent that  $s_{\text{pair}} > \mathfrak{d}$ . □

Since  $\mathfrak{s} \leq \mathfrak{d}$  (see[2]), we have the following corollary.

**Corollary 1.1** *It is consistent that  $\mathfrak{s} < \mathfrak{s}_{pair}$ .*

## 2 pair-reaping number and unbounded number

To show the consistency of  $\mathfrak{r}_{pair} < \mathfrak{b}$ , we shall use the Laver forcing  $\mathbb{L}$ .  $\mathbb{L}$  is defined by  $T \in \mathbb{L}$  if  $T \subset \omega^{<\omega}$  is a tree and for  $s \in T$  with  $stem(T) \subset s$ ,  $|succ_T(s)| = \aleph_0$ .  $\mathbb{L}$  is ordered by inclusion. Then  $\mathbb{L}$  adds an unbounded real.

**Proposition 2.1** *Let  $G$  be a  $\mathbb{L}$ -generic over  $V$  and  $f_G = \bigcup\{stem(T) : T \in G\}$ . Then  $f_G \in \omega^\omega$  and  $f_G$  dominates for all  $g \in \omega^\omega \cap V$ .*

*Therefore if  $\mathbb{L}_{\omega_2}$  is  $\omega_2$ -stage countable support iteration of Laver forcing, then  $V^{\mathbb{L}_{\omega_2}} \models \mathfrak{b} = \mathfrak{c}$ .*

By using  $\omega_2$ -stage countable support iteration of Laver forcing, we shall construct ZFC model which satisfies  $\mathfrak{r}_{pair} < \mathfrak{b}$ .

**Theorem 2.1** *It is consistent  $\mathfrak{r}_{pair} < \mathfrak{b}$ .*

By proposition 2.1 it is enough  $\mathbb{L}$  preserves  $\mathfrak{r}_{pair}$ . We shall use the Laver property.

**Definition 2.1** [4] *A forcing notion  $\mathbb{P}$  have the Laver property if for every  $H : \omega \rightarrow \omega \in V$*

$$\Vdash \forall f \in (\prod_{n \in \omega} H(n)) \cap V[G] \exists A : \omega \rightarrow \omega^{<\omega} \in V \forall n \in \omega (f(n) \in A(n) \wedge |A(n)| \leq 2^n)$$

**Theorem 2.2** [4] *The Laver property is preserved under countable support iteration of proper forcing notions.*

**Theorem 2.3** [1, pp353] *The Laver forcing  $\mathbb{L}$  has the Laver property.*

So  $\mathbb{L}_{\omega_2}$  has the Laver property. If forcing notion  $\mathbb{P}$  has the Laver property, then  $\mathbb{P}$  has the following good property:

**Lemma 2.1** *Let  $\mathbb{P}$  be a forcing notion satisfying the Laver property. Then  $\Vdash_{\mathbb{P}} \forall \dot{X} \in V[G] \exists A \in V (\dot{X} \text{ doesn't pair-split } A)$ .*



**Proof** Let  $p \in \mathbb{P}$ . Let  $\Pi = \langle I_n : n \in \omega \rangle$  be an interval partition of  $\omega$  such that  $|I_n| = 2^{2^n} + 1$ . Then  $\langle \dot{X} \upharpoonright I_n : n \in \omega \rangle \in \Pi_{n \in \omega} 2^{I_n}$ . By the Laver property there exists  $q \leq_{\mathbb{P}} p$  such that  $\langle A_n : n \in \omega \rangle \in V$  such that  $A_n \subset 2^{I_n}$ ,  $|A_n| \leq 2^n$  and  $q \Vdash \forall n \in \omega (\dot{X} \upharpoonright I_n \in A_n)$ . For each  $n \in \omega$   $\{\langle \sigma(k) : \sigma \in A_n \rangle : k \in A_n\}$  is at most  $2^{2^n}$ -many element. But  $|I_n| = 2^{2^n} + 1$ . So there exists  $k_0^n$  and  $k_1^n$  in  $I_n$  such that  $k_0^n \neq k_1^n$  and  $\langle \sigma(k_0^n) : \sigma \in A_n \rangle = \langle \sigma(k_1^n) : \sigma \in A_n \rangle$ . Put  $a_n = \{k_0^n, k_1^n\}$  and  $A = \{a_n : n \in \omega\} \in V$ . Then  $q \Vdash \dot{X} \upharpoonright I_n \cap a_n = \emptyset$  or  $a_n \subset \dot{X} \upharpoonright I_n$  for  $n \in \omega$ . Therefore  $q \Vdash \dot{X}$  doesn't pair-split  $A$ .  $\square$

**Proof of theorem 2.1** Suppose  $V \models CH$ . By theorem 2.2 and 2.3  $\mathbb{L}_{\omega_2}$  has the Laver property. By lemma 2.1 for each  $X \in [\omega]^\omega \cap V^{\mathbb{L}_{\omega_2}}$  there exists an unbounded  $A \subset [\omega]^2$  such that  $V^{\mathbb{L}_{\omega_2}} \models X$  doesn't pair-split  $A$ . So  $\{A \subset [\omega]^2 : A \text{ unbounded}\} \cap V$  is pair-reaping family. Since  $V \models CH$ ,  $\{A \subset [\omega]^2 : A \text{ unbounded}\} \cap V$  has the cardinality at most  $\omega_1$ . Therefore  $V^{\mathbb{L}_{\omega_2}} \models \tau_{pair} < \mathfrak{b}$ .  $\square$

Since  $\tau \geq \mathfrak{b}$  (see[2]), we have the following corollary.

**Corollary 2.1** *It is consistent that  $\tau > \tau_{pair}$ .*

In [5] Masaru Kada introduces a cardinal invariant associated with the Laver property.

Let  $\mathcal{S}$  be the collection of functions  $\phi$  from  $\omega$  to  $[\omega]^{<\omega}$  such that  $|\phi(n)| \leq n + 1$ .  $\mathfrak{l}$  is the smallest cardinal  $\kappa$  such that for every  $h \in \omega^\omega$  there is a set  $\Phi \subset \mathcal{S}$  with cardinality  $\kappa$  so that, for every  $f \in \omega^\omega$  with  $f(n) < h(n)$  for all  $n < \omega$ , there is  $\phi \in \Phi$  such that for all but finitely many  $n \in \omega$  we have  $f(n) \in \phi(n)$ .

As the proof of theorem 2.1 we can prove the following statement.

**Corollary 2.2**  $\tau_{pair} \leq \mathfrak{l}$ .

Pawlikowski shows that the dual notion to the definition of  $\mathfrak{l}$  is the characterization of  $\text{trans-add}(\mathcal{N})$ , transitive additivity of null ideal (see [1, pp91]). That is,  $\text{trans-add}(\mathcal{N})$  is the smallest size of  $\leq^*$ -bounded family  $F \subset \omega^\omega$  such that for every  $\phi \in \mathcal{S}$  there is  $f \in F$  such that for infinitely many  $n \in \omega$  such that  $f(n) \notin \phi(n)$ .

Then the dual inequality to the corollary 2.2 holds.

**Proposition 2.2**  $\mathfrak{s}_{pair} \geq \text{trans-add}(\mathcal{N})$ .

It is known the following relation between  $\text{trans-add}(\mathcal{N})$  and  $\mathfrak{d}$ .

**Theorem 2.4** [6] *It is consistent that  $\text{trans-add}(\mathcal{N}) > \mathfrak{d}$ .*

By theorem 2.4 and proposition 2.2 it is consistent that  $\mathfrak{s}_{\text{pair}} > \mathfrak{d}$ .

### 3 Further results

In this section we mention the development of above results in the paper [3] written by Hrušák, Meza-Alcántara and the author.

Hrušák and Meza-Alcántara study cardinal invariants of ideals on  $\omega$  and they define the pair-splitting number and the pair-reaping number independently of the author and they showed the pair-splitting number and the pair-reaping number are described as cardinal invariants of an ideal on  $\omega$ .

Let  $\mathcal{I}$  be an ideal on  $\omega$ . Define the cardinal invariants associate with  $\mathcal{I}$  by

$$\begin{aligned} \text{cov}^*(\mathcal{I}) &= \min\{|\mathcal{A}| : \mathcal{A} \subset \mathcal{I} \wedge \forall I \in \mathcal{I} \exists A \in \mathcal{A} (|A \cap I| = \aleph_0)\} \\ \text{non}^*(\mathcal{I}) &= \min\{|\mathcal{A}| : \mathcal{A} \subset [\omega]^\omega \wedge \forall I \in \mathcal{I} \exists A \in \mathcal{A} (|A \cap I| < \aleph_0)\}. \end{aligned}$$

**Theorem 3.1** [3] *Let  $\mathcal{G}_{FC}$  be an ideal on  $[\omega]^2$  defined by*

$$\mathcal{G}_{FC} = \{A \subset [\omega]^2 : \chi(\omega, A) < \aleph_0\}$$

where  $\chi(\omega, A) = \min\{k \in \omega : \exists f : \omega \rightarrow k \forall a \in A (|f[a]| = 2)\}$ .

*Then  $\text{non}^*(\mathcal{G}_{FC}) = \mathfrak{r}_{\text{pair}}$  and  $\text{cov}^*(\mathcal{G}_{FC}) = \mathfrak{s}_{\text{pair}}$ .*

From now on we assume  $2^\omega$  is equipped with product topology and the topology of  $\mathcal{P}(\omega)$  is induced by identification of each subset of  $\omega$  with its characteristic function.

Then  $\mathcal{G}_{FC}$  is an  $F_\sigma$ -ideal on  $[\omega]^2$ . As theorem 2.4, 1.1 and theorem 2.1 we can show the following theorem.

**Theorem 3.2** *Suppose  $\mathcal{I}$  is an  $F_\sigma$ -ideal on  $\omega$ .*

1. [6] *It is consistent that  $\mathfrak{d} < \text{cov}^*(\mathcal{I})$ .*
2. [3] *It is consistent that  $\mathfrak{b} > \text{non}^*(\mathcal{I})$ .*

Also the following statement holds as corollary 2.2 and proposition 2.2.

**Corollary 3.1** *Suppose  $\mathcal{I}$  is an  $F_\sigma$ -ideal.*

1. *If  $\text{non}^*(\mathcal{I}) \neq \omega$ , then  $\text{non}^*(\mathcal{I}) \leq \mathfrak{l}$ .*
2. *If  $\text{non}^*(\mathcal{I}) \neq \omega$ , then  $\text{cov}^*(\mathcal{I}) \geq \text{trans-add}(\mathcal{I})$ .*

So many results in section 1 and 2 follows from theorem 3.2 and corollary 3.1.

## Acknowledgment

While carrying out the research for this paper, I discussed my work with Jörg Brendle. He gave me helpful advice. I greatly appreciate his help.

I also thank Shizuo Kamo for pointing out some remarks. I also thank Masaru Kada for pointing out corollary 2.2, proposition 2.2 and another proof for theorem 2.1 from proposition 2.2 and theorem 2.4.

I thank to Michael Hrušák and David Meza-Alcántara who point out the relation between their results and my research. The collaboration produce theorem 3.2 2 and corollary 3.1.

I also thank Teruyuki Yorioka and Noboru Osuga for pointing out some mistake of proof and for suggestions which improved the presentation of this work.

Finally I thank members of Arai Project at Kobe University for much support while carrying out the research.

## References

- [1] Tomek Bartoszyński, Haim Judah, “Set theory. On the structure of the real line”. A K Peters, Ltd., Wellesley, MA, 1995.
- [2] Andreas Blass, “Combinatorial cardinal characteristics of the continuum”, in Handbook of Set Theory (A.Kanamori et al.,eds.),to appear.
- [3] Michael Hrušák David Meza-Alcántara and Hiroaki Minami, “Around pair-splitting and pair-reaping number”, preprint.
- [4] Martin Goldstern, “Tools for your forcing construction”. Set theory of the reals (Ramat Gan, 1991), 305–360, Israel Math. Conf. Proc., 6, Bar-Ilan Univ., Ramat Gan, 1993.

- [5] Masaru Kada, “More on Cichoń’s diagram and infinite games”, *J. Symbolic Logic* 65 (2000), no. 4, 1713–1724.
- [6] Laflamme, Claude, “Zapping small filters”, *Proc. Amer. Math. Soc.* 114 (1992), no. 2, 535–544.
- [7] Hiroaki Minami, “Around splitting and reaping number for partitions of  $\omega$ ”, submitted Aug 2007.
- [8] Saharon Shelah, “Vive la diffe’rence. I. Nonisomorphism of ultrapowers of countable models”. *Set theory of the continuum* (Berkeley, CA, 1989), 357–405, *Math. Sci. Res. Inst. Publ.*, 26, Springer, New York, 1992.