TITLE:

Developable of a Curve and Determinacy Relative to Osculation-Type

AUTHOR(S):

ISHIKAWA, Goo

CITATION:

ISHIKAWA, Goo. Developable of a Curve and Determinacy Relative to Osculation-Type. 数理解析研究所講究録 1991, 764: 131-135

ISSUE DATE:

1991-08

URL:

<http://hdl.handle.net/2433/82262>

RIGHT:

Developable of a Curve and Determinacy Relative to Osculation-Type

Goo ISHIKAWA

Department of Mathematics, Hokkaido University, Sapporo 060, JAPAN

Introduction

The ruled surface by tangent lines to a space curve is called the developable surface of the curve. More generally, the developable of a curve in $(n+1)$ -dimensional projective space is defined as the hypersurface "ruled" by osculating $(n-1)$ -subspaces to the curve.

Consider a C^∞ curve $\gamma : M \rightarrow \mathbf{R}P^{n+1}$, where M is a 1-dimensional manifold. We call the germ γ_p at a point $p \in M$ of finite osculation-type (or simply, of finite type) $\mathbf{a} = (a_1, a_2, \dots, a_{n+1})$ if there exist a C^∞ coordinate t of (M, p) and an affine coordinate $(x_1, \dots, x_{n+1})$ of $\mathbf{R}P^{n+1}$ centered at $\gamma(p)$ such that γ is represented by

$$x_1 = t^{a_1} + o(t^{a_1}), \quad \dots, \quad x_{n+1} = t^{a_{n+1}} + o(t^{a_{n+1}}),$$

where each a_i is a natural number and $1 \leq a_1 < \dots < a_{n+1}$.

A point $p \in M$ is called an ordinary point if γ_p is of type $(1, 2, \dots, n, n+1)$, and, otherwise, it is called a special point.

For each $p \in M$ where γ_p is of finite type and for each i , $(0 \leq i \leq n+1)$, there exists the most osculating linear subspace to γ at p in $T_{\gamma(p)}\mathbf{R}P^{n+1}$ of dimension i . We call it the osculating i -subspace and denote by $O_i(\gamma, p)$. The corresponding projective subspace of $\mathbf{R}P^{n+1}$ through p of dimension i is also denoted by $O_i(\gamma, p)$. The type of a curve therefore describes the order of tangency to each osculating subspace, and it is the simplest local projective invariant of the curve.

We can define the osculating i -bundle $O_i(\gamma)$ in the pullback $\gamma^{-1}T\mathbf{R}P^{n+1}$. The natural parametrization

$$\text{dev}(\gamma) : O_{n-1}(\gamma) \rightarrow \mathbf{R}P^{n+1}$$

defined by $(p, q) \mapsto q$, where $q \in O_{n-1}(\gamma, p) (\subset \mathbf{R}P^{n+1})$, is called also a developable of γ .

There are several results on the classification of developables of curves under the C^∞ right-left equivalence.

For a space curve γ , at each ordinary point p , the developable has cuspidal singularities along γ and $\text{dev}(\gamma)_p$ is equivalent to $(x, t) \mapsto (x, t^2, t^3)$.

Cleave [C], Gaffney-du Plessis [GP] and Shcherbak [S1] prove that, at a point p of type $(1, 2, 4)$, $\text{dev}(\gamma)_p$ is equivalent to $(x, t) \mapsto (x, t^2, xt^3)$.

Mond [M1][M2] gives C^∞ normal forms of developable of curves of type $(1, 2, 2 + r)$, $r \leq 5$, and of type $(1, 3, 4)$.

In the case of arbitrary dimension, Shcherbak, in [S1], shows the the developable of a curve of type $(2, 3, \dots, n+1, n+2)$ is equivalent to the (parametrization of) n -dimensional swallowtail, generalizing the observation of Arnol'd [A] for a curve of type $(2, 3, 4)$ based on the Legendre singularity theory.

In the connection with the study of projections of wave front sets, Shcherbak, further in [S2], gives the C^∞ normal form of the union of the developable of a curve-germ γ_p of type $(1, 2, \dots, n, n+2)$ and the osculating hyperplane $O_n(\gamma, p)$. See also [K].

We can notice that the type of a curve determines the local C^∞ class of the developable of the curve in the above mentioned cases.

Inspired with these previous results, we are led to the natural problem that whether a type of a curve-germ γ_p determines the C^∞ class of map-germ $\text{dev}(\gamma)_p$ or not.

If such determinacy for a type $\mathbf{a}$ is established once, then to have the normal form of developables of curves of type $\mathbf{a}$ is reduced to just a calculation of an example. The purpose of this paper is to announce the complete solution of this determinacy problem.

THEOREM 1. *A type $\mathbf{a}$ of a curve-germ in $\mathbf{R}P^{n+1}$ determines C^∞ class of developable if and only if $\mathbf{a}$ is one of following types:*

- (I) $_{n,r}$ $\mathbf{a} = (1, 2, \dots, n, n+r)$, $r = 1, 2, \dots$,
- (II) $_{n,i}$ $\mathbf{a} = (1, 2, \dots, i, i+2, \dots, n+1, n+2)$, $0 \leq i \leq n-1$,
- (III) $_n$ $\mathbf{a} = (3, 4, \dots, n+2, n+3)$,
- (IV) $\mathbf{a} = (3, 5)$, (V) $\mathbf{a} = (1, 3, 5)$.

Further, in this case, for any γ_p of type **a**, the map-germ $\text{dev}(\gamma)_p$ is C^∞ right left equivalent to $(x', U(x', t), U_r(x', t)) : \mathbb{R}^n, 0 \longrightarrow \mathbb{R}^{n+1}, 0$, where $(x', t) = (x_1, \dots, x_{n-1}, t)$ is a coordinate of $(\mathbb{R}^n, 0)$,

$$U(x', t) = \frac{t^{a_n}}{a_n!} + x_1 \frac{t^{a_n - a_1}}{(a_n - a_1)!} + \dots + x_{n-1} \frac{t^{a_n - a_{n-1}}}{(a_n - a_{n-1})!},$$

$r = a_{n+1} - a_n$ and

$$U_r(x', t) = \int_0^t \frac{t^r}{r!} \frac{\partial U}{\partial t} dt.$$

Notice that the developable appears as a component of the envelope of one-parameter family of osculating hyperplanes to a curve-germ γ_p . In the case $a_{n+1} - a_n > 1$, the envelope also has a component $O_n(\gamma, p)$ itself. In this case therefore it is natural to classify developables by diffeomorphisms preserving $O_n(\gamma, p)$. Then we have

THEOREM 2. A type **a** of a curve-germ in $\mathbb{R}P^{n+1}$ determines C^∞ class of envelope of osculating hyperplanes if and only if **a** is one of types $(I)_{n,r}, r = 1, 2, \dots, (II)_{n,i}$ and $(III)_{n,n} \geq 2$, in Theorem 1.

THEOREM 3. A type **a** of a curve-germ γ_p in $\mathbb{R}P^{n+1}$ determines C^∞ class of the union of developable and $O_n(\gamma, p)$ if and only if **a** is one of types $(I)_{n,r}$ and $(II)_{n,i}$ in Theorem 1.

These results unifies and generalizes the results of [C], [G-P] on $(I)_{2,2}$, the results of [A], [S1], [S2], on $(I)_{n,2}$ and $(II)_{n,0}$, and the results of [M1] [M2] on $(I)_{2,r}, (r \leq 5)$, and $(II)_{2,1}$.

The proofs of Theorems 1, 2 and 3 will be given in a forthcoming paper.

Mond's theorem

Based on Theorem 1, we reprove the following result due to Mond [M1], [M2, Corollary 0.2]:

COROLLARY. *Let $\gamma : \mathbb{R}, 0 \longrightarrow \mathbb{R}P^3$ be a curve-germ of type $(1, 2, 2 + r)$. Then $\text{dev}(\gamma) : \mathbb{R}^2, 0 \longrightarrow \mathbb{R}P^3$ is a topological embedding if r is odd, and $\text{dev}(\gamma)$ has a single curve of selfintersection if r is even.*

PROOF: By Theorem 1, $\text{dev}(\gamma)$ is C^∞ equivalent to the germ at 0 of

$$f(x, t) = (x, \frac{t^2}{2} + xt, \int_0^t \frac{s^r}{r!}(s + x)ds) : \mathbb{R}^2 \longrightarrow \mathbb{R}^3.$$

Now, assume $f(x_1, t_1) = f(x_2, t_2), (x_i, t_i) \in \mathbb{R}^2, i = 1, 2$. Then we see $x_1 = x_2, x_1 = -(1/2)(t_1 + t_2)$ and $\int_{t_1}^{t_2} s^r(s + x_1)ds = 0$. Thus, setting $\sigma = s + x_1$, we have

$$\int_{-a}^a (\sigma - x_1)^r \sigma d\sigma = 0 \quad \dots (*),$$

where $a = (1/2)(t_2 - t_1)$.

If r is odd, then the left hand side of (*) is equal to an integral from $-a$ to a with almost everywhere positive integrand. Hence we have $a = 0$. This means that $(x_1, t_1) = (x_2, t_2)$ and that f is injective.

By a similar argument, if r is even, then we have $x_1 = 0$ or $(x_1, t_1) = (x_2, t_2)$.

Since f is a finite mapping and $f|\{x = 0\} = (0, t^2/2, (r+1)\{t^{r+2}/(r+2)!\})$, we see f is an embedding in the complement of a double point curve $\{x = 0\}$.

REFERENCES

- [A] V.I. Arnol'd, *Lagrangian manifolds with singularities, asymptotic rays, and the open swallowtail*, Funct. Anal. Appl. 15-4 (1981), 235-246.
- [B-M] E. Bierstone, P.D. Milman, *Relations among analytic functions I, II*, Ann. Inst. Fourier 37-1, 37-2 (1987), 187-239, 49-77.

- [C] J.P. Cleave, *The form of the tangent developable at points of zero torsion on space curves*, Math. Proc. Camb. Phil. Soc. **88** (1980), 403–407.
- [G-P] T. Gaffney, A. du Plessis, *More on the determinacy of smooth map-germs*, Invent. Math. **66** (1982), 137–163.
- [K] M.É. Kazaryan, *Singularities of the boundary of fundamental systems, flat points of projective curves, and Schubert cells*, in “Itogi Nauki Tekh., Ser. Sorrem. Probl. Mat. (Comtemporary Problems of Mathematics) 33,” VITINI, 1988, pp. 215–232.
- [M1] D. Mond, *On the tangent developable of a space curve*, Math. Proc. Camb. Phil. Soc. **91** (1982), 351–355.
- [M2] ———, *Singularities of the tangent developable surface of a space curve*, Quart. J. Math. Oxford **40** (1989), 79–91.
- [S1] O.P. Shcherbak, *Projectively dual space curves and Legendre singularities*, Trudy Tbiliss. Univ. **232-233** (1982), 280–336.
- [S2] ———, *Wavefront and reflection groups*, Russian Math. Surveys **43-3** (1988), 149–194.