

Simple K3 singularities which are hypersurface sections
of toric singularities.

Hiroyasu TSUCHIHASHI

東北学院大学
土橋宏康

Let N be a free \mathbf{Z} -module of rank $n+1$. Let ξ^n be the set of pairs (σ, u_0) consisting of an $(n+1)$ -dimensional cone in $N_{\mathbf{R}}$ and a point u_0 in σ satisfying the following conditions (G) and (E), respectively.

(G) There exists the point $v(\sigma)$ in N^* such that σ is generated by finite elements in $\{u \in N \mid \langle v(\sigma), u \rangle = 1\}$.

(E) $\dim \Delta_{\sigma}(u_0) = n$ and $v(\sigma) \in \text{Int}(\Delta_{\sigma}(u_0))$, where $\Delta_{\sigma}(u_0)$ is the convex hull of $\{v \in \sigma^* \cap N^* \mid \langle v, u_0 \rangle = 1\}$.

Note that if an $(n+1)$ -dimensional cone σ in $N_{\mathbf{R}}$ satisfies the condition (G), then the point $v(\sigma)$ is unique and σ is strongly convex rational cone. Let (σ, u_0) be a pair in ξ^n and let $f = \sum_{v \in \Delta_{\sigma}(u_0) \cap N^*} c_v z^v + \text{higher term} \in \mathbb{C}[\sigma^* \cap N^*]$, for certain non-zero coefficients c_v . In the previous paper[2], we show that if f is non-degenerate and the hypersurface section $X = \{f = 0\}$ of $Y = \text{Spec} \mathbb{C}[\sigma^* \cap N^*]$ defined by f has an isolated singularity at $y := \text{orb}(\sigma)$, then (X, y) is a purely elliptic of $(0, n-1)$ -type singularity (for the

definition, see [3]). Especially, when $n = 3$, (X, y) is a simple K3 singularity. We also show in [2] that ξ^3 is a finite set, where ξ^n is the set of equivalence classes of pairs in $\tilde{\xi}^n$ by the following equivalence relation. $(\sigma, u_0) \sim (\sigma', u'_0)$ if and only if there exists an element g in $GL(N)$ such that $g_{\mathbb{R}}\sigma = \sigma'$ and that $g_{\mathbb{R}}(u_0) = u'_0$. In this paper, we show that ξ^n is a finite set for each integer n greater than 2.

Let $\tilde{\mathcal{F}}^n = \{ (\sigma, u_0) \in \tilde{\xi}^n \mid u_0 \in N \}$ and let \mathcal{F}^n be the set of the equivalence classes of the pairs in $\tilde{\mathcal{F}}^n$.

Proposition 1. There exists a map π from ξ^n to \mathcal{F}^n such that $\pi^{-1}(\alpha)$ is a finite set for each α in \mathcal{F}^n .

Proof. Let (σ, u_0) be in $\tilde{\xi}^n$. Then u_0 is in $N_{\mathbb{Q}}$, by the condition (E). Hence the module $N(u_0)$ generated by N and u_0 is also a free \mathbb{Z} -module of rank $n+1$. Therefore, there exists an isomorphism g from $N(u_0)$ to N . First, we show that the pair $(g_{\mathbb{R}}\sigma, g(u_0))$ is in $\tilde{\mathcal{F}}^n$.

Since $\langle v(\sigma), u_0 \rangle = 1$, $v(\sigma)$ is in ${}^t g(N^*) = N(u_0)^* = \{ v \in N^* \mid \langle v, u_0 \rangle \in \mathbb{Z} \}$. Hence ${}^t g_{\mathbb{R}}^{-1}(v(\sigma))$ is in N^* . Therefore, $g_{\mathbb{R}}\sigma$ satisfies (G) and $v(g_{\mathbb{R}}\sigma) = {}^t g_{\mathbb{R}}^{-1}(v(\sigma))$. Since $\{ v \in \sigma^* \cap N^* \mid \langle v, u_0 \rangle = 1 \} = \{ v \in \sigma^* \cap N(u_0)^* \mid \langle v, u_0 \rangle = 1 \} = {}^t g(\{ v' \in (g_{\mathbb{R}}\sigma)^* \cap N^* \mid \langle v', g(u_0) \rangle = 1 \})$, we see that

$\Delta_{g_{\mathbb{R}}\sigma}(g(u_0)) = {}^t g_{\mathbb{R}}^{-1}(\Delta_{\sigma}(u_0))$. Hence $g(u_0)$ satisfies (E).

we easily see that if $(\sigma, u_0) \sim (\sigma', u'_0)$, then $(g_{\mathbb{R}}\sigma, g(u_0)) \sim (g'_{\mathbb{R}}\sigma', g'(u'_0))$, for any isomorphisms $g : N(u_0) \simeq N$ and $g' : N(u'_0) \simeq N$. We denote by π , the map from \mathcal{E}^n to \mathcal{F}^n thus obtained. Next, we show that $\pi^{-1}(\alpha)$ is a finite set for each equivalence class α in \mathcal{F}^n .

Let β and β' be elements in $\pi^{-1}(\alpha)$, let (σ, u_0) , (σ', u'_0) and (τ, t_0) be representatives of β , β' and α , respectively. Then there exist isomorphisms $g : N(u_0) \simeq N$ and $g' : N(u'_0) \simeq N$ such that $g_{\mathbb{R}}\sigma = g'_{\mathbb{R}}\sigma' = \tau$ and that $g(u_0) = g'(u'_0) = t_0$. Assume that $g(N) = g'(N)$. Then the map $h := (g')^{-1}|_{g(N)} \circ g|_N$ is in $GL(N)$, $h_{\mathbb{R}}\sigma = \sigma'$ and $h_{\mathbb{R}}(u_0) = u'_0$. Hence $(\sigma, u_0) \sim (\sigma', u'_0)$. On the other hand, $\langle v(\sigma), g_{\mathbb{R}}^{-1}(u) \rangle = \langle {}^t g_{\mathbb{R}}^{-1}(v(\sigma)), u \rangle = \langle v(\tau), u \rangle = 1$ for primitive elements u in all 1-dimensional faces of τ . Since $g_{\mathbb{R}}^{-1}(u)$ are generators of 1-dimensional faces of σ , $g_{\mathbb{R}}^{-1}(u) \in N$, by the condition (G). Hence $g(N)$ contains the module N' generated by primitive elements in all 1-dimensional faces of τ . Since τ is an $(n+1)$ -dimensional rational cone, N' is also a free \mathbb{Z} -module of rank $n+1$. Hence N/N' is a finite group. Therefore, $\#(\pi^{-1}(\alpha)) \leq \#\{\text{subgroups of } N/N'\} < +\infty$. q.e.d.

Note that $\{u \in \text{Int}(\sigma) \cap N \mid \langle v(\sigma), u \rangle = 1\} = \{u_0\}$ for any pair (σ, u_0) in $\tilde{\mathcal{F}}^n$ (see [2, Proposition 1.8]). Hence we have an injective map from \mathcal{F}^n to $\mathcal{P}^n := \tilde{\mathcal{F}}^n / \sim$, where $\tilde{\mathcal{F}}^n$

is the set of n -dimensional integral convex polytopes P in \mathbb{R}^n with $\text{Int}(P) \cap \mathbb{Z}^n = \{0\}$ and $P \sim P'$ if and only if there exists an element g in $GL(n, \mathbb{Z})$ such that $g_{\mathbb{R}}P = P'$. Hence if \mathcal{P}^n is finite, then \mathcal{E}^n is also finite, by the above proposition.

Proposition 2. \mathcal{P}^n is a finite set.

Proof. There exists a real number L such that $\text{vol}(P) < L$ for any P in $\tilde{\mathcal{P}}^n$, by [1]. Let S be the set of simplices $\overline{0v_1v_2 \dots v_n}$ spanned by $0, v_1 = {}^t(p_{11}, 0, \dots, 0), \dots, v_j = {}^t(p_{j1}, \dots, p_{jj}, 0, \dots, 0) \dots$ and $v_n = {}^t(p_{n1}, \dots, p_{nn})$ in \mathbb{Z}^n such that $0 \leq p_{jk} < p_{jj}$ for $j = 1$ through n and for $k = 1$ through $j-1$ and that $p_{11}p_{22} \dots p_{nn} < n!L$. Clearly, S is a finite set. Let P be in $\tilde{\mathcal{P}}^n$. Then P contains n vertices u_1, u_2, \dots and u_n which are linearly independent. There exists an element g in $GL(n, \mathbb{Z})$ such that $g(u_j) = (p_{j1}, \dots, p_{jj}, 0, \dots, 0)$ ($0 \leq p_{jk} < p_{jj}$ for $k = 1$ through $j-1$). Since $\text{vol}(\overline{0u_1 \dots u_n}) \leq \text{vol}(P) < L$, $g(\overline{0u_1 \dots u_n}) \in S$. On the other hand, each point u in P is a linear combination $a_1u_1 + a_2u_2 + \dots + a_nu_n$ of u_1, u_2, \dots and u_n . If $a_j \neq 0$, then $L > \text{vol}(P) \geq \text{vol}(\overline{0u_1 \dots u_{j-1}uu_{j+1} \dots u_n}) = |a_j| \text{vol}(\overline{0u_1 \dots u_n}) = |a_j| p_{11}p_{22} \dots p_{nn} / n!$. Hence $g(P)$ is contained in the compact set $C := \{ a_1g(u_1) + a_2g(u_2) + \dots + a_n g(u_n) \mid |a_j| \leq$

$n!L/p_{11}\cdots p_{nn}$ }. Since the set of integral convex polytopes contained in C is finite, \mathcal{P}^n is also finite. q.e.d.

Thus we obtain:

Theorem 3. \mathcal{E}^n is finite.

References

- [1] D. Hensley, Lattice vertex polytopes with interior lattice points, Pacific J. of Math. vol. 105, no. 1 (1983), 183-191.
- [2] H. Tsuchihashi, A classification of toric singularities with simple K3 singularities as hypersurface sections, Suuri kaiseki kenkyujo kokyuuroku vol. 742 (1991), 111-125.
- [3] K. Watanabe, Minimal resolution of a quasi-homogeneous simple K3 singularity, in this volume.