

**Some results on Ehrhart polynomials of  
star-shaped triangulations of balls**

**Takayuki Hibi**

日比孝之 (北海道大学)  
理学部

**Department of Mathematics  
Faculty of Science  
Hokkaido University  
Kita-ku, Sapporo 060, Japan**

**In this paper we extend the work [Sta5] and [H4] on Ehrhart polynomials of convex polytopes.**

**First, we recall what the Ehrhart polynomial of a convex polytope is. Let  $\mathcal{P} \subset \mathbb{R}^N$  be an integral convex polytope, i.e., a convex polytope any of whose vertices has integer coordinates, of dimension  $d$  and  $\partial\mathcal{P}$  the boundary of  $\mathcal{P}$ . Given an integer  $n > 0$  we write  $i(\mathcal{P}, n)$  for the number of rational points  $(\alpha_1, \alpha_2, \dots, \alpha_N)$  in  $\mathcal{P}$  such that each  $n\alpha_i$  is an integer. In other words,  $i(\mathcal{P}, n) = \#(n\mathcal{P} \cap \mathbb{Z}^N)$ . Here  $n\mathcal{P} := \{n\alpha; \alpha \in \mathcal{P}\}$  and  $\#(X)$  is the cardinality of a finite set  $X$ . The systematic study of  $i(\mathcal{P}, n)$  originated in the work of Ehrhart (cf. [Ehr]), who established that the function  $i(\mathcal{P}, n)$  possesses the following fundamental properties :**

**(0.1)  $i(\mathcal{P}, n)$  is a polynomial in  $n$  of degree  $d$ .**

**(Thus  $i(\mathcal{P}, n)$  can be defined for every integer  $n$ .)**

**(0.2)  $i(\mathcal{P}, 0) = 1$ .**

**(0.3) ("loi de réciprocité")  $(-1)^d i(\mathcal{P}, -n) = \#(n(\mathcal{P} - \partial\mathcal{P}) \cap \mathbb{Z}^N)$   
for every integer  $n > 0$ .**

**We say that  $i(\mathcal{P}, n)$  is the Ehrhart polynomial of  $\mathcal{P}$ .**

We define the sequence  $\delta_0, \delta_1, \delta_2, \dots$  of integers by the formula

$$(1 - \lambda)^{d+1} \left[ 1 + \sum_{n=1}^{\infty} i(\mathcal{P}, n) \lambda^n \right] = \sum_{i=0}^{\infty} \delta_i \lambda^i. \quad (*)$$

Thus, in particular,  $\delta_0 = 1$  and  $\delta_1 = \#(\mathcal{P} \cap \mathbb{Z}^N) - (d + 1)$ . Also,  $\delta_i = 0$  for every  $i > d$  by, e.g., [Sta3, Corollary 4.3.1], and  $\delta_d = \#((\mathcal{P} - \partial\mathcal{P}) \cap \mathbb{Z}^N)$  by (0.3). Moreover, each  $\delta_i$  is non-negative [Sta1]. When  $d = N$ ,  $(\sum_{0 \leq i \leq d} \delta_i) / d!$  is equal to the volume of  $\mathcal{P}$  ([Sta3, Proposition 4.6.30]). We say that the sequence  $\delta(\mathcal{P}) := (\delta_0, \delta_1, \dots, \delta_d)$  which appears in Eq. (\*) is the  $\delta$ -vector of  $\mathcal{P}$ .

Some linear inequalities on the  $\delta$ -vector of  $\mathcal{P}$  are known, e.g.,

(1.1) ([Sta5]) If  $\delta_j \neq 0$  and  $\delta_{j+1} = \delta_{j+2} = \dots = \delta_d = 0$ , then  $\sum_{0 \leq \ell \leq i} \delta_\ell \leq \sum_{0 \leq \ell \leq i} \delta_{j-\ell}$  for every  $0 \leq i \leq [j/2]$ .

(1.2) ([H4]) When  $(\mathcal{P} - \partial\mathcal{P}) \cap \mathbb{Z}^N$  is non-empty, then  $\delta_1 \leq \delta_i$  for every  $1 \leq i < d$ .

See also [Sta6], [B-M], [H1], [H2] and [H3] for further results on Ehrhart polynomials of convex polytopes.

On the other hand, let  $\Gamma$  be an integral polyhedral complex in  $\mathbb{R}^N$ , i.e., a finite set of integral convex polytopes in  $\mathbb{R}^N$  such that (a) if  $\mathcal{P} \in \Gamma$  and  $\mathcal{Q}$  is a face of  $\mathcal{P}$ , then  $\mathcal{Q} \in \Gamma$ , and (b) if  $\mathcal{P}, \mathcal{P}' \in \Gamma$ , then  $\mathcal{P} \cap \mathcal{P}'$  is a face of  $\mathcal{P}$  and of  $\mathcal{P}'$ . We write  $|\Gamma|$  for the underlying space  $\cup_{\mathcal{P} \in \Gamma} \mathcal{P}$  ( $\subset \mathbb{R}^N$ ) of  $\Gamma$ , and let  $\partial|\Gamma|$  be the boundary of  $|\Gamma|$  (in the usual topological sense with respect to the relative topology on  $|\Gamma|$  inherited from the standard topology on  $\mathbb{R}^N$ ). We call  $d := \max\{\dim \mathcal{P}; \mathcal{P} \in \Gamma\}$  the dimension of  $\Gamma$ . In analogy with  $i(\mathcal{P}, n)$ , we define for  $n > 0$   $i(\Gamma, n)$  to be the number of rational points  $(\alpha_1, \alpha_2, \dots, \alpha_N)$  in  $|\Gamma|$  for which each  $n\alpha_j \in \mathbb{Z}$ . Thus, thanks to (0.3), we easily see

$$i(\Gamma, n) = \sum_{\mathcal{P} \in \Gamma} (-1)^{\dim \mathcal{P}} i(\mathcal{P}, -n).$$

Hence  $i(\Gamma, n)$  is a polynomial in  $n$  of degree  $d$ , however,  $i(\Gamma, 0) = \chi(\Gamma)$ , the Euler characteristic of  $\Gamma$ .

Now, suppose that  $\Gamma$  is an integral polyhedral complex in  $\mathbb{R}^N$  of dimension  $d$  such that the underlying space  $|\Gamma|$  is homeomorphic to the  $d$ -ball. Then  $\chi(\Gamma) = 1$ . Hence the  $\delta$ -vector  $\delta(\Gamma) = (\delta_0, \delta_1, \dots, \delta_d)$  of  $\Gamma$  can be defined by replacing  $i(\mathcal{P}, n)$  with  $i(\Gamma, n)$  in Eq. (\*). Thus  $\delta_0 = 1$ ,  $\delta_1 = \#(|\Gamma| \cap \mathbb{Z}^N) - (d + 1)$  and  $\delta_d = \#((|\Gamma| - \partial|\Gamma|) \cap \mathbb{Z}^N)$ . Here  $\partial|\Gamma|$  is the boundary, which is homeomorphic to the  $(d-1)$ -sphere, of  $|\Gamma|$ . Also, each  $\delta_i$  is non-negative [Sta4].

We are now in the position to state our main result in this paper.

**THEOREM.** Let  $\Gamma$  be an integral polyhedral complex in  $\mathbb{R}^N$  of dimension  $d$  such that the underlying space  $|\Gamma|$  is homeomorphic to the  $d$ -ball. Suppose that  $(|\Gamma| - \partial|\Gamma|) \cap \mathbb{Z}^N$  is non-empty, i.e.,  $\delta_d \neq 0$ , and that  $|\Gamma|$  is star-shaped relative to some  $\alpha \in (|\Gamma| - \partial|\Gamma|) \cap \mathbb{Z}^N$ , i.e., for each  $\beta$  in  $|\Gamma|$ , the open line segment  $\{(1-t)\alpha + t\beta; 0 < t < 1\}$  from  $\alpha$  to  $\beta$  lies in  $|\Gamma| - \partial|\Gamma|$ . Let  $\delta(\Gamma) = (\delta_0, \delta_1, \dots, \delta_d)$  be the  $\delta$ -vector of  $\Gamma$ .

(2.1) We have the lower bound inequality

$$\delta_1 \leq \delta_i$$

for every  $1 \leq i < d$ .

(2.2) Moreover, the linear inequality

$$\delta_0 + \delta_1 + \dots + \delta_i \leq \delta_d + \delta_{d-1} + \dots + \delta_{d-i}$$

holds for every  $0 \leq i \leq [d/2]$ .

The proof of (2.1) is based on the same combinatorial technique as in [H4]. On the other hand, in [Sta5], the canonical ideal  $\Omega(A(\mathcal{P}))$  of a Cohen-Macaulay integral domain  $A(\mathcal{P})$  associated with a convex polytope  $\mathcal{P}$  plays an essential role for the proof of (1.1).

More generally, in [Sta<sub>4</sub>, p.202], given an integral polyhedral complex  $\Gamma$  in  $\mathbb{R}^N$ , a noetherian graded ring  $A(\Gamma)$  is defined as follows. Let  $X_1, \dots, X_N$  and  $T$  be indeterminates over a field  $k$ . A basis of  $A(\Gamma)$  as a vector space over  $k$  consists of 1 together with all monomials  $X^{\alpha}T^n = X_1^{\alpha_1} \dots X_N^{\alpha_N}T^n$ , where  $n > 0$  is an integer and  $\alpha = (\alpha_1, \dots, \alpha_N) \in (n|\Gamma| \cap \mathbb{Z}^N)$ . In  $A(\Gamma)$ , multiplication of two monomials  $X^{\alpha}T^n$  and  $X^{\beta}T^m$  is defined by (i)  $(X^{\alpha}T^n)(X^{\beta}T^m) = X^{\alpha+\beta}T^{n+m}$  if  $\alpha \in n\mathcal{P}$  and  $\beta \in m\mathcal{P}$  for some  $\mathcal{P} \in \Gamma$ , and (ii)  $(X^{\alpha}T^n)(X^{\beta}T^m) = 0$  otherwise. (Note that  $A(\Gamma)$  is never an integral domain unless  $|\Gamma| = \mathcal{P}$  for some  $\mathcal{P} \in \Gamma$ .) We define a grading on  $A(\Gamma)$  by setting  $\deg(X^{\alpha}T^n) = n$ . Then the Hilbert function  $H(A(\Gamma), n)$  of  $A(\Gamma)$  is equal to  $i(\Gamma, n)$  for every  $n > 0$ .

Thanks to [Sta<sub>4</sub>, Lemma 4.6], if  $|\Gamma|$  is homeomorphic to the  $d$ -ball, then the algebra  $A(\Gamma)$  is Cohen-Macaulay. Moreover, the canonical ideal  $\Omega(A(\Gamma))$  of  $A(\Gamma)$  can be expressed easily by virtue of Hochster's theorem ([Sta<sub>2</sub>, p.81]). The key lemma for our proof of (2.2) is the existence of an integral polyhedral complex  $\Gamma'$  in  $\mathbb{R}^N$  with  $|\Gamma| = |\Gamma'|$  such that the canonical ideal  $\Omega(A(\Gamma'))$  possesses a certain non-zero divisor on  $A(\Gamma')$  which is required in, e.g., [H<sub>5</sub>, Proposition (1.3)].

**EXAMPLE.** Let  $\mathcal{P} \subset \mathbb{R}^3$  be the simplex with vertices  $A = (1,0,0)$ ,  $B = (0,1,0)$ ,  $C = (0,0,1)$  and  $D = (-1,-1,-1)$ . Also, let  $\mathcal{Q} \subset \mathbb{R}^3$  be the simplex whose vertices are  $A$ ,  $B$ ,  $C$  and  $E = (1,1,0)$ . We define  $\Gamma$  to be the integral polyhedral complex in  $\mathbb{R}^3$  of dimension 3 which consists of  $\mathcal{P}$ ,  $\mathcal{Q}$ , all faces of  $\mathcal{P}$  and all faces of  $\mathcal{Q}$ . Then the underlying space  $|\Gamma|$  is homeomorphic to the 3-ball, and  $(|\Gamma| - \partial|\Gamma|) \cap \mathbb{Z}^3 = \{(0,0,0)\}$ . However,  $|\Gamma|$  is not star-shaped relative to the origin of  $\mathbb{R}^3$ . We have  $\delta(\Gamma) = (1,2,1,1)$ , which fails to satisfy the inequality in (2.1) for  $i = 2$  and that in (2.2) for  $i = 1$ .

The author is grateful to Professor Richard P. Stanley for valuable suggestion on the original version of the present paper.

## REFERENCES

- [B-M] U. Betke and P. McMullen, Lattice points in lattice polytopes, *Monatsh. Math.* **99** (1985), 253-265.
- [Ehr] E. Ehrhart, "Polynômes arithmétiques et Méthode des Polyèdres en Combinatoire," Birkhäuser, Basel and Stuttgart, 1977.
- [H<sub>1</sub>] T. Hibi, Some results on Ehrhart polynomials of convex polytopes, *Discrete Math.* **83** (1990), 119-121.
- [H<sub>2</sub>] ———, Dual polytopes of rational convex polytopes, *Combinatorica*, in press.
- [H<sub>3</sub>] ———, A combinatorial self-reciprocity theorem for Ehrhart quasi-polynomials of rational convex polytopes, *European J. Combin.*, to appear.
- [H<sub>4</sub>] ———, A lower bound theorem for Ehrhart polynomials of convex polytopes, *Advances in Math.*, to appear.
- [H<sub>5</sub>] ———, Face number inequalities for matroid complexes and Cohen-Macaulay types of Stanley-Reisner rings of distributive lattices, *Pacific J. Math.*, in press.
- [Sta<sub>1</sub>] R. Stanley, Decompositions of rational convex polytopes, *Annals of Discrete Math.* **6** (1980), 333-342.
- [Sta<sub>2</sub>] ———, "Combinatorics and Commutative Algebra," *Progress in Math.*, Vol. 41, Birkhäuser, Boston/Basel/Stuttgart, 1983.
- [Sta<sub>3</sub>] ———, "Enumerative Combinatorics, Volume I," Wadsworth & Brooks/Cole, Monterey, Calif., 1986.
- [Sta<sub>4</sub>] ———, Generalized h-vectors, intersection cohomology of toric varieties, and related results, in "Commutative Algebra and Combinatorics" (M. Nagata and H. Matsumura, eds.), *Advanced Studies in Pure Math.*, Vol 11, North-Holland, Amsterdam, 1987, pp. 187-213.
- [Sta<sub>5</sub>] ———, On the Hilbert function of a graded Cohen-Macaulay domain, *J. of Pure and Appl. Algebra* **73** (1991), 307-314.
- [Sta<sub>6</sub>] ———, A monotonicity property of h-vectors and h\*-vectors, preprint (March, 1991).