

Title	A Fixed Point Theorem for Noncommutative Families of Nonexpansive Mappings in Banach spaces(Nonlinear Analysis and Convex Analysis)
Author(s)	LAU, Anthony To-Ming; TAKAHASHI, Wataru
Citation	数理解析研究所講究録 (1995), 897: 133-142
Issue Date	1995-02
URL	<a href="http://hdl.handle.net/2433/84466">http://hdl.handle.net/2433/84466</a>
Right	
Type	Departmental Bulletin Paper
Textversion	publisher

# A Fixed Point Theorem for Noncommutative Families of Nonexpansive Mappings in Banach spaces

by

Anthony To-Ming LAU and Wataru TAKAHASHI  
(高橋 渉)

Department of Mathematical Sciences,  
University of Alberta,  
Edmonton, Alberta, Canada T6G-2G1

and

Department of Information Sciences,  
Tokyo Institute of Technology,  
Oh-Okayama, Meguro-ku, Tokyo 152, Japan

## Abstract

Let  $C$  be a nonempty weakly compact convex subset of a Banach space which has normal structure and let  $S$  be a semitopological semigroup such that  $RUC(S)$  has a left invariant mean. Then we prove a fixed point theorem for a continuous representation of  $S$  as nonexpansive mappings on  $C$ .

---

1991 Mathematics Subject Classification. Primary 47H10.

Key words and phrases. Fixed point, nonexpansive mapping, invariant mean.

## 1 Introduction.

Let  $S$  be a semitopological semigroup, i.e.,  $S$  is a semigroup with a Hausdorff topology such that for each  $a \in S$ , the mappings  $s \rightarrow sa$  and  $s \rightarrow as$  from  $S$  into  $S$  are continuous and let  $RUC(S)$  be the space of bounded right uniformly continuous functions on  $S$ . Let  $C$  be a nonempty subset of a Banach space and let  $\mathcal{S} = \{T_t : t \in S\}$  be a family of self-maps of  $C$ .  $\mathcal{S}$  is said to be a continuous representation of  $S$  as nonexpansive mappings on  $C$  if the following conditions are satisfied :

- (1)  $T_{st}x = T_sT_tx$  for all  $t, s \in S$  and  $x \in C$ ;
- (2) for each  $x \in C$ , the mapping  $s \rightarrow T_sx$  from  $S$  into  $C$  is continuous.

Let  $F(\mathcal{S})$  denote the set of common fixed points of  $T_s$ ,  $s \in S$ . Fixed point theorems for noncommutative families of nonexpansive mappings on  $C$  have been investigated by several authors ; see, for example, Bartoszek[1], Holmes-Lau[2,3], Lau[4,5,6], Lau-Takahashi[7,8], Lim[9,10], Mitchell[11,12], Takahashi[13,14,15,16], Takahashi-Jeong[17] and others. Among these, Lim[9] proved that if  $S$  is left reversible (i.e., any two closed right ideals in  $S$  have non-void intersection) and  $C$  is weakly compact, convex, and has normal structure, then  $\mathcal{S}$  has a common fixed point in  $C$ .

In this paper, we prove a fixed point theorem for a continuous representation of  $S$  as nonexpansive mappings on  $C$  in the case of which  $RUC(S)$

has a left invariant mean and  $C$  is weakly compact, convex, and has normal structure. It is well known that left reversibility and existence of a left invariant mean on  $RUC(S)$  do not imply each other.

## 2 Fixed point theorem.

Let  $S$  be a set and  $m(S)$  be the Banach space of all bounded real-valued functions on  $S$  with the supremum norm. Let  $X$  be a subspace of  $m(S)$  containing constants. Then  $\mu \in X^*$  is called a mean on  $X$  if  $\|\mu\| = \mu(1) = 1$ . Let  $\mu \in X^*$  be a mean on  $X$  and  $f \in X$ . Then we denote by  $\mu(f)$  the value of  $\mu$  at the function  $f$ . According to time and circumstances, we write  $\mu_t(f(t))$  the value  $\mu(f)$ . As is well known,  $\mu \in X^*$  is a mean on  $X$  if and only if

$$\inf_{s \in S} f(s) \leq \mu(f) \leq \sup_{s \in S} f(s)$$

for every  $f \in X$ . If  $S$  is a semigroup,  $a \in S$ , and  $f \in m(S)$ , define  $(\ell_a f)(t) = f(at)$  and  $(r_a f)(t) = f(ta)$ ,  $t \in S$ . If  $\ell_a(X) \subseteq X$  for all  $a \in S$ , then a mean  $\mu$  on  $X$  is left invariant if  $\mu(\ell_a f) = \mu(f)$  for all  $a \in S$  and  $f \in X$ . Let  $S$  be a semitopological semigroup. Let  $C(S)$  be the Banach space of bounded continuous real-valued functions on  $S$ . Let  $RUC(S)$  denote the space of bounded right uniformly continuous functions on  $S$ , i.e., all  $f \in C(S)$  such that the mapping  $s \rightarrow r_s f$  of  $S$  into  $C(S)$  is continuous. Then  $RUC(S)$  is a closed subalgebra of  $C(S)$  containing constants and invariant under left and right translations (see [12] for details). A closed convex subset  $C$  of a Banach

space is said to have normal structure if for each closed bounded convex subset  $K$  of  $C$ , which contains at least two points, there exists an element of  $K$  which is not a diametral point of  $K$ . It is well known that a closed convex subset of a uniformly convex Banach space has normal structure and a compact convex subset of a Banach space has normal structure. Lim[9] also proved the following.

Lemma[9]. A closed convex subset  $C$  of a Banach space has normal structure if and only if it does not contain a sequence  $\{x_n\}$  such that for some  $c > 0$ ,  $\|x_n - x_m\| \leq c$ ,  $\|x_{n+1} - \bar{x}_n\| \geq c - \frac{1}{n^2}$  for all  $n \geq 1$ ,  $m \geq 1$ , where  $\bar{x}_n = \frac{1}{n} \sum_{i=1}^n x_i$ .

Now we can prove a fixed point theorem for noncommutative families of nonexpansive mappings in Banach spaces.

*Theorem.* Let  $S$  be a semitopological semigroup, let  $D$  be a weakly compact subset of a Banach space  $B$  which has normal structure and let  $\mathcal{S} = \{T_s : s \in S\}$  be a continuous representation of  $S$  as nonexpansive mappings on  $D$ . Suppose  $RUC(S)$  has a left invariant mean. Then  $\mathcal{S}$  has a common fixed point in  $D$ .

*Proof.* We first prove that for any  $x \in D$  and  $y \in B$ , a function  $h$  defined by  $h(t) = \|T_t x - y\|$  for all  $t \in S$  is in  $RUC(S)$ . In fact, we have, for

$s, u \in S,$

$$\begin{aligned} \|r_s h - r_u h\| &= \sup_{t \in S} |(r_s h)(t) - (r_u h)(t)| = \sup_{t \in S} |h(ts) - h(tu)| \\ &= \sup_{t \in S} |\|T_{ts}x - y\| - \|T_{tu}x - y\|| \leq \sup_{t \in S} \|T_{ts}x - T_{tu}x\| \\ &\leq \|T_s x - T_u x\|. \end{aligned}$$

Let

$$E = \{K \subset D : K \text{ is nonempty, closed, convex, and } T_s\text{-invariant}\}.$$

Then by Zorn's Lemma, there exists a minimal element  $C$  of  $E$ . Let  $\delta(C) > 0$  and let  $\mu$  be a left invariant mean. Then, for any  $x \in C$ ,

$$A_x = \{z \in C : \mu_t \|T_t x - z\| = \min_{y \in C} \mu_t \|T_t x - y\|\}$$

is nonempty, closed, convex, and  $T_s$ -invariant (see [8,13] for details). So, we have  $A_x = C$  from minimality of  $C$ . Since  $\mu$  is a mean, there exists a net of finite means  $\lambda_\alpha$  such that  $\lambda_\alpha \xrightarrow{w^*} \mu$ . Let  $x_0 \in C$ ,  $\epsilon > 0$ , and  $x_1, x_2, \dots, x_n \in C$ . Since  $A_{x_0} = C$ , there exists  $\alpha_0$  such that

$$(\mu_{\alpha_0})_t \|T_t x_0 - x_i\| \leq r + \epsilon, \quad \forall i = 1, 2, \dots, n,$$

where  $r = \min_{y \in C} \mu_t \|T_t x_0 - y\|$ . That is, there exists  $z = \sum_{j=1}^{n_{\alpha_0}} \lambda_j T_{s_j} x_0$  with  $\lambda_1, \dots,$

$\lambda_{n_{\alpha_0}} \geq 0$  and  $\sum_{j=1}^{n_{\alpha_0}} \lambda_j = 1$  such that

$$\|z - x_i\| \leq \sum_{j=1}^{n_{\alpha_0}} \lambda_j \|T_{s_j} x_0 - x_i\| \leq r + \epsilon, \quad \forall i = 1, 2, \dots, n. \quad (1)$$

Let  $C_{y,\epsilon} = \{z \in C : \|z - y\| \leq r + \epsilon\}$  for each  $y \in C$ . Then by (1),

$$\{C_{y,\epsilon} : y \in C\}$$

has finite intersection property. Since  $C$  is weakly compact, there is  $z_0 \in C$  such that  $\|z_0 - y\| \leq r + \epsilon$  for every  $y \in C$ . Since  $\{T_t x_0\} \subset C$ , we have

$$\sup_{t \in S} \|z_0 - T_t x_0\| \leq \sup_{y \in C} \|z_0 - y\| \leq r + \epsilon. \text{ Since}$$

$$r = \mu_t \|T_t x_0 - z_0\| \leq \sup_t \|T_t x_0 - z_0\| \leq \sup_{y \in C} \|z_0 - y\| \leq r + \epsilon$$

and

$$r = \mu_t \|T_t x_0 - x\| \leq \sup_t \|T_t x_0 - x\| \leq \sup_{y \in C} \|y - x\|, \quad \forall x \in C,$$

we have

$$r \leq \inf_{x \in C} \sup_t \|T_t x_0 - x\| \leq \inf_{x \in C} \sup_{y \in C} \|y - x\| \leq r + \epsilon.$$

Since  $\epsilon > 0$  is arbitrary, we have

$$r = \mu_t \|T_t x_0 - x\| = \inf_{z \in C} \sup_{y \in C} \|y - z\|, \quad \forall x \in C. \quad (2)$$

Since  $x_0 \in C$  is arbitrary, for any  $x, z \in C$ , we have

$$r = \mu_t \|T_t x - z\| = \inf_{u \in C} \sup_{t \in S} \|T_t x - u\| = \inf_{u \in C} \sup_{y \in C} \|y - u\|.$$

So, let

$$A_0 = \{z \in C : \sup_{t \in S} \|z - T_t x\| \leq r, \quad \forall x \in C\}.$$

By (2), since there exists  $z_0 \in C$  such that

$$\sup_{y \in C} \|y - z_0\| = r,$$

we have that  $A_0$  is nonempty. Let  $z_0 \in A_0$  and  $s \in S$ . Then putting

$$A_s = \{z \in C : \sup_{t \in S} \|T_{st}x - z\| \leq r, \forall x \in C\},$$

we have  $z_0, T_s z_0 \in A_s$ . Further, for any  $x \in C$ ,

$$\begin{aligned} r = \mu_t \|T_t x - z_0\| &= \mu_t \|T_{st} x - z_0\| \leq \sup_{t \in S} \|T_{st} x - z_0\| \\ &\leq \sup_{t \in S} \|T_t x - z_0\| \leq r. \end{aligned}$$

and

$$\begin{aligned} r = \mu_t \|T_t x - T_s z_0\| &= \mu_t \|T_{st} x - T_s z_0\| \leq \sup_{t \in S} \|T_{st} x - T_s z_0\| \\ &\leq \sup_{t \in S} \|T_t x - z_0\| \leq r. \end{aligned}$$

For using Lim's Lemma, fix  $z_0 \in A_0$ . Then since  $r = \mu_t \|T_t z_0 - z_0\|$ , there exists  $s_1 \in S$  such that  $\|T_{s_1} z_0 - z_0\| \geq r - 1$ . Since  $z_0, T_{s_1} z_0 \in A_{s_1}$  and  $A_{s_1}$  is convex,

$$\bar{x}_2 = \frac{1}{2} z_0 + \frac{1}{2} T_{s_1} z_0 \in A_{s_1}.$$

Let  $x_1 = z_0$  and  $x_2 = T_{s_1} z_0$ . Since  $r = \mu_t \|T_t z_0 - \bar{x}_2\| = \mu_t \|T_{s_1 t} z_0 - \bar{x}_2\|$ , there exists  $s_2 \in S$  such that  $\|T_{s_1 s_2} z_0 - \bar{x}_2\| \geq r - \frac{1}{2^2}$ . So, let  $x_3 = T_{s_1 s_2} z_0$ . Then, we have

$$\|x_1 - x_2\| = \|z_0 - T_{s_1} z_0\| \leq \sup_{t \in S} \|z_0 - T_t z_0\| = r,$$



$$\|x_2 - x_3\| = \|T_{s_1}z_0 - T_{s_1s_2}z_0\| \leq \|z_0 - T_{s_1}z_0\| \leq r,$$

and

$$\|x_3 - x_1\| = \|T_{s_1s_2}z_0 - z_0\| \leq \sup_{t \in S} \|T_t z_0 - z_0\| = r.$$

Similarly, let

$$\bar{x}_3 = \frac{1}{3}x_1 + \frac{1}{3}x_2 + \frac{1}{3}x_3.$$

Then,  $r = \mu_t \|T_t z_0 - \bar{x}_3\| = \mu_t \|T_{s_1s_2t}z_0 - \bar{x}_3\|$ , there exists  $s_3 \in S$  such that  $\|T_{s_1s_2s_3}z_0 - \bar{x}_3\| \geq r - \frac{1}{3^2}$ . So, let  $x_4 = T_{s_1s_2s_3}z_0$ . Then, we have

$$\|x_4 - x_1\| = \|T_{s_1s_2s_3}z_0 - z_0\| \leq \sup_{t \in S} \|T_t z_0 - z_0\| = r,$$

$$\|x_4 - x_2\| = \|T_{s_1s_2s_3}z_0 - T_{s_1}z_0\| \leq \|T_{s_2s_3}z_0 - z_0\| \leq \sup_{t \in S} \|T_t z_0 - z_0\| = r,$$

and

$$\|x_4 - x_3\| = \|T_{s_1s_2s_3}z_0 - T_{s_1s_2}z_0\| \leq \|T_{s_3}z_0 - z_0\| \leq \sup_{t \in S} \|T_t z_0 - z_0\| = r.$$

By mathematical induction, let  $x_5 = T_{s_1s_2s_3s_4}z_0, x_6 = T_{s_1s_2s_3s_4s_5}z_0, \dots$ . Then we have

$$\|x_n - x_m\| \leq r, \quad \forall n, m \quad \text{and} \quad \|x_{n+1} - \bar{x}_n\| \geq r - \frac{1}{n^2}.$$

Using Lim's Lemma,  $C$  has not normal structure. This is a contradiction.

## References

- [1 ] W.Bartoszek, Nonexpansive actions of topological semigroups on strictly convex Banach space and fixed points, Proc.Amer.Math.Soc., 104(1988), 809-811.
- [2 ] R.H.Holmes and A.T.Lau, Asymptotically non-expansive actions of topological semigroups and fixed points, Bull. London Math. Soc., 3(1971), 343-347.
- [3 ] R.H.Holmes and A.T.Lau, Nonexpansive actions of topological semigroups and fixed points, J. London Math. Soc., 5(1972), 330-336.
- [4 ] A.T.Lau, Invariant means on almost periodic functions and fixed point properties, Rocky Mountain J. Math., 3(1973), 69-76.
- [5 ] A.T.Lau, Some fixed point theorems and their applications to  $W^*$ -algebras, in Fixed Point Theory and Its Applications (S. Swaminathan Ed.), Academic Press, Orlando, FL, (1976), 121-129.
- [6 ] A.T.Lau, Amenability and fixed point property for semigroup of non-expansive mappings, in Fixed Point Theory and Applications (M.A.Théra and J.B.Baillon Eds.), Pitman Research Notes in Mathematics Series #252 (1991), 303-313.
- [7 ] A.T.Lau and W.Takahashi, Weak convergence and non-linear ergodic theorems for reversible semigroups of nonexpansive mappings, Pacific J. Math., 126(1987), 277-294.
- [8 ] A.T.Lau and W.Takahashi, Invariant means and semigroups of nonexpansive mappings on uniformly convex Banach spaces, J. Math. Anal. Appl., 153(1990), 497-505.

- [9 ] T.C.Lim, A fixed point theorem for families of nonexpansive mappings, Pacific J. Math., 53(1974), 484-493.
- [10 ] T.C.Lim, Asymptotic centers and nonexpansive mappings in conjugate Banach spaces, Pacific J. Math., 90(1980), 135-143.
- [11 ] T.Mitchell, Fixed points of reversible semigroups of non-expansive mappings, Kōdai Math. Sem. Rep., 2(1970), 322-323.
- [12 ] T.Mitchell, Topological semigroups and fixed points, Illinois J. Math., 14(1970), 630-641.
- [13 ] W.Takahashi, Fixed point theorem for amenable semigroups of non-expansive mappings, Kōdai Math. Sem. Rep., 21(1969), 383-386.
- [14 ] W.Takahashi, A nonlinear ergodic theorem for an amenable semigroup of nonexpansive mappings in a Hilbert space, Proc. Amer. Math. Soc., 81(1981), 253-256.
- [15 ] W.Takahashi, Fixed point theorems for families of nonexpansive mappings on unbounded sets, J. Math. Soc. Japan, 36(1984), 543-553.
- [16 ] W.Takahashi, Fixed point theorem and nonlinear ergodic theorem for nonexpansive semigroups without convexity, Can. J. Math., 35(1992), 1-8.
- [17 ] W.Takahashi and D.H.Jeong, Fixed point theorem for nonexpansive semigroups on Banach space, Proc. Amer. Math. Soc., 122(1994), 1175-1179.