TITLE:

Synthetic Application of Intramolecular
Cyanoboration on the Basis of Removal and
Conversion of a Tethering Group by
Palladium-Catalyzed Retro-Allylation

AUTHOR(S):

Ohmura, Toshimichi; Awano, Tomotsugu;
Suginome, Michinori; Yorimitsu, Hideki; Oshima,
Koichiro

CITATION:

Ohmura, Toshimichi ...[et al]. Synthetic Application of Intramolecular Cyanoboration on the Basis of Removal and Conversion of a Tethering Group by Palladium-Catalyzed Retro-Allylation. *Synlett* 2008, 2008(3): 423-427: U10107ST.

ISSUE DATE:

2008-02-12

URL:

<http://hdl.handle.net/2433/88971>

RIGHT:

© 2008 Georg Thieme Verlag Stuttgart New York; This is not the published version. Please cite only the published version.; この論文は出版社版ではありません。引用の際には出版社版をご確認ご利用ください。

Synthetic Application of Intramolecular Cyanoboration on the Basis of Removal and Conversion of a Tethering Group by Palladium-Catalyzed Retro-allylation

Toshimichi Ohmura,^a Tomotsugu Awano,^a Michinori Suginome,^{a*} Hideki Yorimitsu,^{b*} Koichiro Oshima^{b*}

^a Department of Synthetic Chemistry and Biological Chemistry, Graduate School of Engineering, Kyoto University, Katsura, Nishikyo-ku, Kyoto 615-8510, Japan

^b Department of Material Chemistry, Graduate School of Engineering, Kyoto University, Katsura, Nishikyo-ku, Kyoto 615-8510, Japan
Fax: +81-75-383-2722 (Suginome); +81-75-383-2438 (Yorimitsu, Oshima)

E-mail: suginome@sbchem.kyoto-u.ac.jp, yori@orgrxn.mbox.media.kyoto-u.ac.jp, oshima@orgrxn.mbox.media.kyoto-u.ac.jp

Received: The date will be inserted once the manuscript is accepted.

Abstract: A new synthetic strategy, involving utilization of a tethered intramolecular reaction with a removable tether, was demonstrated by the intramolecular cyanoboration/retro-allylation sequence.

Key words: alkenes, arylation, nitriles, palladium, retro-allylation

Silicon- and boron-tethered intramolecular reactions are attractive methods for the synthesis of organic molecules that are not easily accessible directly via the corresponding intermolecular reactions.^{1,2,3} Attractive features of the intramolecular reaction involve high reaction efficiency, regioselectivity, and stereoselectivity. On the other hand, a major drawback of the intramolecular reactions is the need for the tethering groups, which significantly limit the variation of the substrate scope. It seems to be highly desirable to remove and convert the tethering group into other functional groups after the intramolecular cyclization step.

We expected that the removal and conversion of the tethering group would be possible by palladium-catalyzed retro-allylation, in which allylic C–C bonds in homoallylic alcohols are cleaved and converted into other C–C bonds.⁴ Because many silicon- and boron-tethered reactions are designed for homopropargylic alcohol-derived substrates,^{3g,5} the retro-allylation would work well. Herein we demonstrate the strategy in palladium-catalyzed intramolecular cyanoboration, which allows the addition of boryl and cyano groups to alkynes.^{6,7} The Suzuki-Miyaura coupling of the cyanoboration products **II** followed by retro-allylative coupling allows the synthesis of highly substituted α,β -unsaturated nitriles **IV**, in which no tethering group is left (Scheme 1).

Scheme 1 A Synthetic Application of Intramolecular Cyanoboration Based on Removal of Tethering Unit

We first carried out Suzuki-Miyaura coupling of borylalkene **1**, which was prepared via palladium-catalyzed intramolecular cyanoboration of cyanoboryl homopropargyl ether. The coupling reactions with aryl bromides took place under the Fu's conditions,⁸ giving cyano-substituted homoallylic alcohols **2a-c** in high yields (Equation 1).

Equation 1

We then examined palladium-catalyzed retro-allylative coupling of **2** with aryl bromides **3**. Reaction of cyano-substituted homoallylic alcohol **2a** with 4-bromotoluene (**3a**) was carried out in the presence of Pd(OAc)₂ (5 mol %), PCy₃ (10 mol %), and Cs₂CO₃ (1.2 equiv.) (entry 1 in Table 1).⁴ Retro-allylative coupling took place smoothly by heating the reaction mixture at 110 °C, giving alkene **4a** in 90% yield after 3 h. It should be noted that the reaction proceeded efficiently even for the homoallylic alcohol bearing the trisubstituted C–C

double bond, because the retro-allylation has been found difficult with internal alkenes.⁴ It is also noted that the geometry of the double bond isomerized from *Z* to *E* (*E*:*Z* = 85:15) in the course of the reaction, indicating that the reaction proceeded via a π -allyl palladium intermediate (see below). A small amount of protonated product **5a** (9%) was formed along with **4a**.

As reported recently,^{4b} PCy₃ was found to be the most effective ligand for the reaction of **2a** with **3a** (entry 1). Slower reaction rate or formation of a significant amount of **5a** was observed with PCy₂Ph, PCyPh₂, and PPh₃ (entries 2-4). The reaction was also catalyzed well by electron-donating P(4-MeOC₆H₄)₃, whereas the reaction rate was reduced by the use of electron-deficient P(4-CF₃C₆H₄)₃ (entries 5 and 6). It is interesting to note that phosphite ligands tended to promote protonative retro-allylation (entries 7 and 8). Alkene **5a** was selectively obtained by a palladium catalyst bearing P(OPh)₃ (entry 8). The protonative retro-allylation took place with a Pd/P(OPh)₃ catalyst even in the absence of aryl halides, giving **5a** in 89% yield (entry 9).

Table 1 Screening of Ligand in Palladium-Catalyzed Retro-Allylative Coupling of **2a** with **3a**^a

entry	ligand	time (h)	yield [%] ^b , (<i>E</i> : <i>Z</i>) ^c	
			4a	5a
1	PCy ₃	3	90 (77) ^d (85:15)	9 (na)
2	PCy ₂ Ph	18	31 (84:16)	37 (88:12)
3	PCyPh ₂	18	74 (84:16)	23 (87:13)
4	PPh ₃	18	57 (na)	7 (na)
5	P(4-MeOC ₆ H ₄) ₃	6	87 (75:25)	8 (na)
6	P(4-CF ₃ C ₆ H ₄) ₃	18	78 (85:15)	15 (90:10)
7	P(Oi-Pr) ₃	6	31 (76:24)	52 (91:9)
8	P(OPh) ₃	2	0	85 (88:12)
9 ^e	P(OPh) ₃	2	-	(89) ^d (88:12)

^a Pd(OAc)₂ (0.010 mmol), ligand (0.020 mmol), Cs₂CO₃ (0.24 mmol), **2a** (0.20 mmol), and **3a** (0.24 mmol) were stirred at 110 °C unless otherwise noted.

^b GC yield.

^c Determined by ¹H NMR of crude mixture.

^d Isolated yield.

^e Carried out in the absence of **3a**.

When bulky aryl bromide was used, in the coupling reaction, we observed different effects on the phosphine ligands on reaction selectivity. Thus, retro-allylative coupling of **2a** with sterically hindered 2,6-dimethylbromobenzene (**3b**) afforded **4b** in only 19% yield along with major formation of **5a** under the conditions using PCy₃ (entry 1 in Table 2). Improvement of the reaction yield was achieved by the use of Pd/PCyPh₂ or PPh₃ catalysts, with which **4b** was obtained in good yields (entries 3 and 4).

Table 2 Screening of Ligand in Palladium-Catalyzed Retro-Allylative Coupling of **2a** with Sterically Hindered **3b**^a

entry	ligand	time (h)	yield [%] ^b , (<i>E</i> : <i>Z</i>) ^c	
			4b	5a
1	PCy ₃	3	19 (90:10)	44 (92:8)
2	PCy ₂ Ph	3	40 (89:11)	29 (94:6)
3	PCyPh ₂	1	70 (89:11)	4 (na)
4	PPh ₃	2	84 (87:13)	4 (na)

^a Pd(OAc)₂ (0.010 mmol), ligand (0.020 mmol), Cs₂CO₃ (0.24 mmol), **2a** (0.20 mmol), and **3b** (0.24 mmol) were stirred at 110 °C.

^b GC yield.

^c Determined by ¹H NMR of crude mixture.

Retro-allylative coupling of **2a-c** with various aryl bromides **3** was then examined (Table 3). The coupling with bromobenzene (**3c**) and *para*-substituted bromobenzene **3a** and **3d-f** were carried out in the presence of Pd/PCy₃ catalyst (entries 1-4, 7, and 9).⁹ These reactions were completed within 4-10 h, and the corresponding **4** was isolated in 61-77% yields. The results indicate that the electronic property of the aryl groups (R⁴ and R⁵) did not affect the reaction. On the other hand, coupling reactions with bulky aryl bromide **3b**, **3g**, and **3h** also gave the coupling product in high yields for 7-15 h in the presence of the Pd/PPh₃ catalyst (entries 5, 6, 8, and 10).

Table 3 Palladium-Catalyzed Retro-Allylative Coupling of **2**^a

entry	2	R ⁵ -Br	ligand, time (h)	product	yield [%] ^b , (<i>E</i> : <i>Z</i>) ^c
1	2a	PhBr (3c)	PCy ₃ , 9	4c	68 (86:14)
2	2a	4-MeOC ₆ H ₄ Br (3d)	PCy ₃ , 4	4d	63 (85:15)
3	2a	4-EtO ₂ CC ₆ H ₄ Br (3e)	PCy ₃ , 4	4e	67 (82:18)
4	2a	4-CF ₃ C ₆ H ₄ Br (3f)	PCy ₃ , 6	4f	77 (83:17)
5	2a	2-MeC ₆ H ₄ Br (3g)	PPh ₃ , 15	4g	61 (90:10)
6	2a	1-naphthylBr (3h)	PPh ₃ , 11	4h	60 (92:8)
7	2b	3a	PCy ₃ , 10	4i	68 (84:16)
8	2b	3b	PPh ₃ , 7	4j	89 (90:10)
9	2c	3a	PCy ₃ , 8	4k	61 (82:18)
10	2c	3b	PPh ₃ , 7	4l	73 (90:10)

^a Pd(OAc)₂ (0.010 mmol), ligand (0.020 mmol), Cs₂CO₃ (0.24 mmol), **2** (0.20 mmol), and **3** (0.24 mmol) were stirred at 110 °C.

^b Isolated yield.

^c Determined by ¹H NMR of crude mixture.

Homoallylic alcohols **6** bearing a tetrasubstituted double bond were subjected to retro-allylative coupling with **3a** in the presence of Pd/PCy₃ catalyst (Equation 2). The reaction of **6a** (R³ = Ph) completed in 6 h and gave desired product **7a** in 30% yield along with a significant

amount of the protonated product **8a**. On the other hand, **6b** ($R^3 = \text{Me}$) gave **7b** selectively (72%), despite the much slower reaction rate.

Equation 2

Homoallylic alcohols **2** and **6** were also subjected to the Pd/P(OPh)₃ catalyst system that promotes the protonative retro-allylation (Table 4).¹⁰ The reactions of **2b**, **2c**, and **6a** took place at 110 °C within 2 h to give crotononitrile derivatives **5b**, **5c**, and **8a** in good to high yields (entries 1-3). Although a slow reaction rate was observed for the reaction of **6b**, **8b** was finally isolated in 97% yield after 42 h (entry 4). It should be noted that the palladium-catalyzed retro-allylative coupling described above and protonative retro-allylation worked effectively for the homoallylic alcohols bearing a tetrasubstituted double bond (Equation 2 and entries 3 and 4 in Table 4), while the previous study demonstrated the failure of the retro-allylation of homoallylic alcohols bearing *E*-alkene moiety.⁴

Table 4 Palladium-Catalyzed Protonative Retro-Allylation of **2** and **6**^a

entry	substrate	R^3	R^4	time (h)	product	yield [%] ^b , (<i>E</i> : <i>Z</i>) ^c
1	2b	H	Ph	2	5b	67 (89:11)
2	2c	H	4-EtO ₂ CC ₆ H ₄	2	5c	63 (89:11)
3	6a	Ph	4-MeOC ₆ H ₄	6	8a	92 (62:38)
4	6b	Me	4-MeOC ₆ H ₄	42	8b	97 (33:67)

^a Pd(OAc)₂ (0.010 mmol), P(OPh)₃ (0.020 mmol), Cs₂CO₃ (0.24 mmol), and **2** or **6** (0.20 mmol) were stirred at 110 °C.

^b Isolated yield.

^c Determined by ¹H NMR of crude mixture.

The present retro-allylative coupling reaction is remarkable in that a substituted C=C bond including tetrasubstituted alkene is involved. To understand the high reactivity of the present retro-allylation, we examined the reaction of **9** (Equation 3). Homoallylic alcohol **9**, which does not have the aryl groups on the double bond, was subjected to the retro-allylative coupling with **3b** in the presence of the Pd/PPh₃ catalyst (Equation 3). The reaction proceeded smoothly to afford alkenes **10** in 69% yield, in contrast to the poor reactivity of internal disubstituted alkenes such as the (*E*)-pent-3-

en-1-ol derivatives observed previously.⁴ This result clearly indicates that the cyano group enhances the reactivity of homoallylic alcohol in palladium-catalyzed retro-allylation.

Equation 3

The possible reaction mechanism for the retro-allylative coupling is shown in Scheme 2. Oxidative addition of Ar¹-Br to Pd(0) followed by the substitution of the halogen atom on Pd with the alkoxide gives intermediate **T**. Retro-allylation of **T** provides σ -allylpalladium complex **U**, which isomerizes to π -allylpalladium complex **V**. Following reductive elimination affords product **W** with regeneration of Pd(0). The cyano group may decrease the rate of reductive elimination from **U**, leading to **V** via σ - π isomerization that is not involved in the retro-allylation of cyano-free homoallylic alcohols.⁴

Scheme 2 Possible Mechanism for Retro-Allylative Coupling

In summary, we have established a synthetic transformation of intramolecular cyanoboration products via C–C bond formation by Suzuki-Miyaura coupling followed by palladium-catalyzed retro-allylation. Efficient removal of a tethering group used in intramolecular cyanoboration was achieved through retro-allylation. Our new strategy, which utilizes “intramolecular reaction with removable tether”, allows collecting advantages of both the intra- and intermolecular reactions such as high reaction efficiency, selectivity, and wide reaction scope.

Acknowledgment

This work was supported by Grants-in-Aid for Scientific Research on Priority Areas “Advanced Molecular Transformations of Carbon Resources”, No 17065012 and 18037030, from Ministry of Education, Culture, Sports, Science and Technology, Japan.

References and Notes

- (1) For reviews of temporary-tethering strategy, see: (a) Bols, M.; Skrydstrup, T. *Chem. Rev.* **1995**, *95*, 1253. (b) Fensterbank, L.; Malacria, M.; Sieburth, S. M. *Synthesis* **1997**, 813. (c) Gauthier, Jr. D. R.; Zandi, K. S.; Shea, K. J. *Tetrahedron* **1998**, *54*, 289.
- (2) For selected recent examples of silicon-tethered intramolecular reaction, see: (a) Evans, P. A.; Baum, E. W. *J. Am. Chem. Soc.* **2004**, *126*, 11150. (b) Chouraqui, G.; Petit, M.; Aubert, C.; Malacria, M. *Org. Lett.* **2004**, *6*, 1519. (c) Someya, H.; Kondoh, A.; Sato, A.; Ohmiya, H.; Yorimitsu, H.; Oshima, K. *Synlett* **2006**, 3061. (d) Ohmura, T.; Furukawa, H.; Sugimoto, M. *J. Am. Chem. Soc.* **2006**, *128*, 13366. (e) Chen, C.-L.; Sparks, S. M.; Martin, S. F. *J. Am. Chem. Soc.* **2006**, *128*, 13696. (f) Kim, Y. J.; Lee, D. *Org. Lett.* **2006**, *8*, 5219.
- (3) For examples of boron-tethered intramolecular reaction, see: (a) Shimada, S.; Osoda, K.; Narasaka, K. *Bull. Chem. Soc. Jpn.* **1993**, *66*, 1254. (b) Nicolaou, K. C.; Liu, J.-J.; Yang, Z.; Ueno, H.; Sorensen, E. J.; Claiborne, C. F.; Guy, R. K.; Hwang, C.-K.; Nakada, M.; Nantermet, P. G. *J. Am. Chem. Soc.* **1995**, *117*, 634. (c) Batey, R. A.; Thadani, A. N.; Lough, A. J. *J. Am. Chem. Soc.* **1999**, *121*, 450. (d) Batey, R. A.; Smil, D. V. *Angew. Chem. Int. Ed.* **1999**, *38*, 1798. (e) Micalizio, G. C.; Schreiber, S. L. *Angew. Chem. Int. Ed.* **2002**, *41*, 3272. (f) Yamamoto, Y.; Ishii, J.; Nishiyama, H.; Itoh, K. *J. Am. Chem. Soc.* **2005**, *127*, 9625. (g) Yamamoto, A.; Sugimoto, M. *J. Am. Chem. Soc.* **2005**, *127*, 15706.
- (4) (a) Hayashi, S.; Hirano, K.; Yorimitsu, H.; Oshima, K. *J. Am. Chem. Soc.* **2006**, *128*, 2210. (b) Iwasaki, M.; Hayashi, S.; Hirano, K.; Yorimitsu, H.; Oshima, K. *J. Am. Chem. Soc.* **2007**, *129*, 4463. (c) Hayashi, S.; Hirano, K.; Yorimitsu, H.; Oshima, K. *J. Am. Chem. Soc.* **2007**, *129*, 12650.
- (5) For examples, see: (a) Tamao, K.; Maeda, K.; Tanaka, T.; Ito, Y. *Tetrahedron Lett.* **1988**, *29*, 6955. (b) Murakami, M.; Oike, H.; Sugawara, M.; Sugimoto, M.; Ito, Y. *Tetrahedron* **1993**, *49*, 3933. (c) Sugimoto, M.; Kinugasa, H.; Ito, Y. *Tetrahedron Lett.* **1994**, *35*, 8635. (d) Ojima, I.; Vidal, E.; Tzamarioudaki, M.; Matsuda, I. *J. Am. Chem. Soc.* **1995**, *117*, 6797. (e) O'Malley, S. J.; Leighton, J. L. *Angew. Chem. Int. Ed.* **2001**, *40*, 2915. (f) Denmark, S. E.; Pan, W. *Org. Lett.* **2002**, *4*, 4163. (g) Trost, B. M.; Ball, Z. T. *J. Am. Chem. Soc.* **2003**, *125*, 30.
- (6) (a) Sugimoto, M.; Yamamoto, A.; Murakami, M. *J. Am. Chem. Soc.* **2003**, *125*, 6358. (b) Sugimoto, M.; Yamamoto, A.; Murakami, M. *J. Organomet. Chem.* **2005**, *690*, 5300.
- (7) We have also developed intermolecular cyanoboration of alkynes, see: Sugimoto, M.; Yamamoto, A.; Murakami, M. *Angew. Chem. Int. Ed.* **2005**, *44*, 2380.
- (8) Littke, A. F.; Dai, C.; Fu, G. C. *J. Am. Chem. Soc.* **2000**, *122*, 4020.
- (9) **General Procedure for the Palladium-Catalyzed Retro-Allylative Coupling of 2 with 3 (Tables 1-3):** Cesium carbonate was dried in vacuo with heating with heat gun prior to use. Under a nitrogen atmosphere, a mixture of Pd(OAc)₂ (0.010 mmol), PCy₃ or PPh₃ (0.020 mmol), Cs₂CO₃ (0.24 mmol), **2** (0.20 mmol), and **3** (0.24 mmol) was heated at 110 °C. The reaction was monitored by GC.

Heating was stopped as soon as **2** was consumed, since prolonged heating led drop of the product yield. Volatiles were removed and the crude product was purified by column chromatography on silica gel.

(*E*)-3-(4-Methoxyphenyl)-4-(4-methylphenyl)but-2-enenitile (**4a**): ¹H NMR (400 MHz, CDCl₃) δ 7.38 (d, *J* = 8.8 Hz, 2H), 7.02-7.12 (m, 4H), 6.83 (d, *J* = 8.8 Hz, 2H), 5.65 (s, 1H), 4.16 (s, 2H), 3.80 (s, 3H), 2.28 (s, 3H). ¹³C NMR (126 MHz, CDCl₃) δ 161.6, 161.2, 136.3, 134.0, 129.5, 129.4 (2C), 128.20 (2C), 128.18 (2C), 118.0, 114.1 (2C), 94.8, 55.3, 39.0, 21.0. IR (neat) 2211 (ν_{CN}), 1605 (ν_{C=C}) cm⁻¹. LRMS (EI) *m/z* 263 (M⁺, 100), 248 (18), 158 (38), 133 (40), 105 (41). HRMS (EI) *m/z* calcd for C₁₈H₁₇NO (M⁺): 263.1310, found: 263.1311. The geometry of the double bond was assigned as *E* by NOE experiments.

(*E*)-4-(2,6-Dimethylphenyl)-3-(4-methoxyphenyl)but-2-enenitile (**4b**): ¹H NMR (400 MHz, CDCl₃) δ 7.06-7.11 (m, 3H), 6.98-7.03 (m, 2H), 6.78 (d, *J* = 8.8 Hz, 2H), 5.49 (t, *J* = 1.2 Hz, 1H), 4.17 (d, *J* = 1.2 Hz, 2H), 3.78 (s, 3H), 2.29 (s, 6H). ¹³C NMR (126 MHz, CDCl₃) δ 162.7, 160.6, 137.5 (2C), 133.5, 131.1, 128.3 (2C), 127.8 (2C), 127.1, 116.6, 113.7 (2C), 96.3, 55.2, 35.7, 20.5 (2C). IR (KBr): 2207 (ν_{CN}), 1605 (ν_{C=C}) cm⁻¹. LRMS (EI) *m/z* 277 (M⁺, 80), 237 (100), 119 (40). HRMS (EI) *m/z* calcd for C₁₉H₁₉NO (M⁺): 277.1467, found: 277.1474. The geometry of the double bond was assigned as *E* by NOE experiments.

(10) **General Procedure for the Palladium-Catalyzed Protonative Retro-Allylation of 2 (Entry 9 in Table 1 and Table 4):** Cesium carbonate was dried in vacuo with heating with heat gun prior to use. Under a nitrogen atmosphere, a mixture of Pd(OAc)₂ (0.010 mmol), P(OPh)₃ (0.020 mmol), Cs₂CO₃ (0.24 mmol), and **2** (0.20 mmol) was heated at 110 °C. The reaction was monitored by GC. After consumption of **2**, the mixture was extracted with Et₂O, and organic layer was dried over anhydrous Mg₂SO₄. The product was isolated by column chromatography on silica gel.

(*E*)-3-(4-Methoxyphenyl)but-2-enenitile (**5a**): ¹H NMR (400 MHz, CDCl₃) δ 7.41-7.45 (m, 2H), 6.89-6.93 (m, 2H), 5.55 (d, *J* = 0.8 Hz, 1H), 3.85 (s, 3H), 2.45 (d, *J* = 0.8 Hz, 3H). ¹³C NMR (126 MHz, CDCl₃) δ 161.3, 158.8, 130.4, 127.3 (2C), 118.1, 114.1 (2C), 93.2, 55.4, 20.0. IR (KBr) 2205 (ν_{CN}), 1603 (ν_{C=C}) cm⁻¹. LRMS (EI) *m/z* 173 (M⁺, 100), 158 (41), 103 (35). HRMS (EI) *m/z* calcd for C₁₁H₁₁NO (M⁺): 173.0841, found: 173.0841. The geometry of the double bond was assigned as *E* by NOE experiments.

Synthetic Application of Intramolecular Cyanoboration