TITLE:

Direct and indirect electrochemical generation of alkoxycarbenium ion pools from thioacetals

AUTHOR(S):

Matsumoto, Kouichi; Ueoka, Koji; Suzuki, Shinkiti; Suga, Seiji; Yoshida, Jun-ichi

CITATION:

Matsumoto, Kouichi ...[et al]. Direct and indirect electrochemical generation of alkoxycarbenium ion pools from thioacetals. Tetrahedron 2009, 65(52): 10901-10907

ISSUE DATE:

2009-12-26

URL:

<http://hdl.handle.net/2433/89513>

RIGHT:

c 2009 Elsevier Ltd. All rights reserved.; この論文は出版社版ではありません。引用の際には出版社版をご確認ご利用ください。; This is not the published version. Please cite only the published version.

Graphical Abstract

To create your abstract, type over the instructions in the template box below.
 Fonts or abstract dimensions should not be changed or altered.

Direct and Indirect Electrochemical Generation of Alkoxycarbenium Ion Pools from Thioacetals

Kouichi Matsumoto, Koji Ueoka, Shinkiti Suzuki, Seiji Suga¹ and Jun-ichi Yoshida*
*Department of Synthetic Chemistry and Biological Chemistry, Graduate School of Engineering, Kyoto University,
 Nishikyo-ku, Kyoto 615-8510, Japan*

Direct and Indirect Electrochemical Generation of Alkoxycarbenium Ion Pools from Thioacetals

Kouichi Matsumoto, Koji Ueoka, Shinkiti Suzuki, Seiji Suga¹ and Jun-ichi Yoshida^{a*}

^aDepartment of Synthetic Chemistry and Biological Chemistry, Graduate School of Engineering, Kyoto University, Nishikyo-ku, Kyoto 615-8510, Japan.

Abstract—Thioacetals were found to be effective precursors to generate and accumulate alkoxycarbenium ions based on direct and indirect cation pool methods. Alkoxycarbenium ions thus-generated reacted with carbon nucleophiles such as allylsilanes and enol silyl ethers to give C-C bond formation products in good yields. © 2009 Elsevier Science. All rights reserved

1. Introduction

Alkoxycarbenium ions, carbocations stabilized by a neighboring alkoxy group, are important intermediates in organic synthesis.² Usually alkoxycarbenium ions are generated from acetals using Lewis acids such as $\text{BF}_3\cdot\text{OEt}_2$, SnCl_4 and TiCl_4 in the presence of a nucleophile (Scheme 1). Although benzylic alkoxycarbenium ions and di- and tri(alkoxy)carbenium ions are stable and are well characterized spectroscopically,³ simple alkylalkoxycarbenium ions are unstable and are difficult to characterize spectroscopically. The generation processes for such unstable alkoxycarbenium ions are reversible, and their equilibrium concentrations are usually very low. In fact, extensive NMR studies on the reaction of acetals with Lewis acids revealed the presence of Lewis acid-acetal complexes, but alkoxycarbenium ions were not detected.⁴

Scheme 1.

The electrochemical method is also effective for generation of alkoxycarbenium ions.⁵ Recently we have developed the “cation pool” method, in which organic cations are generated and accumulated in the absence of nucleophiles by low-temperature electrolysis.⁶ The cation pool method serves as a powerful tool not only for mechanistic studies on highly reactive organic cations but also for rapid parallel synthesis. The method has been successfully applied to alkoxycarbenium ions,⁷ in addition to *N*-acyliminium ions,⁸ diarylcarbenium ions,⁹ organosilicon cations,¹⁰ and iodine cations.¹¹ α -Silyl ethers serve as effective precursors of alkoxycarbenium ion pools (Scheme 2). Silyl groups serve as effective electroauxiliaries, which lower the oxidation potentials

and control the regiochemistry by virtue of selective C-Si bond cleavage.¹² Alkylalkoxycarbenium ions thus-generated are well characterized by NMR spectroscopy, which exhibits very similar spectra to those generated in super acid media.¹³ In the next step, the cations react with carbon nucleophiles such as allylsilanes to give the corresponding carbon-carbon bond formation products.

Scheme 2.

We have been searching for alternative precursors of alkoxycarbenium ion pools and have been interested in thioacetals as precursors because of the following reasons: 1) An arylthio group is also an effective electroauxiliary for electrochemical oxidation.¹⁴ 2) Thioacetals can be easily prepared. In preliminary communications,¹⁵ we have reported generation of alkoxycarbenium ion pools from thioacetals by low-temperature electrolysis (direct cation pool method) and by the action of $\text{ArS}(\text{ArSSAr})^+$, which is generated by low-temperature electrolysis of ArSSAr (indirect cation pool method) (Scheme 3). In this paper we report full details of these studies.

Scheme 3.

* Corresponding author. Tel.: +81-75-383-2726; fax: +81-75-383-2727; e-mail: yoshida@sbchem.kyoto-u.ac.jp.

2. Result and Discussion

2.1 Direct Cation Pool Method

At first, we examined the direct electrochemical method using thioacetal **1** as a precursor of an alkoxy-carbenium ion. Thus, a solution of **1** was electrolyzed in $\text{Bu}_4\text{NBF}_4/\text{CD}_2\text{Cl}_2$ at -78°C using an H-type divided cell (Scheme 4). After the electrolysis (1.4 F/mol), the resulting solution was analyzed by NMR spectroscopy at -80°C . ^1H and ^{13}C NMR spectra (-80°C) indicated the formation of alkoxy-carbenium ion (**2**) by selective cleavage of the C-S bond. ^1H NMR exhibited a signal at δ 9.55 ppm due to the methine proton. ^{13}C NMR exhibited a signal at δ 230.9 ppm due to the methine carbon. These chemical shifts are quite similar to those obtained by the direct electrochemical oxidation of $\text{C}_8\text{H}_{17}\text{CH}(\text{OMe})\text{SiMe}_3$ (9.55 and 231.0 ppm),^{7a} suggesting that the alkoxy-carbenium ion is efficiently generated and accumulated as a single species in irreversible fashion. Though PhSSPh is generated by the oxidation of **1**, the oxidation potential of **1** (RDE decomposition potential: 1.30 V vs. SCE) is lower than that of PhSSPh (1.42 V), indicating that PhSSPh cannot be a mediator for the oxidation of **1**.

Scheme 4.

The reaction of **2** with allyltrimethylsilane gave the allylated product **3** in 60% (Scheme 5). The electrochemical method was found to be applicable to cyclic thioacetals (**4** and **5**), which were effectively oxidized under similar conditions. After the electrolysis, the resulting solutions were allowed to react with an allylsilane to give the corresponding C-C bond formation products **6** and **7** in 82% and 38% yields, respectively.

Scheme 5.

2.2 Indirect Cation Pool Method

Because electrochemical reactions take place only on the surface of the electrode, alkoxy-carbenium ions generated in an early stage of the electrolysis should stay in the solution until the electrolysis is completed. This means the application of direct cation pool method strongly depends on the stability of accumulated alkoxy-carbenium ions.¹⁶ During the accumulation, unstable alkoxy-carbenium ions might decompose. To solve this problem, we examined indirect electrochemical method.¹⁷ In the

first step, an active chemical reagent is generated by the electrolysis, which is allowed to react with a precursor to generate an alkoxy-carbenium ion pool in the second step. The second reaction can be faster than the electrolysis because it takes place in a homogeneous solution.

We envisaged that ArS^{+} ¹⁸ generated by the electrochemical oxidation of ArSSAr might be suitable for the conversion of thioacetals to alkoxy-carbenium ion pools. It is known in the literature that ArS^{+} ¹⁹ or its equivalent $\text{ArS}(\text{ArSSAr})^+$ ²⁰ serves as a powerful and highly thiophilic reagent, though the nature of the generated species has not yet been fully elucidated.²¹

Thus, the electrolysis of ArSSAr (Ar = $p\text{-XC}_6\text{H}_4$) (oxidation potential: X = F: 1.47 V, X = CH_3 : 1.34 V, X = OMe: 1.47 V, X = Cl: 1.27 V, 2 equiv based on thioacetal **1**) was carried out in $\text{Bu}_4\text{NBF}_4/\text{CH}_2\text{Cl}_2$ at -78°C using an H-type divided cell (Scheme 6, step 1). In the next step, the electrogenerated $\text{ArS}(\text{ArSSAr})^+$ was reacted with thioacetal **1** to generate alkoxy-carbenium ion **2** (step 2). This process requires only 5 min at -78°C . In the third step, allyltrimethylsilane (10 equiv) was added to obtain the final product **3**.

Scheme 6.

The results obtained under various conditions are summarized in Table 1. Electrolysis with 1.34 F/mol based on **1** (0.67 F/mol based on ArSSAr) gave the best yield of **3** (entry 2). 0.67 F/mol is the theoretical amount of electricity to convert ArSSAr to $\text{ArS}(\text{ArSSAr})^+$. This means that 1.34 equiv of $\text{ArS}(\text{ArSSAr})^+$ based on **1** was formed, because 2 equiv of ArSSAr based on **1** was used. Almost quantitative yield of **3** (98%) suggests that alkoxy-carbenium ion **2** was generated almost quantitatively. High reactivity of $\text{ArS}(\text{ArSSAr})^+$ and homogeneity of the reaction system seem to be responsible for fast and quantitative generation of **2**. Use of 1.5 equiv of ArSSAr led to a decrease in the yield of **3** (entry 5). Use of 3.0 equiv of ArSSAr also led to a decrease in the yield of **3** (entry 6). Other ArSSAr (Ar = C_6H_5 , $p\text{-CH}_3\text{C}_6\text{H}_4$, $p\text{-MeOC}_6\text{H}_4$ and $p\text{-ClC}_6\text{H}_4$) were also effective, although the yield of **3** depends on the nature of the substituent on the aryl group (entries 7-10).

Table 1. Optimization of the indirect cation pool method

Entry	X	ArSSAr (eq based on 1)	Electricity (F/mol based on ArSSAr)	Yield (%) of 3
1	F	2	0.50	69
2	F	2	0.67	98
3	F	2	0.75	91
4	F	2	1.0	87
5	F	1.5	0.67	73
6	F	3	0.67	86
7	H	2	0.67	79
8	CH ₃	2	0.67	79
9	OMe	2	0.67	69
10	Cl	2	0.67	96

The formation of **2** was confirmed by NMR spectroscopy at -80 °C. A solution obtained by the reaction of **1** with the electrogenerated $\text{ArS}(\text{ArSSAr})^+$ (Ar = *p*-FC₆H₄) exhibited signals at 9.53 and 4.92 ppm due to the methine proton and methyl protons, respectively (¹H NMR), and a signal at 230.6 ppm due to the methine carbon (¹³C NMR) (Scheme 7a). These chemical shifts are quite similar to those obtained by the direct electrochemical oxidation of C₈H₁₇CH(OMe)SiMe₃ (9.55, 4.95 and 231.0 ppm).^{7a} Such similarity in chemical shifts indicated that the sulfur-containing byproducts, such as PhSSAr and ArSSAr, which should be present in the solution, did not change the nature of alkoxy-carbenium ion **2** appreciably.²² We also generated and analyzed an alkoxy-carbenium ion by the reaction of **5** with $\text{ArS}(\text{ArSSAr})^+$ (Ar = *p*-FC₆H₄). The cation exhibited a signal at 9.86 ppm due to the methine proton (¹H NMR) and a signal at 227.8 ppm due to the methine carbon (¹³C NMR) (Scheme 7b).

Scheme 7.

The detailed mechanism for the reaction of **1** with $\text{ArS}(\text{ArSSAr})^+$ (step 2) has not been clarified as yet, but **2** seems to be generated according to Scheme 8. Though the possibility of a single electron-transfer mechanism cannot be ruled out, ionic mechanism seems to be more plausible. In the ionic mechanism $\text{ArS}(\text{ArSSAr})^+$ acted as a thiophilic Lewis acid.

Scheme 8.

In order to get a deeper insight into the mechanism, reactions of **1** with other Lewis acids such as BF₃·OEt₂ and SnCl₄ were

examined.²³ The resulting solution was analyzed by NMR spectroscopy at -80 °C, but alkoxy-carbenium ion **2** was not detected at all.⁴ Presumably these Lewis acids are not strong enough to generate **2** in a significant concentration. The equilibrium between **1** and **2** lies to **1**.

Much lower reactivity of BF₃·OEt₂ was confirmed by the following experiments. The reaction of thioacetal (**1**) with $\text{ArS}(\text{ArSSAr})^+$ (Ar = *p*-FC₆H₄) in 5 min followed by treatment with allyltrimethylsilane in 1 min at -78 °C gave **3** in 90% yield (eq 1). In contrast, the reaction using a BF₃·OEt₂ did not give **3** in an appreciable amount under similar conditions (eq 2).

As described above, alkoxy-carbenium ion **2** generated by the indirect method exhibited NMR spectra similar to that generated by the direct anodic oxidation of C₈H₁₇CH(OMe)SiMe₃.^{7a} Thus, we next compared the thermal stability of **2** as follows: The cation pool generated at -78 °C was allowed to warm to a second temperature. After being kept there for 30 min, the pool was allowed to react with allyltrimethylsilane. The yield of **3** is plotted against the temperature in Figure 1. The cation pool of **2** generated by the indirect method exhibited thermal stability similar to that generated by the direct method of C₈H₁₇CH(OMe)SiMe₃.^{7a}

Figure 1. Thermal stability of alkoxy-carbenium ion **2** generated by the direct method and the present indirect method.

To test the applicability of the present indirect method, reactions with several carbon nucleophiles were examined (Table 2). Allylsilanes, enol silyl ethers, and ketene silyl acetals, and enol acetate were effective as carbon nucleophiles, and the corresponding C-C bond formation products were obtained. 1,3-dicarbonyl compounds such as acetylacetone were also effective to form carbon-carbon bond.

Table 2. The reaction of alkoxy-carbenium ion with various carbon nucleophiles^{a)}

Nu	Product	Yield (%) ^{b)}
		89 ^{c)} (98) ^{d), e)}
		91 (1.25:1) ^{e)}
		79
		42 (2.45:1) ^{e)}
		67
		63 ^{f)}
		43 ^{d)}

a) ArSSAr (Ar = *p*-FC₆H₄, 0.40 mmol) was oxidized electrochemically in 0.3 M Bu₄NBF₄/CH₂Cl₂ at -78 °C using 0.67 F/mol of electricity. The solution thus obtained was allowed to react with 0.20 mmol of thioacetal at -78 °C for 5 min. Then a nucleophile (0.50 mmol, 2.5 equiv) was added, and the resulting solution was stirred at -78 °C for 15 min and the reaction was quenched with Et₃N (1 mL). b) Isolated yield. c) GC yield. d) 10 eq of Nu was used. e) Diastereomer ratio. f) The reaction time with a nucleophile was 30 min.

Table 3. Indirect generation of various alkoxy-carbenium ion pools followed by reactions with carbon nucleophiles

Substrate	Nu	Product	Yield (%) ^{a)}
			82 ^{b)}
			83 ^{b)}
			72 ^{b)}
			91 ^{c)}
			68
			86 (ca 9:1) ^{d)}
			85 (ca 7:1) ^{d)}

a) Isolated yield. b) GC yield. c) The acetal was reacted with ArS(ArSSAr)⁺ for 10 min, the resulting alkoxy-carbenium ion was reacted with allyltrimethylsilane for 1 h. d) Diastereomer ratio.

Finally, we examined generation of alkoxy-carbenium ion pools from several thioacetals. The *p*-ClC₆H₄S, *p*-FC₆H₄ and *p*-MeC₆H₄S groups were also effective as electroauxiliaries. Aryl and alkyl substituted thioacetals, including cyclic substrates, were effective for generation of alkoxy-carbenium ion pools. The resulting alkoxy-carbenium ion pools reacted with carbon nucleophiles such as allylsilanes to give the corresponding C-C bond formation products in good yields as shown in Table 3.

3. Conclusions

In conclusion, we have developed a method for generation and accumulation of alkoxy-carbenium ions from thioacetals by direct low-temperature electrolysis. We have also developed a sequential one-pot indirect method for generation and accumulation of alkoxy-carbenium ions. In the first step, ArSSAr is oxidized by low temperature electrolysis to generate and accumulate ArS(ArSSAr)⁺, which is allowed to react with a thioacetal to generate and accumulate an alkoxy-carbenium ion. The cation pool thus-generated reacts with various carbon nucleophiles such as allylsilanes and silyl enol ethers. The direct and indirect methods for generation of alkoxy-carbenium ion pools from thioacetals add a new dimension to organic cation chemistry and organic electrochemistry.

4. Experimental Section

4.1. General Remarks

GC analysis was performed on a gas chromatograph (SHIMADZU GC-14B) equipped with a flame ionization detector using a fused silica capillary. ¹H and ¹³C NMR spectra were recorded in CDCl₃ on a Varian Gemini 2000 (¹H 300 MHz, ¹³C 75 MHz), Varian MERCURY plus-400 (¹H 400 MHz, ¹³C 100 MHz), JEOL A-500 (¹H 500 MHz, ¹³C 125 MHz), or JEOL ECA-600P (¹H 600 MHz, ¹³C 150 MHz) spectrometer with Me₄Si as an internal standard unless otherwise noted. Mass spectra were obtained on JEOL JMS SX-102A mass spectrometer. IR spectra were measured with a SHIMADZU FTIR 1600 spectrometer. Thinlayer chromatography (TLC) was carried out by using Merck precoated silica gel F254 plates (thickness 0.25 mm). Flash chromatography was carried out on a column of silica gel (Kanto Chem. Co., Silica Gel N, spherical, neutral, 40-100 mm). Gel permeation chromatography (GPC) was carried out on Japan Analytical Industry LC-908 equipped with JAIGEL-1H and 2H using CHCl₃ as eluent. All reactions were carried out under Ar atmosphere unless otherwise noted.

4.2. Materials

Dichloromethane was washed with water, distilled from P₂O₅, redistilled from dried K₂CO₃ to remove a trace amount of acid, and stored over molecular sieves 4A. Dry THF was used as obtained (Kanto Chemical Co., Inc.). ArSSAr (Ar = *p*-FC₆H₄) was prepared according to the procedures in the literatures,²⁴ and identified by the comparison of its spectral data with that of authentic sample.²⁵

4.3. Thioacetals

1-Methoxy-1-phenylthiononane (**1**),^{14c} 2-phenylthiotetrahydrofuran (**4**)²⁶ and 2-phenylthiotetrahydropyran (**5**)²⁶ were prepared according to the literature procedures.

4.3.1. 1-Methoxy-1-(4-chlorophenylthio)nonane (**14**).

To a solution of 4-chlorobenzenethiol (5.19 g, 36.0 mmol) in tetrahydrofuran (THF) (64 mL), was added triethylamine (7.2 mL, 52 mmol) and chloromethyl methyl ether (3.3 mL, 43 mmol) at 0

°C. After being stirred at room temperature for 2 h, the reaction mixture was partitioned between saturated aqueous NH_4Cl and ether. The organic phase was separated, washed with saturated aqueous NaHCO_3 , and was dried over MgSO_4 . The solvent was removed under reduced pressure and the residue was purified by distillation (130 °C, 14 mmHg), to obtain (4-chlorophenylthio)methyl methyl ether (5.76 g, 30.5 mmol).

To a solution of (4-chlorophenylthio)methyl methyl ether (1.51 g, 8.0 mmol) in THF (20 mL), was added butyllithium (1.54 M in hexane, 6.4 mL, 10 mmol) at -78 °C. The mixture was stirred at -45 °C for 3 h and was cooled to -78 °C. 1-Iodooctane (2.62 g, 10.9 mmol) was added, and the mixture was stirred at -78 °C for 20 min. Saturated aqueous NH_4Cl was added. The organic materials were extracted with ether, and the organic extracts were washed with brine and dried over MgSO_4 . After removal of the solvent, the residue was purified by flash chromatography (hexane) to obtain the title compound (1.60 g, 5.3 mmol, 66 %): TLC R_f 0.73 (hexane/EtOAc 20:1); ^1H NMR (400 MHz, CDCl_3) δ 0.87 (t, J = 6.8 Hz, 3H), 1.21-1.28 (m, 10H), 1.40-1.43 (m, 2H), 1.36-1.44 (m, 2H), 1.62-1.78 (m, 2H), 3.46 (s, 3H), 4.57 (t, J = 6.8 Hz, 1H), 7.23-7.26 (m, 2H), 7.36-7.40 (m, 2H); ^{13}C NMR (100 MHz, CDCl_3) δ 14.2, 22.8, 26.3, 29.2, 29.3, 29.5, 31.9, 35.6, 55.4, 90.9, 128.7, 131.6, 133.5, 134.6; IR (neat) 2953, 1476, 1130, 1092, 623 cm^{-1} ; LRMS (EI) m/z 300 (M^+), 269 ($\text{M}^+ - \text{OCH}_3$), 187 ($\text{M}^+ - \text{C}_8\text{H}_{17}$), 157 ($\text{M}^+ - \text{SC}_6\text{H}_4p\text{-Cl}$); HRMS (EI) calcd for $\text{C}_{16}\text{H}_{25}\text{OClS}$: 300.1315, found: 300.1313.

4.3.2. 1-Methoxy-1-(4-fluorophenylthio)nonane (15).

To a solution of pelargonaldehyde dimethyl acetal (4.03 g, 21.4 mmol) and *p*-fluorothiophenol (2.30 mL, 21.5 mmol) in toluene (180 mL), was added $\text{BF}_3 \cdot \text{OEt}_2$ (2.7 mL, 21.5 mmol) at -78 °C. The solution was stirred for 2 h. Saturated aqueous NaHCO_3 was added. The reaction mixture was partitioned between Et_2O and saturated aqueous NaHCO_3 . The organic phase was separated and washed with saturated aqueous NaHCO_3 and dried over MgSO_4 . After removal of the solvent, the crude product was purified by flash chromatography (hexane/EtOAc 30:1) to give the title compound (15) (4.23 g, 14.9 mmol, 70 %): TLC R_f 0.28 (hexane/EtOAc 20:1); ^1H NMR (400 MHz, CDCl_3) δ 0.87 (t, J = 6.8 Hz, 3H), 1.18-1.32 (m, 10H), 1.34-1.46 (m, 2H), 1.58-1.74 (m, 2H), 3.47 (s, 3H), 4.51 (t, J = 6.4 Hz, 1H), 6.94-7.02 (m, 2H), 7.39-7.45 (m, 2H); ^{13}C NMR (75 MHz, CDCl_3) δ 14.1, 22.6, 26.2, 29.1, 29.2, 29.4, 31.8, 35.5, 55.5, 91.0, 115.8 (d, J = 21.7 Hz), 127.9 (d, J = 3.4 Hz), 136.1 (d, J = 8.0 Hz), 162.7 (d, J = 246.0 Hz); IR (neat) 2926, 1590, 1491, 830 cm^{-1} ; LRMS (EI) m/z 284 (M^+), 157 ($\text{M}^+ - \text{SC}_6\text{H}_4p\text{-F}$); HRMS (EI) calcd for $\text{C}_{16}\text{H}_{25}\text{FOS}$ (M^+) 284.1610, found 284.1605.

4.3.3. 1-Methoxy-1-(4-methylphenylthio)nonane (16).

To a solution of 4-methylbenzenethiol (6.13 g, 49.4 mmol) in THF (100 mL), was added triethylamine (10 mL, 73 mmol) and chloromethyl methyl ether (4.69 g, 58.3 mmol) at 0 °C. After being stirred at room temperature for 2 h, the reaction mixture was partitioned between saturated aqueous NH_4Cl and ether. The organic phase was separated, washed with saturated aqueous NaHCO_3 , and was dried over MgSO_4 . The solvent was removed under reduced pressure and the residue was purified by distillation to obtain (4-methylphenylthio)methyl methyl ether (6.71 g, 39.9 mmol).

To a solution of (4-methylphenylthio)methyl methyl ether (3.21 g, 19.1 mmol) in THF (45 mL), was added butyllithium (1.54 M in hexane, 15 mL, 23.1 mmol) at -78 °C. The mixture was stirred

at -45 °C for 3 h and cooled to -78 °C. 1-Iodooctane (5.90 g, 24.6 mmol) was added, and the mixture was stirred at -78 °C for 30 min. Saturated aqueous NH_4Cl was added. The organic materials were extracted with ether, and the organic extracts were washed with brine and dried over MgSO_4 . After removal of the solvent, the residue was purified by flash chromatography (hexane/EtOAc 50:1) to obtain the title compound (4.43 g, 15.8 mmol, 83 %): TLC R_f 0.42 (hexane/EtOAc 20:1); ^1H NMR (300 MHz, CDCl_3) δ 0.87 (t, J = 6.6 Hz, 3H), 1.22-1.34 (m, 10H), 1.36-1.52 (m, 2H), 1.64-1.80 (m, 2H), 2.32 (s, 3H), 3.46 (s, 3H), 4.54 (t, J = 6.6 Hz, 1H), 7.09 (m, 2H), 7.35 (m, 2H); ^{13}C NMR (75 MHz, CDCl_3) δ 14.1, 21.1, 22.6, 26.2, 29.1, 29.2, 29.4, 31.8, 35.6, 55.4, 91.1, 129.4, 129.4, 134.0, 137.5; IR (neat) 2924, 1466, 1088, 644 cm^{-1} ; LRMS (EI) m/z 280 (M^+), 157 ($\text{M}^+ - \text{SC}_6\text{H}_4p\text{-CH}_3$); HRMS (EI) calcd for $\text{C}_{17}\text{H}_{28}\text{OS}$: 280.1861, found: 280.1859.

4.3.4. (Methoxy(phenylthio))methylbenzene (17).

To a toluene solution of benzaldehyde dimethyl acetal (786.8 mg, 5.17 mmol) and thiophenol (612.0 mg, 5.55 mmol), was added $\text{BF}_3 \cdot \text{OEt}_2$ (736.0 mg, 5.19 mmol) at -78 °C. The solution was stirred for 2 h. Dry pyridine (2.4 mL) was added. The mixture was diluted with ether and was washed with 1 M NaOH solution and water. After drying over Na_2SO_4 and removal of solvent, colorless oil thus obtained was purified by flash chromatography (hexane/EtOAc 50:1 to 10:1) to obtain the title compound (757.8 mg, 3.29 mmol, 64 %).²⁷

4.3.5. (Methoxy(phenylthio)methyl)cyclohexane (19).

To a toluene solution of cyclohexanecarbaldehyde dimethyl acetal (774.0 mg, 4.89 mmol) and thiophenol (600 mg, 5.45 mmol) was added $\text{BF}_3 \cdot \text{OEt}_2$ (901.3 mg, 6.35 mmol) at -78 °C. The solution was stirred for 2 h, and dry pyridine (0.5 mL) was added. The mixture was diluted with ether and was washed with 1 M NaOH solution and water. After drying by Na_2SO_4 and removal of solvent, colorless oil thus obtained was purified by flash chromatography (hexane/EtOAc 50:1 to 10:1) and GPC to give the title compound (918.6 mg, 3.89 mmol, 79 %).²⁷

4.4 Generation of Alkoxy-carbenium Ions (Direct Cation Pool Method). Typical Procedure.

The anodic oxidation was carried out in an H-type divided cell (4G glass filter) equipped with a carbon felt anode (Nippon Carbon JF-20-P7, ca. 320 mg, dried at 250 °C/1 mmHg for 1 h before use) and a platinum plate cathode (40 mm x 20 mm). In the anodic chamber was placed a solution of 1-methoxy-1-phenylthiononane (1) (94.9 mg, 0.356 mmol) in 0.3 M $\text{Bu}_4\text{NBF}_4/\text{CH}_2\text{Cl}_2$ (8.0 mL). In the cathodic chamber were placed 0.3 M $\text{Bu}_4\text{NBF}_4/\text{CH}_2\text{Cl}_2$ (8.0 mL) and trifluoromethanesulfonic acid (150.7 mg, 1.00 mmol). The constant current electrolysis (8 mA) was carried out at -78 °C with magnetic stirring until 2.5 F/mol of electricity was consumed.

4.5. NMR Analysis of Alkoxy-carbenium Ion (2) based on Direct Cation Pool Method.

^1H and ^{13}C NMR spectra were recorded in CD_2Cl_2 on a JEOL A-500 spectrometer. Chemical shifts are reported using methylene signals of CH_2Cl_2 at δ 5.32 (^1H NMR) and δ 53.80 (^{13}C NMR) as standards. The anodic oxidation was carried out in a divided cell equipped with a carbon felt anode and a platinum plate cathode. In the anodic chamber were placed a solution of 1-methoxy-1-phenylthiononane (1) (49.9 mg, 0.187 mmol) in 3.7 mL of 0.3 M

$\text{Bu}_4\text{NBF}_4/\text{CD}_2\text{Cl}_2$. In the cathodic chamber were placed trifluoromethanesulfonic acid (71.1 mg, 0.474 mmol) and 0.3 M $\text{Bu}_4\text{NBF}_4/\text{CD}_2\text{Cl}_2$ (3.7 mL). The constant current electrolysis (3.7 mA) was carried out at -78°C with magnetic stirring. After 1.4 F/mol of electricity was consumed, the reaction mixture of the anodic chamber was transferred to a 5 mm ϕ NMR tube with a septum cap under Ar atmosphere at -78°C . The NMR measurement was carried out at -80°C : ^1H NMR (500 MHz, CD_2Cl_2 , selected) δ 4.95 (s, 3H), 9.55 (s, 1H); ^{13}C NMR (125 MHz, CD_2Cl_2 , selected) δ 40.7 (CH_3OCH_2), 76.1 (CH_3OCH), 230.9 (CH_3OCH). In the NMR spectra the other signals could not be assigned because of overlap of the signals of Bu_4NBF_4 used as electrolyte. The signals due to the thiophenyl group leaving from **1** were also observed: ^1H NMR δ 7.2–8.0 (m); ^{13}C NMR δ 116.5, 119.0, 124.1, 130.3.

4.6. Reaction of Alkoxy-carbenium Ion and Carbon Nucleophiles. 4-Methoxy-dodec-1-ene (**3**) (A Typical Procedure of Direct Cation Pool Method).

The electrolysis of **1** (94.9 mg, 0.356 mmol) was carried out as described above. To the “cation pool” thus generated in the anodic chamber, was added allyltrimethylsilane (99.7 mg, 0.873 mmol) at -78°C and the reaction mixture was stirred for 15 min. The solvent was removed under reduced pressure and the residue was quickly filtered through a short column (2 x 3 cm) of silica gel to remove Bu_4NBF_4 . The silica gel was washed with ether (150 mL). The GC analysis of the combined filtrate indicated that **3** was formed in 60% yield (GC t_R 6.3 min, column, OV-1; 0.25 mm x 25 m; initial oven temperature, 100°C ; rate of temperature increase, $10^\circ\text{C}/\text{min}$). The isolated product was identified by the comparison of its spectral data with those of an authentic sample.^{7a}

4.7. Generation of Alkoxy-carbenium Ions (Indirect Cation Pool Method). Typical Procedure.

The anodic oxidation was carried out in an Htype divided cell (4G glass filter) equipped with a carbon felt anode and a platinum plate cathode (40 mm x 20 mm). In the anodic chamber was placed a solution of ArSSAr (Ar = *p*- FC_6H_4) (101.9 mg, 0.401 mmol) in 0.3 M $\text{Bu}_4\text{NBF}_4/\text{CH}_2\text{Cl}_2$ (8.0 mL). In the cathodic chamber were placed 0.3 M $\text{Bu}_4\text{NBF}_4/\text{CH}_2\text{Cl}_2$ (8.0 mL) and trifluoromethanesulfonic acid (41.0 mg, 0.273 mmol). The constant current electrolysis (8 mA) was carried out at -78°C with magnetic stirring until 0.67 F/mol of electricity was consumed. To the anodic chamber containing electrogenerated $\text{ArS}(\text{ArSSAr})^+$, was added 1-methoxy-1-phenylthiononane (**1**) (53.3 mg, 0.200 mmol) and the mixture was stirred for 5 min at -78°C . The solution thus obtained was used for the subsequent reaction.

4.8. NMR Analysis of Alkoxy-carbenium Ion based on Indirect Cation Pool Method.

4.8.1. Alkoxy-carbenium Ion (**2**) derived from **1**.

^1H and ^{13}C NMR spectra were recorded in $\text{CH}_2\text{Cl}_2/\text{CD}_2\text{Cl}_2$ (10:1) on a JEOL ECA600P (^1H 600 MHz, ^{13}C 150 MHz) spectrometer. The anodic oxidation was carried out in a divided cell equipped with a carbon felt anode and a platinum plate cathode. In the anodic chamber were placed a solution of ArSSAr (Ar = *p*- FC_6H_4) (81.9 mg, 0.322 mmol) in 4.0 mL of 0.3 M $\text{Bu}_4\text{NBF}_4/\text{CH}_2\text{Cl}_2/\text{CD}_2\text{Cl}_2$ (10:1). In the cathodic chamber were placed trifluoromethanesulfonic acid (32.4 mg, 0.216 mmol) and 0.3 M $\text{Bu}_4\text{NBF}_4/\text{CH}_2\text{Cl}_2/\text{CD}_2\text{Cl}_2$ (10:1) (4.0 mL). The constant

current electrolysis (4.0 mA) was carried out at -78°C with magnetic stirring. After 0.67 F/mol of electricity was consumed, the reaction mixture of the anodic chamber (0.8 mL) was transferred to a 5 mm ϕ NMR tube with a septum cap under Ar atmosphere at -78°C . 1-Methoxy-1-phenylthiononane (**1**) (10.2 mg, 0.0383 mmol) was added and the tube was shaken at the same temperature. The NMR measurement was carried out at -80°C . Chemical shifts are reported using methylene signals of CH_2Cl_2 at δ 5.32 (^1H NMR) and δ 53.80 (^{13}C NMR) as standards. The huge signal coming from CH_2Cl_2 is reduced by usual pulse techniques²⁸. ^1H NMR (600 MHz, $\text{CH}_2\text{Cl}_2/\text{CD}_2\text{Cl}_2$ (10:1), selected) δ 3.26 (t, J = 6.3 Hz, 2H), 4.92 (s, 3H), 9.53 (s, 1H); ^{13}C NMR (150 MHz, $\text{CH}_2\text{Cl}_2/\text{CD}_2\text{Cl}_2$ (10:1), selected) δ 40.3 (CH_3OCH_2), 75.6 (CH_3OCH), 230.6 (CH_3OCH). Other signals could not be assigned because of overlap with the signals of Bu_4NBF_4 used as the electrolyte.

4.8.2. Alkoxy-carbenium Ion derived from **5**.

The anodic oxidation was carried out in a divided cell equipped with a carbon felt anode and a platinum plate cathode. In the anodic chamber were placed a solution of ArSSAr (Ar = *p*- FC_6H_4) (153.0 mg, 0.602 mmol) in 4.0 mL of 0.3 M $\text{Bu}_4\text{NBF}_4/\text{CH}_2\text{Cl}_2/\text{CD}_2\text{Cl}_2$ (10:1). In the cathodic chamber were placed trifluoromethanesulfonic acid (80.2 mg, 0.534 mmol) and 0.3 M $\text{Bu}_4\text{NBF}_4/\text{CH}_2\text{Cl}_2/\text{CD}_2\text{Cl}_2$ (10:1) (4.0 mL). The constant current electrolysis (4.0 mA) was carried out at -78°C with magnetic stirring. After 0.67 F/mol of electricity was consumed, the reaction mixture of the anodic chamber (0.8 mL) was transferred to a 5 mm ϕ NMR tube equipped with a septum cap under Ar atmosphere at -78°C . 2-(phenylthio)tetrahydropyran (**5**) (16.4 mg, 0.084 mmol) was added and the tube was shaken at the same temperature. The NMR measurement was carried out at -80°C . Chemical shifts are reported using methylene signals of CH_2Cl_2 at δ 5.32 (^1H NMR) and δ 53.80 (^{13}C NMR) as standards. The huge signal coming from CH_2Cl_2 is reduced by usual pulse techniques²⁸. ^1H NMR (600 MHz, $\text{CH}_2\text{Cl}_2/\text{CD}_2\text{Cl}_2$ (10:1), selected) δ 1.88 (br t, J = 5.8 Hz, 2H), 2.14 (br s, 2H), 3.49 (br t, J = 5.2 Hz, 2H), 5.24 (br s, 2H), 9.86 (s, 1H); ^{13}C NMR (150 MHz, $\text{CH}_2\text{Cl}_2/\text{CD}_2\text{Cl}_2$ (10:1), selected) δ 12.1, 19.5, 35.4, 82.5, 227.8. Other signal could not be assigned because of overlap with the signals of Bu_4NBF_4 used as the electrolyte or CH_2Cl_2 , which was used as the solvent.

4.9. Reaction of Alkoxy-carbenium Ion and Carbon Nucleophiles. 4-Methoxy-dodec-1-ene (**3**) (A Typical Procedure of Indirect Cation Pool Method).

The electrolysis of ArSSAr (Ar = *p*- FC_6H_4) (102 mg, 0.401 mmol) was carried out as described above. To the $\text{ArS}(\text{ArSSAr})^+$ thus generated in the anodic chamber, was added 1-methoxy-1-phenylthiononane (**1**) (52.7 mg, 0.198 mmol) at -78°C and the reaction mixture was stirred for 5 min. Then allyltrimethylsilane (56.6 mg, 0.495 mmol) was added and the resulting solution was stirred at -78°C for 15 min. The reaction was quenched with Et_3N . The solvent was removed under reduced pressure and the residue was quickly filtered through a short column (2 x 3 cm) of silica gel to remove Bu_4NBF_4 . The silica gel was washed with ether (150 mL). The GC analysis of the combined filtrate indicated that the title compound was formed in 89 % yield (GC t_R 7.1 min, column, CBP1; 0.22 mm ϕ x 0.25 mm x 25 m; initial oven temperature, 100°C ; rate of temperature increase, $10^\circ\text{C}/\text{min}$). The isolated product was identified by the comparison of its spectral data with those of an authentic sample.^{7a}

The other products, **6**, **7^a**, **8**, **9**, **10**, **11**, **12**, **13**, **18^c** and **20^a** were identified by the comparison of their spectral data with those of authentic samples.

Acknowledgments.

This work was financially supported in part by a Grant-in-Aid for Scientific Research from the Japan Society for the Promotion of Science. The authors thank Prof. Dr. Hendrik Zipse of Ludwig-Maximilians-Universität München for fruitful discussions. K.M acknowledges JSPS for financial support.

References

- Current address: Division of Chemistry and Biochemistry, Graduate School of Natural Science and Technology, Okayama University, 3-1-1 Tsushima-naka, Kita-ku, Okayama 700-8530, Japan
- For example, Santelli, M.; Pons, J.-M. *Lewis Acids and Selectivity in Organic Synthesis*; CRC Press: Boca Raton, 1995; Chap 4.
- Benzylic alkoxy-carbenium ions: (a) Rabinovitz, M.; Bruck, D. *Tetrahedron Lett.* **1971**, 245-246. (b) Amyes, T. L.; Jencks, W. P. *J. Am. Chem. Soc.* **1989**, *111*, 7888-7900. (c) Jagannadham, V.; Amyes, T. L.; Richard, J. P. *J. Am. Chem. Soc.* **1993**, *115*, 8465-8466. (d) Mayr, H.; Gorath, G. *J. Am. Chem. Soc.* **1995**, *117*, 7862-7868. Di- and tri-alkoxy-carbenium ions: (e) Ramsey, B. G.; Taft, R. W. *J. Am. Chem. Soc.* **1966**, *88*, 3058-3063. (f) Steenken, S.; Buschek, J.; McClelland, R. A. *J. Am. Chem. Soc.* **1986**, *108*, 2808-2813. (g) McClelland, R. A.; Steenken, S. *J. Am. Chem. Soc.* **1988**, *110*, 5860-5866. (h) Steenken, S.; McClelland, R. A. *J. Am. Chem. Soc.* **1989**, *111*, 4967-4973.
- Denmark, S. E.; Willson, T. M. In *Selectivities in Lewis Acid Promoted Reactions*; Schinzer, D. Ed.; Kluwer Academic Publishers: Dordrecht, 1989; p 247.
- (a) Yoshida, J.; Kataoka, K.; Horcajada, R.; Nagaki, A. *Chem. Rev.* **2008**, *108*, 2265-2299. (b) Sperry, J. B.; Wright, D. L. *Chem. Soc. Rev.* **2006**, *35*, 605-621. (c) Lund, H. *J. Electrochem. Soc.* **2002**, *149*, S21-S33. (d) Moeller, K. D. *Tetrahedron* **2000**, *56*, 9527-9554.
- Yoshida, J.; Suga, S. *Chem. Eur. J.* **2002**, *8*, 2650-2658.
- (a) Suga, S.; Suzuki, S.; Yamamoto, A.; Yoshida, J. *J. Am. Chem. Soc.* **2000**, *122*, 10244-10245. (b) Suga, S.; Suzuki, S.; Yoshida, J. *Org. Lett.* **2005**, *6*, 4717-4720. (c) Okajima, M.; Suga, S.; Itami, K.; Yoshida, J. *J. Am. Chem. Soc.* **2005**, *127*, 6930-6931.
- (a) Yoshida, J.; Suga, S.; Suzuki, S.; Kinomura, N.; Yamamoto, A.; Fujiwara, K. *J. Am. Chem. Soc.* **1999**, *121*, 9546-9549. (b) Suga, S.; Okajima, M.; Yoshida, J. *Tetrahedron Lett.* **2001**, *42*, 2173-2176. (c) Suga, S.; Suzuki, S.; Yoshida, J. *J. Am. Chem. Soc.* **2002**, *124*, 30-31. (d) Suga, S.; Watanabe, M.; Yoshida, J. *J. Am. Chem. Soc.* **2002**, *124*, 14824-14825. (e) Suga, S.; Nagaki, A.; Yoshida, J. *Chem. Commun.* **2003**, 354-355. (f) Suga, S.; Nagaki, A.; Tsutsui, Y.; Yoshida, J. *Org. Lett.* **2003**, *5*, 945-947. (g) Suga, S.; Kageyama, Y.; Babu, G.; Itami, K.; Yoshida, J. *Org. Lett.* **2004**, *6*, 2709-2711. (h) Suga, S.; Suzuki, S.; Maruyama, T.; Yoshida, J. *Bull. Chem. Soc. Jpn.* **2004**, *77*, 1545-1554. (i) Suga, S.; Nishida, T.; Yamada, D.; Nagaki, A.; Yoshida, J. *J. Am. Chem. Soc.* **2004**, *126*, 14338-14339. (j) Nagaki, A.; Kawamura, K.; Suga, S.; Ando, T.; Sawamoto, M.; Yoshida, J. *J. Am. Chem. Soc.* **2004**, *126*, 14702-14703. (k) Suga, S.; Tsutsui, Y.; Nagaki, A.; Yoshida, J. *Bull. Chem. Soc. Jpn.* **2005**, *78*, 1206-1271. (l) Maruyama, T.; Suga, S.; Yoshida, J. *J. Am. Chem. Soc.* **2005**, *127*, 7324-7325. (m) Nagaki, A.; Togai, M.; Suga, S.; Aoki, N.; Mae, K.; Yoshida, J. *J. Am. Chem. Soc.* **2005**, *127*, 11666-11675. (n) Maruyama, T.; Suga, S.; Yoshida, J. *Tetrahedron* **2006**, *62*, 6519-6525. (o) Suga, S.; Watanabe, M.; Song, C.-H.; Yoshida, J. *Electrochemistry* **2006**, *74*, 672-679. (p) Maruyama, T.; Mizuno, Y.; Shimizu, I.; Suga, S.; Yoshida, J. *J. Am. Chem. Soc.* **2007**, *129*, 1902-1903. (q) Suga, S.; Shimizu, I.; Ashikari, Y.; Mizuno, Y.; Maruyama, T.; Yoshida, J. *Chem. Lett.* **2008**, *37*, 1008-1009.
- (a) Okajima, M.; Soga, K.; Nokami, T.; Suga, S.; Yoshida, J. *Org. Lett.* **2006**, *8*, 5005-5007. (b) Nokami, T.; Ohata, K.; Inoue, M.; Tsuyama, H.; Shibuya, A.; Soga, K.; Okajima, M.; Suga, S.; Yoshida, J. *J. Am. Chem. Soc.* **2008**, *130*, 10864-10865. (c) Okajima, M.; Soga, K.; Watanabe, T.; Terao, K.; Nokami, T.; Suga, S.; Yoshida, J. *Bull. Chem. Soc. Jpn.* **2009**, *82*, 594-599.
- Nokami, T.; Soma, R.; Yamamoto, Y.; Kamei, T.; Itami, K.; Yoshida, J. *Beilstein J. Org. Chem.* **2007**, *3*, 7.
- Midorikawa, K.; Suga, S.; Yoshida, J. *Chem. Commun.* **2006**, 3794-3796.
- (a) Yoshida, J.; Maekawa, T.; Murata, T.; Matsunaga, S.; Isoe, S. *J. Am. Chem. Soc.* **1990**, *112*, 1962-1970. (b) Yoshida, J.; Nishiwaki, K. *J. Chem. Soc., Dalton Trans.* **1998**, 2589-2596.
- (a) Olah, G. A.; Bollinger, J. M. *J. Am. Chem. Soc.* **1967**, *89*, 2993-2996. (b) Olah, G. A.; Sommer, J. *J. Am. Chem. Soc.* **1968**, *90*, 4323-4327. (c) Forsyth, D. A.; Osterman, V. M.; DeMember, J. R. *J. Am. Chem. Soc.* **1985**, *107*, 818-822.
- (a) Yoshida, J.; Sugawara, M.; Kise, N. *Tetrahedron Lett.* **1996**, *37*, 3157-3160. (b) Sugawara, M.; Mori, K.; Yoshida, J. *Electrochim. Acta*, **1997**, *42*, 1995-2003. (c) Yoshida, J.; Sugawara, M.; Tatsumi, M.; Kise, N. *J. Org. Chem.* **1998**, *63*, 5950-5961.
- (a) Suzuki, S.; Matsumoto, K.; Kawamura, K.; Suga, S.; Yoshida, J. *Org. Lett.* **2004**, *6*, 3755-3758. (b) Suga, S.; Matsumoto, K.; Ueoka, K.; Yoshida, J. *J. Am. Chem. Soc.* **2006**, *128*, 7710-7711.
- To solve the problem, we have developed the cation flow method. (a) Suga, S.; Okajima, M.; Fujiwara, K.; Yoshida, J. *J. Am. Chem. Soc.* **2001**, *123*, 7941-7942. (b) Suga, S.; Okajima, M.; Fujiwara, K.; Yoshida, J. *QSAR Comb. Sci.* **2005**, *24*, 728-741.
- For indirect electrochemical methods, see: Steckhan, E. *Angew. Chem., Int. Ed. Engl.* **1986**, *25*, 683-701.
- (a) Bewick, A.; Coe, D. E.; Mellor, J. M.; Walton, D. J. *J. Chem. Soc., Chem. Commun.* **1980**, 51-52. (b) Do, Q. T.; Elthmani, D.; Simonet, J.; Le Guillanton, G. *Electrochim. Acta* **2005**, *50*, 4792-4799, and references cited therein.
- For example, (a) Smit, W. A.; Krimer, M. Z.; Vorobieva, E. A. *Tetrahedron Lett.* **1975**, *16*, 2451-2454. (b) Capozzi, G.; Lucchini, V.; Modena, G.; Rivetti, F. *J. Chem. Soc., Perkin Trans. 2* **1975**, 361-366.
- For example, (a) Gybin, A. S.; Smit, W. A.; Bogdanov, V. S.; Krimer, M. Z.; Kalyan, J. B. *Tetrahedron Lett.* **1980**, *21*, 383-386. (b) Bogdanov, V. S.; Gybin, A. S.; Cherepanova, E. G.; Smith, W. A. *Izv. Akad. Nauk Ser. Khim.* **1981**, 2681-2693.
- (a) Larsen, A. G.; Holm, A. H.; Roberson, M.; Daasbjerg, K. *J. Am. Chem. Soc.* **2001**, *123*, 1723-1729. (b) Brinck, T.; Carlqvist, P.; Holm, A. H.; Daasbjerg, K. *J. Phys. Chem. A* **2002**, *106*, 8827-8833. (c) Holm, A. H.; Yusta, L.; Carlqvist, P.; Brinck, T.; Daasbjerg, K. *J. Am. Chem. Soc.* **2003**, *123*, 2148-2157.
- Reactivity of cations: (a) Mayr, H.; Ofial, A. R. *Angew. Chem., Int. Ed.* **2006**, *45*, 1844-1854. (b) Hofmann, M.; Hampel, N.; Kanzian, T.; Mayr, H. *Angew. Chem., Int. Ed.* **2004**, *43*, 5402-5405.

23. (a) Braga, A. L.; Dornelles, L.; Silveira, C. C.; Wessjohann, L. A. *Synthesis* **1999**, 562-564. (b) Sato, T.; Okura, S.; Otera, J.; Nozaki, H. *Tetrahedron Lett.* **1987**, 28, 6299-6302. (c) Nishiyama, H.; Narimatsu, S.; Sakuta, K.; Itoh, K. *J. Chem. Soc., Chem. Commun.* **1982**, 459-460.
24. Leino, R.; Lönnqvist, J. E. *Tetrahedron Lett.* **2004**, 45, 8489-8491.
25. Becker, D. P.; Villamil, C. I.; Barta, T. E.; Bedell, L. J.; Boehm, T. L.; DeCrescenzo, G. A.; Freskos, J. N.; Getman, D. P.; Hockerman, S.; Heintz, R.; Howard, S. C.; Li, M. H.; McDonald, J. J.; Carron, C. P.; Funckes-Shippy, C. L.; Mehta, P. P.; Munie, G. E.; Swearingen, C. A. *J. Med. Chem.* **2005**, 48, 6713-6730.
26. Braga, A. L.; Silveira, C. C.; Dornelles, L.; Zeni, G.; Galarza, F. A. D.; Wessjohann, L. A. *Syn. Commun.* **1995**, 25, 3155-3162.
27. Sato, T.; Otera, J.; Nozaki, H. *Tetrahedron* **1989**, 45, 1209-1218.
28. Guéron, M.; Plateau, P.; Decors, M. *Prog. Nucl. Magn. Reson. Spectrosc.* **1991**, 23, 135-209.