

1 **Increased visceral fat mass and insulin signaling in colitis-related colon**
2 **carcinogenesis model mice**

3

4 **Shingo Miyamoto^a, Takuji Tanaka^b, and Akira Murakami^{a,*}**

5

6 **^aDivision of Food Science and Biotechnology, Graduate School of Agriculture,**
7 **Kyoto University, Kyoto 606-8502, Japan**

8 **^bDepartment of Oncologic Pathology, Kanazawa Medical University, 1-1 Daigaku,**
9 **Uchinada, Ishikawa 920-0293, Japan**

10 ***Corresponding author. Fax: +81-75-753-6284; E-mail: cancer@kais.kyoto-u.ac.jp**

11

12

13 **Abbreviations:** Akt, protein kinase B; AOM, azoxymethane; CD, Crohn's disease;
14 CRC, colorectal cancer; DSS, dextran sulfate sodium; ERK, extracellular
15 signal-regulated protein kinase; IBD, inflammatory bowel disease; IL-6, interleukin-6;
16 mTOR, mammalian target of rapamycin; Ob-R, leptin receptor; TNF- α , tumor necrosis
17 factor- α

1 Abstract

2

3 Leptin, a pleiotropic hormone regulating food intake and metabolism, plays an
4 important role in the regulation of inflammation and immunity. We previously
5 demonstrated that serum leptin levels are profoundly increased in mice which received
6 azoxymethane (AOM) and dextran sulfate sodium (DSS) as tumor-initiator and
7 -promoter, respectively, in a colon carcinogenesis model. In this study, we attempted to
8 address underlying mechanism whereby leptin is up-regulated in this rodent model.
9 Five-week-old male ICR mice were given a single intraperitoneal injection of AOM
10 (week 0), followed by 1% DSS in drinking water for 7 days. Thereafter, the weights of
11 visceral fats and the serum concentration of leptin were determined at week 20. Of
12 interest, the relative epididymal fat pad and mesenteric fat weights, together with serum
13 leptin levels in the AOM and/or DSS-treated mice were markedly increased compared
14 to that in untreated mice. In addition, leptin protein production in epididymal fat pad
15 with AOM/DSS-treated mice was 4.7-fold higher than that of control. Further, insulin
16 signaling molecules, such as protein kinase B (Akt), S6, mitogen-activate protein
17 kinase/ extracellular signaling-regulated kinase 1/2, and extracellular signaling-
18 regulated kinase 1/2, were concomitantly activated in epididymal fat of AOM/DSS-
19 treated mice. This treatment also increased the serum insulin and IGF-1 levels. Taken
20 together, our results suggest that higher levels of serum insulin and IGF-1 promote the
21 insulin signaling in epididymal fat and thereby increasing serum leptin, which may play
22 an crucial role in, not only obesity-related, but also -independent colon carcinogenesis.

23

24 **Key words:** leptin, colon carcinogenesis, epididymal fat pad, insulin signaling

1 **Introduction**

2

3 Leptin, a 16-kDa protein encoded by the *ob* gene, was discovered in 1994 and
4 shown to be a regulator of body weight and energy balance, functioning in the
5 hypothalamus [1]. Although this hormone is produced mainly by WAT, its expression
6 has also been detected in other tissues, including the gastrointestinal tract [2]. In humans,
7 leptin levels are proportional to body mass index and are elevated in obese individuals
8 [3]. Since obesity is known to increase the risk of certain cancers, great effort has been
9 directed to elucidating the possible role of leptin in cancer development [4], and several
10 *in vitro* studies showed that leptin can act as a mitogenic, anti-apoptotic, and
11 tumorigenic factor with various cancer cell lines [5-7]. In addition, data obtained with
12 clinical samples revealed that leptin and its receptor (Ob-R) are expressed in cancer
13 tissues [8]. Thus, the leptin system might promote cancer progression in an autocrine
14 and paracrine manner.

15 Our previous investigation demonstrated that serum leptin levels were profoundly
16 increased in male ICR mice that received azoxymethane (AOM) and dextran sulfate
17 sodium (DSS) for colitis-related colon carcinogenesis [9]. In addition, a citrus flavonoid,
18 nobiletin, reduced not only those levels, but also colonic tumor development, whereas
19 did not cause significant alterations in serum levels of other adipocytokines (TNF- α ,
20 IL-6, and adiponectin) and triglyceride [9]. These findings led us to address and
21 investigate an issue why and how serum leptin level is elevated in the mice treated with
22 AOM and DSS in a colitis-related colon carcinogenesis model.

23

24 **Materials and Methods**

1

2 *Animals and chemicals*

3 Male Crj:CD-1 (ICR) mice (Japan SLC, Shizuoka, Japan) aged 5 weeks were
4 obtained for this study. The animals were handled according to the Guidelines for the
5 Regulation of Animals provided by the Experimentation Committee of Kyoto
6 University and the experimental protocol was approved by the Committee. Upon arrival,
7 all mice were randomized and transferred to plastic cages (5 mice/cage), with free
8 access to drinking water and a pelleted basal diet MF diet (Oriental Yeast, Kyoto, Japan),
9 and controlled conditions of humidity ($60 \pm 5\%$), light (12/12-hour light/dark cycle) and
10 temperature ($24 \pm 2^\circ\text{C}$). All mice were quarantined for 1 week before starting the
11 experiments.

12 The colonic carcinogen AOM was purchased from Sigma Chemical Co. (St. Louis,
13 MO). DSS with a molecular weight of 36,000 to 50,000 (Cat. No. 160110) was
14 purchased from the MP Biochemicals, LLC (Aurora, OH) and used for induction of
15 colitis, after being dissolved in distilled water to a concentration of 1% (w/v).

16

17 *Antibodies*

18 Antibodies directed against phospho-protein kinase B (Akt) (Ser483), phospho-S6
19 (Ser240/244), phospho-mitogen-activate protein kinase/ extracellular
20 signaling-regulated kinase (MEK)1/2 (Ser217/221) and phospho- extracellular
21 signaling-regulated kinase (ERK)1/2 (Thr202/Tyr204), as well as horseradish
22 peroxidase (HRP)-conjugated anti-rabbit antibody were obtained from Cell Signaling
23 Technology (Beverly, MA).

24

1 *Animal treatment*

2 A total of 80 male ICR mice were divided into four (3 experimental and 1 control)
3 groups. Group 1 served as an untreated control. Mice in groups 2 and 4 were given a
4 single i.p. injection of AOM (10 mg/kg body weight), then starting 1 week after that
5 injection, those in group 4 received 1% DSS in drinking water for 7 days. Mice on
6 group 3 received DSS alone according to the same time schedule. All groups were fed
7 the basal diet throughout the study. At the end of weeks 2, 5 mice from each group were
8 euthanized, then 15 mice from each group were euthanized after week 20. Euthanasia
9 was performed under ether anesthesia, then blood samples were collected from the
10 abdominal aorta. During the study, body weights of all animals were weighted weekly.
11 At sacrifice, all organs, including adipose tissue, were removed and weighted, and then
12 obtained adipose tissues were frozen using liquid nitrogen, until later use.

13

14 *Reverse transcription-polymerase chain reaction (RT-PCR)*

15 Total RNA was extracted from the epididymal fat pads using TRIzol reagent
16 (Invitrogen, Carlsbad, CA), according to the manufacturer's instructions. cDNA was
17 synthesized using 1 µg of total RNA with an RNA PCR Kit (AMV). PCR amplification
18 was performed using a thermal cycler (PTC-100TM, MJ Research, Watertown, MA)
19 and conducted with each sense and antisense primer, with *GAPDH* used as the internal
20 standard. The primers for *leptin* were 5'-CCAAAACCCTCATCAAGACC-3' and 5'-
21 GTCCAACCTGTTGAAGAATGTCCC-3' (Proligo, Kyoto, Japan), which yielded an
22 expected band size of 395 bp, while those for *GAPDH* were 5'-GTGAAGGTCGGA-
23 GTCAACG-3' and 5'-GGTGAAGACGCCAGTGGACTC-3' (Proligo), which yielded
24 an expected band size of 300 bp. The primers were used at final concentrations of 0.5

1 and 0.05 μM respectively. PCR was performed under the following conditions: 43
2 cycles at 95°C for 45 seconds, 57°C for 45 seconds, and 72°C for 45 seconds for *leptin*,
3 and 27 cycles at 94°C for 30 seconds, 55°C for 30 seconds, and 72°C for 30 seconds for
4 *GAPDH*. The PCR products were subjected to electrophoresis on 3% agarose gels and
5 stained with SYBR[®] Gold. The intensity of each band was analyzed using Scion
6 Image.

7

8 *Western blotting*

9 To determine the protein expressions of phospho-Akt, phospho-S6,
10 phospho-MEK1/2, and phospho-ERK1/2 in the epididymal fat pads, tissue supernatants
11 were subjected to Western blotting. Frozen adipose tissue was lysed in lysis buffer [10
12 nM Tris, pH 7.4, 1% sodium dodecyl sulfate (SDS), 1 mM sodium metavanadate (V)],
13 and centrifuged at 3200 \times g for 5 minutes. Denatured proteins (40 μg) were separated
14 using SDS-PAGE on a 10% polyacrylamide gel and then transferred to Immobilon-P
15 membranes (Millipore, Billerica, MA). After blocking with Block Ace (Snow Brand
16 Milk Products, Tokyo, Japan) for 1 hour, the membranes were reacted with the
17 corresponding specific primary antibody (1:1000), followed by the corresponding
18 HRP-conjugated secondary antibody (1:1000). The blots were developed using ECL
19 Western blotting detection reagents. The intensity of each band was analyzed using
20 Scion Image.

21

22 *Clinical chemistry*

23 The collected blood samples were used for clinical chemistry measurements of
24 leptin (Quantikine Mouse leptin, ELISA/Assay Kit, R&D Systems Inc.), insulin (Insulin

1 measurement kit, Morinaga Institute of Biological Science), and insulin-like growth
2 factor (IGF)-1 (Quantikine Mouse IGF-1, ELISA/Assay Kit, R&D Systems Inc.). The
3 samples were used without dilution for insulin measurements, while they were diluted
4 20- and 500-fold for leptin and IGF-1.

5

6 *Statistical analysis*

7 Data were analyzed using Student's *t*-test (two-sided) when appropriate, with $P <$
8 0.05 considered to indicate significance.

9

10 **Results**

11

12 *General observations of mice*

13 Food intake was significantly ($P < 0.05$) lower in group 4 as compared to group 1
14 (Figure 1A). The mean body weights of mice in groups 2 through 4 are higher than
15 group 1, though the differences did not reach statistical significance (Figure 1B). In
16 addition, the AOM and/or DSS treatment led to notable increases in mean relative
17 epididymal fat pad weight (g/100 g body weight), which was most notable in group 4
18 (AOM and DSS), as the weight was significantly ($P < 0.05$) increased by 1.7-fold as
19 compared to group 1 (Figure 1C). However, the mean relative mesenteric fat weight
20 (g/100 g body weight) did not significantly differ among the groups (Figure 1D).

21

22 *Leptin level in the serum and epididymal fat pad*

23 As shown in Figure 2, the serum concentration of leptin in group 4 was significantly
24 ($P < 0.05$) elevated by 3.0-fold as compared to group 1, which is consistent with our

1 previous findings [9]. The serum leptin levels in groups 2 and 3 were also increased by
2 1.8- and 2.2-fold as compared to group 1, respectively, without statistical significance.
3 Since leptin is produced mainly by WAT, we assessed the level of leptin mRNA and
4 protein expression in epididymal fat pads. Leptin mRNA expression in the mice treated
5 with both AOM and DSS (group 4) was increased by 2.0-fold as compared to the
6 untreated control (Figure 3A), while leptin protein production was also increased by
7 4.7-fold ($P < 0.05$) in group 4 mice (Figure 3B).

8

9 *Activation of insulin signaling pathway in epididymal fat pads*

10 Insulin is an important physiological factor that activates mammalian target of
11 rapamycin (mTOR), which is a master regulator of protein synthesis [10], adipose tissue
12 morphogenesis [11], and leptin synthesis/secretion [12] in adipocytes. To determine
13 whether the insulin signaling pathway was activated in the epididymal fat pads of the
14 present ICR mice, we determined the phosphorylation status of Akt, S6, MEK1/2, and
15 ERK1/2, all of which are involved in that pathway [13]. As shown in Figure 3, AOM
16 and DSS treatment substantially increased the phosphorylation of Akt (3.5-fold, $P <$
17 0.05 , Figure 3C), S6 (38-fold, $P < 0.05$, Figure 3D), MEK (5.4-fold, $P < 0.05$, Figure
18 3E), and ERK (8.8-fold, $P < 0.05$, Figure 3F), respectively, as compared to the untreated
19 control mice.

20

21 *Serum levels of insulin and IGF-1*

22 We also assessed the serum levels of insulin and IGF-1 at two time-points, week 2
23 (just after the cessation of the DSS treatment) and 20 (the termination of the study). As
24 shown in Figure 4A, serum insulin was elevated in a time-dependent manner by 2.1-

1 and 7.8-fold in groups 1 and 4, respectively. Meanwhile, the level of IGF-1 in group 4
2 was significantly ($P < 0.05$) higher at week 20 than that in group 1, which was in
3 contrast to the findings at week 2 (Figure 4B).

4

5 **Discussion**

6

7 Colorectal cancer (CRC) is the most serious complication associated with long
8 standing inflammatory bowel disease (IBD), which includes ulcerative colitis and
9 Crohn's disease (CD) [14, 15]. Mesenteric WAT is mainly hypertrophic around the
10 intestines in IBD patients, with axial polarity of inflammation the mainstay contributor
11 [16]. To better understand the pathogenesis of IBD-related CRC, one (T.T.) of the
12 present authors and colleagues developed a novel colitis-related CRC model (a
13 two-stage mouse colon carcinogenesis model), which were initiated with AOM and
14 promoted by DSS [17]. In the current study, such sequential treatment significantly
15 increased serum leptin and epididymal fat pad weight, whereas mesenteric fat was not
16 altered (Figure 1C and D). The individual potencies were less than their combination,
17 suggesting that AOM and DSS act additively to increase the concentration of leptin in
18 serum.

19 Bendet et al. reported that insulin secretion during a 3-hour oral glucose-tolerance
20 test was increased and homeostatic model assessment values were decreased in CD
21 patients [18]. In a later study, Bregenzer et al. found increased insulin resistance and β
22 cell function in CD patients [19]. Furthermore, it should be noted that the concentration
23 of serum C-peptide, a stable indicator of steady-state insulin secretion, was reported to
24 be higher in AOM-treated male rats [20]. Since insulin resistance increases the level of

1 serum IGF-1 [21], the increased level of IGF-1 in serum of the present group 4 mice
2 (Figure 4B) may have been associated with putative insulin resistance caused by
3 AOM/DSS-induced inflammation with other pathologic conditions, including oxidative
4 stress.

5 Under physiological conditions, serum leptin levels are affected by several factors,
6 and leptin secretion from adipocytes is dominantly dependent on the status of adipocyte
7 signaling molecules. A previous study showed that both insulin and IGF-1 have been
8 reported to be important physiological factors activating mTOR. Activation of this
9 pathway mediates multiple insulin effects, such as stimulation of adipogenesis or
10 lipogenesis [22], and potentiation of leptin secretion. In the present study, the
11 insulin/IGF signaling pathway was consistently activated, and leptin induction and
12 production were promoted in the epididymal fat of mice treated with AOM and DSS
13 (Figure 3A-F). Thus, it is suggested that these hormones stimulate growth and
14 differentiation of visceral fat tissue, thereby accelerating leptin production.

15 In conclusion, we found that higher levels of serum insulin and IGF-1 promote
16 insulin signaling in epididymal fat, which may lead to increased amounts of visceral fat
17 tissue and a resulting elevation of serum leptin. These events may play certain critical
18 roles in the development of CRC. However, further studies are warranted for
19 understanding the precise mechanisms underlying AOM and DSS-induced adipose
20 tissue alterations.

21

22 **Acknowledgements**

23

24 This study was supported in part by a Grant-in-Aid for Cancer Research from the

- 1 Ministry of Health, Labor and Welfare of Japan (A.M. and T.T.), a Grant-in Aid for
- 2 Scientific Research (C) (A.M.), and the Japan Society for the Promotion of Science
- 3 Fellows (S.M.).
- 4

1 **References**

- 2
- 3 [1] Y. Zhang, R. Proenca, M. Maffei, M. Barone, L. Leopold, J. Friedman,
4 Positional cloning of the mouse obese gene and its human homologue., *Nature*
5 372(6505) (1994) 425-432.
- 6 [2] S. Sitaraman, X. Liu, L. Charrier, L. Gu, T. Ziegler, A. Gewirtz, D. Merlin,
7 Colonic leptin: source of a novel proinflammatory cytokine involved in IBD., *FASEB J*
8 18(6) (2004) 696-698.
- 9 [3] J. Kolaczynski, J. Ohannesian, R. Considine, C. Marco, J. Caro, Response of
10 leptin to short-term and prolonged overfeeding in humans., *J Clin Endocrinol Metab*
11 81(11) (1996) 4162-4165.
- 12 [4] C. Garofalo, E. Surmacz, Leptin and cancer., *J Cell Physiol* 207(1) (2006)
13 12-22.
- 14 [5] J. Hardwick, G. Van Den Brink, G. Offerhaus, S. Van Deventer, M.
15 Peppelenbosch, Leptin is a growth factor for colonic epithelial cells., *Gastroenterology*
16 121(1) (2001) 79-90.
- 17 [6] P. Rouet-Benzineb, T. Aparicio, S. Guilmeau, C. Pouzet, V. Descatoire, M.
18 Buyse, A. Bado, Leptin counteracts sodium butyrate-induced apoptosis in human colon
19 cancer HT-29 cells via NF-kappaB signaling., *J Biol Chem* 279(16) (2004)
20 16495-16502.
- 21 [7] O. Ogunwobi, I. Beales, The anti-apoptotic and growth stimulatory actions of
22 leptin in human colon cancer cells involves activation of JNK mitogen activated protein
23 kinase, JAK2 and PI3 kinase/Akt., *Int J Colorectal Dis* 22(4) (2007) 401-409.
- 24 [8] M. Koda, M. Sulkowska, L. Kanczuga-Koda, E. Surmacz, S. Sulkowski,
25 Overexpression of the obesity hormone leptin in human colorectal cancer., *J Clin Pathol*
26 60(8) (2007) 902-906.
- 27 [9] S. Miyamoto, Y. Yasui, T. Tanaka, H. Ohigashi, A. Murakami, Suppressive
28 effects of nobiletin on hyperleptinemia and colitis-related colon carcinogenesis in male
29 ICR mice., *Carcinogenesis* 29(5) (2008) 1057-1063.
- 30 [10] C. Lynch, B. Patson, J. Anthony, A. Vaval, L. Jefferson, T. Vary, Leucine is a
31 direct-acting nutrient signal that regulates protein synthesis in adipose tissue., *Am J*
32 *Physiol Endocrinol Metab* 283(3) (2002) E503-513.
- 33 [11] H. Fox, S. Kimball, L. Jefferson, C. Lynch, Amino acids stimulate
34 phosphorylation of p70S6k and organization of rat adipocytes into multicellular
35 clusters., *Am J Physiol* 274(1 Pt 1) (1998) C206-213.
- 36 [12] C. Roh, J. Han, A. Tzatsos, K. Kandror, Nutrient-sensing mTOR-mediated

- 1 pathway regulates leptin production in isolated rat adipocytes., *Am J Physiol Endocrinol*
2 *Metab* 284(2) (2003) E322-330.
- 3 [13] R. Bradley, B. Cheatham, Regulation of ob gene expression and leptin
4 secretion by insulin and dexamethasone in rat adipocytes., *Diabetes* 48(2) (1999)
5 272-278.
- 6 [14] J. Eaden, K. Abrams, J. Mayberry, The risk of colorectal cancer in ulcerative
7 colitis: a meta-analysis., *Gut* 48(4) (2001) 526-535.
- 8 [15] R. van Hogezaand, R. Eichhorn, A. Choudry, R. Veenendaal, C. Lamers,
9 Malignancies in inflammatory bowel disease: fact or fiction?, *Scand J Gastroenterol*
10 *Suppl*(236) (2002) 48-53.
- 11 [16] K. Karmiris, I. Koutroubakis, E. Kouroumalis, Leptin, adiponectin, resistin,
12 and ghrelin--implications for inflammatory bowel disease., *Mol Nutr Food Res* 52(8)
13 (2008) 855-866.
- 14 [17] T. Tanaka, H. Kohno, R. Suzuki, Y. Yamada, S. Sugie, H. Mori, A novel
15 inflammation-related mouse colon carcinogenesis model induced by azoxymethane and
16 dextran sodium sulfate., *Cancer Sci* 94(11) (2003) 965-973.
- 17 [18] N. Bendet, E. Scapa, O. Cohen, O. Bloch, D. Aharoni, Y. Ramot, M. Weiss, A.
18 Halevi, M. Rapoport, Enhanced glucose-dependent glucagon-like peptide-1 and insulin
19 secretion in Crohn patients with terminal ileum disease is unrelated to disease activity or
20 ileal resection., *Scand J Gastroenterol* 39(7) (2004) 650-656.
- 21 [19] N. Bregenzler, A. Hartmann, U. Strauch, J. Schölmerich, T. Andus, L.
22 Bollheimer, Increased insulin resistance and beta cell activity in patients with Crohn's
23 disease., *Inflamm Bowel Dis* 12(1) (2006) 53-56.
- 24 [20] R. Xiao, J. Carter, A. Linz, M. Ferguson, T. Badger, F. Simmen, Dietary whey
25 protein lowers serum C-peptide concentration and duodenal SREBP-1c mRNA
26 abundance, and reduces occurrence of duodenal tumors and colon aberrant crypt foci in
27 azoxymethane-treated male rats., *J Nutr Biochem* 17(9) (2006) 626-634.
- 28 [21] E. Giovannucci, Nutrition, insulin, insulin-like growth factors and cancer.,
29 *Horm Metab Res* 35(11-12) (2003) 694-704.
- 30 [22] H. Cho, J. Park, H. Lee, Y. Lee, J. Kim, Regulation of adipocyte differentiation
31 and insulin action with rapamycin., *Biochem Biophys Res Commun* 321(4) (2004)
32 942-948.

33

34

35

1 **Figure legends**

2

3 Figure 1. The mean food intake (A) by the mice during the study, body (B), epididymal
4 fat pad (C), and mesenteric fat (D) weights. The weights were measured at the
5 end of the study.

6

7 Figure 2. Serum leptin levels of all groups. The levels were quantified by ELISA, as
8 described in Materials and Methods. Each values is shown as the mean \pm SD
9 (n=15). Statistical analysis was performed using Student's *t*-test: **P* < 0.05 vs.
10 the untreated control (group 1).

11

12 Figure 3. Leptin production and activation of insulin signaling pathway in epididymal
13 fat pads at the end of the study. (A) Leptin mRNA expression was determined by
14 RT-PCR, as described in Materials and Methods, with the levels corrected by
15 those of GAPDH. Each values is shown as the mean \pm SD (n=5), with 3
16 representative results presented. (B) Leptin protein amounts were quantified by
17 ELISA, as described in Materials and Methods. Each values is shown as the
18 mean \pm SD (n=10). (C-F) Activation of the insulin signaling pathway was
19 examined by Western blotting, as described in Materials and Methods. Each
20 values is shown as the mean \pm SD (n=10), with 3 representative results
21 presented. Statistical analysis was performed using Student's *t*-test: **P* < 0.05 vs.
22 the untretated control (group 1).


23

24 Figure 4. Levels of serum insulin (A) and IGF-1 (B) after weeks 2 and 20. Serum


1 insulin and IGF-1 levels were quantified by ELISA, as described in Materials
2 and Methods. Open bars; at week 2, closed bars; at week 20. Each values is
3 shown as the mean \pm SD (n=5 or 15). Statistical analysis was performed using
4 Student's *t*-test: **P* < 0.05 vs. the untreated control (group 1) for the
5 corresponding time point.

Fig. 1 (Miyamoto et al.)


A



B



C



D




Fig. 2 (Miyamoto et al.)




Fig. 3 (Miyamoto et al.)




Fig. 4 (Miyamoto et al.)

