TITLE:

LONG TIME TAILS IN DIFFUSION-CONTROLLED RECOMBINATIONS

AUTHOR(S):

Ohtsuki, T.

CITATION:

Ohtsuki, T.. LONG TIME TAILS IN DIFFUSION-CONTROLLED RECOMBINATIONS. 物性研究 1991, 57(3): 396-396

ISSUE DATE:

1991-12-20

URL:

<http://hdl.handle.net/2433/94844>

RIGHT:

LONG TIME TAILS IN DIFFUSION-CONTROLLED RECOMBINATIONS

T. Ohtsuki

Department of Applied Physics, Fukui University, Fukui 910,
Japan

Abstract. Recently, much attention has been paid to slow asymptotic relaxation in simple diffusion-controlled recombination processes. Below an upper critical dimension d_c , particle densities $\rho(t)$ exhibit nonanalytic decay in the long time limit $t \rightarrow \infty$. Typical examples are

$$\rho_A(t) \propto t^{-d/2} \quad (d < d_c = 2), \quad (1)$$

$$\rho_A(t) = \rho_B(t) \propto t^{-d/4} \quad (d < d_c = 4), \quad (2)$$

$$\rho_A(t) \propto \exp[-t^{d/2}] \quad (d < d_c = 2). \quad (3)$$

These behaviors are investigated on the basis of a field theoretical renormalization group technique where initial conditions are taken into account explicitly. It becomes evident that the method is a powerful tool in understanding these phenomena systematically. Complete scaling relations are derived and concrete forms of scaling functions are calculated. Logarithmic corrections at $d = d_c$ are also discussed.