

Characteristic classes of surface bundles

By SHIGEYUKI MORITA

東大教養 森田茂之

In this paper we define characteristic classes of surface bundles, namely smooth fibre bundles whose fibres are a closed orientable surface Σ_g of genus $g \geq 2$ and announce some non-triviality results for them. As a consequence we obtain lower bounds for the Betti numbers of the mapping class group $M(g)$ of Σ_g .

It is known [EE] that the connected component of the identity of $\text{Diff}_+\Sigma_g$, the group of orientation preserving diffeomorphisms of Σ_g , is contractible. Therefore $B\text{Diff}_+\Sigma_g$ is a $K(M(g), 1)$. Now let ξ be the tangent bundle along the fibres of an oriented surface bundle and let $e(\xi)$ be its Euler class. If we apply the Gysin homomorphism to $e^{i+1}(\xi)$, we obtain an integral cohomology class of the base space of degree $2i$. By the naturality, this defines certain cohomology classes

$$e_i \in H^{2i}(M(g); \mathbb{Z}) \quad (i=1, 2, \dots).$$

$M(g)$ acts on $H^1(\Sigma_g; \mathbb{Z})$ preserving the symplectic form given by the cup product and so we obtain a homomorphism $M(g) \longrightarrow \text{Sp}(2g; \mathbb{Z})$, where $\text{Sp}(2g; \mathbb{Z})$ is the group of all $2g \times 2g$ symplectic matrices with integral entries. This induces a homomorphism $M(g) \longrightarrow \text{Sp}(2g; \mathbb{R})$. Since $\text{Sp}(2g; \mathbb{R})$ has $U(g)$ as a maximal compact subgroup, we have a g -dimensional complex vector bundle η on $K(M(g), 1)$. Let $c_i(\eta) \in H^{2i}(M(g); \mathbb{Z})$ be its i -th Chern class. From the argument of Atiyah in [A] and the fact that η is

1980 Mathematics Subject Classification. Primary 55R40, 57N05; Secondary 32G15.

flat as a real vector bundle, we can conclude

$$e_{2i-1} = \frac{(-1)^i}{B_i} s_{2i-1}(c(\eta)) \quad (i=1,2,\dots \text{ and coefficients are in } \mathbb{Q})$$

$$s_{2i}(c(\eta)) = 0$$

where $s_i(c(\eta))$ stands for the characteristic class of η corresponding to the formal sum $\sum_j t_j^i$ and B_i is the i -th Bernoulli number. These two relations induce those among monomials of e_{2i-1} 's and the quotient $\mathbb{Q}[e_1, e_3, \dots]/(\text{relations})$ is naturally isomorphic to the relative Lie algebra cohomology $H^*(\mathfrak{sp}(2g; \mathbb{R}), u(g))$ which in turn is additively isomorphic to $H^*(S^2 \times S^4 \times \dots \times S^{2g}; \mathbb{Q})$ (see [BH]). It is known that $M(g)$ acts properly discontinuously on the Teichmüller space $T(g) \cong \mathbb{R}^{6g-6}$ with non-compact quotient \mathcal{M}_g , the moduli space for Riemann surfaces of genus g . Hence $\text{vcd}(M(g)) \leq 6g-7$. Thus any monomial of e_i 's of degree $\geq 6g-6$ vanishes. To sum up we have a homomorphism

$$\phi: \mathbb{Q}[e_1, e_2, \dots]/(\text{above relations}) \longrightarrow H^*(M(g); \mathbb{Q}),$$

here we use the letter e_i for both symbolical and actual meanings. Since $\text{vcd}(M(g))$ is conjectured to be $3g-3$ ([Hv]), ϕ will surely have a still large kernel. Our main results are

THEOREM 1. For any $k \in \mathbb{N}$, there exists a natural number $g(k)$ such that the elements e_1, \dots, e_k are all non-trivial in $H^*(M(g); \mathbb{Q})$ if $g \geq g(k)$.

COROLLARY 2. The natural surjective homomorphism $\text{Diff}_{+\Sigma_g} \longrightarrow M(g)$ does not have a right inverse if $g \geq g(3)$ (we can take 86 for $g(3)$). In fact the induced homomorphism $H_{2i}(\text{Diff}_{+\Sigma_g}) \longrightarrow H_{2i}(M(g))$ is not

surjective for $i=3,4,\dots,k$ if $g \geq g(k)$ (here we consider $\text{Diff}_+ \Sigma_g$ as a discrete group).

This result should be compared with the recent affirmative solution of the Nielsen realization problem by Kerckhoff [Ke].

THEOREM 3. For any $k \in \mathbb{N}$, there exists a natural number $g'(k)$ such that the $2i$ -th Betti number $b_{2i}(M(g))$ of $M(g)$, which is equal to $b_{2i}(\mathcal{M}_g)$, is at least i for all $i=1,\dots,k$ if $g \geq g'(k)$.

We mention that Harer [Ha] has proved that $b_2(M(g)) = 1$ for $g \geq 5$. Since "half" of our characteristic classes come from $S_p(2g; \mathbb{Z})$, we also have informations on the homomorphism $H_{2i}(M(g)) \longrightarrow H_{2i}(S_p(2g; \mathbb{Z}))$. We omit the precise statement.

Sketch of Proofs. The proofs of the above results are given by constructing sufficiently many surface bundles with non-trivial characteristic classes. Roughly speaking we apply the method of Atiyah [A] (see also [Ko]) iteratively. To be more precise, let $\pi: E \longrightarrow X$ be a surface bundle with fibre Σ_g . We assume that X is an iterated surface bundle. Given $(n, n') \in \mathbb{N} \times \mathbb{N}$, an " (n, n') -construction on $\pi: E \longrightarrow X$ " is described by the following diagram of surface bundles:

$$\begin{array}{ccccccccc}
 F & \longrightarrow & E_2^* & \longrightarrow & E_1^* & \longrightarrow & E_1^* & \longrightarrow & E^* \\
 \downarrow \Sigma'' & & \downarrow \Sigma' & & \downarrow \Sigma' & & \downarrow \Sigma & & \downarrow \Sigma \\
 E_2 & \xlongequal{\quad} & E_2 & \longrightarrow & E_1 & \xlongequal{\quad} & E_1 & \longrightarrow & E \\
 & & & & & & & & \downarrow \Sigma \\
 & & & & & & & & X .
 \end{array}$$

Here $E^* \longrightarrow E$ is the pull back bundle $\pi^*(E)$. E^* contains a cross-section D as the diagonal. $E_1 \longrightarrow E$ is a covering map which kills

first the action of $\pi_1(E)$ on $H^1(\text{fibre}; \mathbb{Z}/n\mathbb{N}')$ and then kills $H^1(\ ; \mathbb{Z}/n\mathbb{N}')$. $E_1^* \rightarrow E^*$ is the pull back by this map. $E_1^* \rightarrow E_1^*$ is a fibre-wise $n\mathbb{N}'$ -fold covering map. $E_2 \rightarrow E_1$ is a covering map which satisfies the condition: the homology class of the inverse image D' of D under the map $E_2^* \rightarrow E^*$ is divisible by n . The assumption that X is an iterated surface bundle guarantees the existence of such covering. Finally $F \rightarrow E_2^*$ is an n -fold cyclic ramified covering ramified along D' . $F \rightarrow E_2$ is a surface bundle with fibre Σ'' whose genus is $n^2 n' g - \frac{1}{2} n(n+1)n'+1$. The $(2,1)$ -construction on the trivial surface bundle $\Sigma_g \rightarrow \text{pt.}$ is nothing but Atiyah's method in [A]. Theorem 1 is proved by calculating e_k of surface bundles which are defined by applying (n_j, n'_j) -constructions on $\Sigma_g \rightarrow \text{pt.}$ successively ($j=1, \dots, k$). It turns out that e_k of such a surface bundle is $(g-1)$ times a non-trivial polynomial of n_j, n'_j 's. Since such surface bundles admit multi-valued cross-sections, once the statement of Theorem 1 is proved for one g_0 , it holds for all $g \geq g_0$. Corollary 2 follows from Theorem 1 and the Bott vanishing theorem [B]. We can also compute other characteristic classes than e_k . It turns out that $e_{i_1}^{d_1} \dots e_{i_s}^{d_s}$ ($\sum_j i_j d_j = k$) is a linear combination of $(g-1), (g-1)^2, \dots, (g-1)^{d_1 + \dots + d_s}$ with coefficients in polynomials of n_j, n'_j 's. Theorem 3 follows from this.

It is very likely that these examples of surface bundles are enough to prove the injectivity of ϕ in small degrees. However necessary computations for that are extremely complicated. Also it seems to be interesting to test the surjectivity of ϕ by examining these examples because it is by no means clear that characteristic numbers of a surface bundle which is obtained by applying an (n, n') -construction on another

surface bundle depend only on those of the latter. The details together with these points will appear elsewhere.

REFERENCES

- [A] M.F. Atiyah, The signature of fibre-bundles, Global Analysis, Papers in Honor of K. Kodaira, Tokyo Univ. Press, 1969, 73-84.
- [BH] A. Borel and F. Hirzebruch, Characteristic classes and homogeneous spaces, I, Amer. Jour. Math. 80 (1958), 459-538.
- [B] R. Bott, On a topological obstruction to integrability, Proc. Symp. Pure Math. vol XVI, Global Analysis, AMS 1970, 127-131.
- [EE] C.J. Earle and J. Eells, The diffeomorphism group of a compact Riemann surface, Bull. Amer. Math. Soc. 73 (1967), 557-559.
- [Ha] J. Harer, The second homology group of the mapping class group of an orientable surface, Invent. Math. 72 (1983), 221-239.
- [Hv] W.J. Harvey, Geometric structures of surface mapping class groups, Homological Group Theory, LMS Lecture Notes No. 36, Cambridge Univ. Press, 1979, 255-269.
- [Ke] S.P. Kerckhoff, The Nielsen realization problem, Ann. of Math. 117 (1983), 235-265.
- [Ko] K. Kodaira, A certain type of irregular algebraic surface, Jour. Analyse Math. 19 (1967), 207-215.

Department of Mathematics, College of Arts and Sciences,
University of Tokyo, Tokyo, Japan

以上は 1983年7月現在のまとめである。その後の進展も含めて詳細な結果は現在論文を準備中。とくに定理3に關しては、準同型 ϕ は 1次の次数が injective であることが証明された。