

## グラフの星部分グラフ分解

Star decomposition indexes of graphs

加納 幹 雄 明石工高専

Mikio KANO, Akashi Technological College

We deal with finite simple graphs, which have neither multiple edges nor loops. Let  $G$  be a graph with vertex set  $V(G)$  and edge set  $E(G)$ . The arboricity  $a(G)$  of  $G$  is the minimum integer  $n$  for which  $E(G)$  can be decomposed into  $n$  forests. A formula for the arboricity of a graph was obtained by Nash-William [5],[6]. The formula is the following:

$$a(G) = \max \left[ \frac{|E(H)|}{|V(H)| - 1} \right]$$

where the maximum is taken over all subgraphs  $H$  of  $G$ , and  $[x]$  denotes the least integer not less than  $x$ . If we impose some conditions to forests, then we obtain new invariants. A graph is called a linear forest if each component of it is a path, and linear arboricity  $\equiv(G)$  of  $G$  is defined to be the minimum  $n$  for which  $E(G)$  can be decomposed into  $n$  linear forests. Some results on linear arboricity can be found in [1],[4]. We call a graph  $H$  a star if  $H$  is isomorphic to the complete bipartite graph  $K_{1,n}$  for some  $n$  (Fig. 1). We call a graph  $G$




Figure 1.  $K_{1,4}$ .




Figure 2. A star-forest.




Figure 3. A graph  $G$  with  $*(G)=3$ . Figure 4.  $A$  and  $B$ .

a star-forest if each component of  $G$  is a star (Fig. 2). We define the star decomposition index  $*(G)$  of  $G$  by the minimum  $n$  for which  $E(G)$  can be decomposed into  $n$  star-forests (Fig. 3). In this paper we shall investigate star decomposition indexes.

We begin with the following easy result.

Proposition 1. Let  $T$  be a tree. If  $T$  is not a star, then  $*(T)=2$ .

Proof Let  $T$  be a tree that is not a star. Then it is obvious that  $*(T) \geq 2$ . Let  $L(T)$  be the line graph of  $T$  (i.e.  $V(L(T))=E(T)$  and two vertices of  $L(T)$  are adjacent if and only if corresponding edges of  $T$  are adjacent.). For two vertices  $x$  and  $y$  of  $L(T)$ , we denote by  $d(x,y)$  the distance between  $x$  and  $y$  in  $L(T)$ . Choose any vertex  $v$  of  $L(T)$ , and set

$$A = \{x \in V(L(T)) \mid d(v,x) \text{ is odd}\} \quad \text{and}$$

$$B = \{x \in V(L(T)) \mid d(v,x) \text{ is even}\} \ni v \quad (\text{Fig. 4}).$$

Then  $A$  and  $B$  are star-forests of  $T$ , and thus  $*(T) \leq 2$ . Therefore  $*(T)=2$ .  $\square$

By  $K_n$  and  $K_{n,m}$ , we denote the complete graph of order  $n$  and the complete graph of order  $n+m$ , respectively. Let  $A$  be a graph. Then an  $A$ -factor of a graph is its spanning subgraph each component of which is isomorphic to  $A$ .

Theorem 1. [2]  $*(K_{2n-1})=*(K_{2n})=n+1$ , where  $n \geq 3$ .

Proof We first show that  $*(K_{2n}) \geq *(K_{2n-1}) \geq n+1$ . It is obvious that  $*(K_{2n-1}) \leq *(K_{2n})$ . Since  $K_{2n-1}$  is a  $2(n-1)$ -regular graph and does not have a  $K_{1,n-1}$ -factor, we obtain  $*(K_{2n-1}) \geq n+1$  by Theorem 3, which will be given later.

We next show that  $*(K_{2n}) \leq n+1$ . Let  $V(K_{2n}) = \{v_1, v_2, \dots, v_{2n}\}$  and put

$$F_t = \{v_t v_i \mid t < i' < t+n, i \equiv i' \pmod{2n}\} \\ \cup \{v_{n+t} v_j \mid n+t < j' < t+2n, j \equiv j' \pmod{2n}\} \subset E(K_{2n})$$

for  $t=1, \dots, n$ , and define

$$F_{n+1} = \{v_1 v_{n+1}, v_2 v_{n+2}, \dots, v_n v_{2n}\} \quad (\text{Fig. 5}).$$

Then  $K_{2n} = F_1 \cup F_2 \cup \dots \cup F_{n+1}$ , and we conclude that  $*(K_{2n}) \leq n+1$ .

Consequently,  $*(K_{2n-1})=*(K_{2n})=n+1$ .  $\square$




Figure 5.  $F_1, F_2, F_3, F_4$  and  $F_5$  of  $K_8$ .

The star decomposition index of the complete bipartite graph  $K_{n,n}$  was determined by Egawa, Fukuda, Nagoya and Urabe [3], and  $*(K_{n,m})$  for some classes of  $n, m$  are obtained by Enomoto and etc.

Theorem 2. [3]  $*(K_{2n,2n})=*(K_{2n-1,2n-1})=n+2$ , where  $n \geq 4$ .

Proof We prove only that  $*(K_{2n-1,2n-1}) \leq *(K_{2n,2n}) \leq n+2$ . For the proof of  $*(K_{2n-1,2n-1}) \geq n+2$ , the reader should refer to [3]. It is trivial that  $*(K_{2n-1,2n-1}) \leq *(K_{2n,2n})$ . Let  $V(K_{2n,2n}) = \{a_1, \dots, a_n, b_1, \dots, b_n\} \cup \{c_1, \dots, c_n, d_1, \dots, d_n\}$ . For every  $k$ ,  $1 \leq k \leq n$ , we define

$$F_k = \{a_k c_i, b_k d_i, a_i d_k, b_i c_k \mid 1 \leq i \leq n, i \neq k\},$$

and put

$$F_{n+1} = \{a_i c_i, a_i d_i \mid 1 \leq i \leq n\}, \text{ and}$$

$$F_{n+2} = \{b_i c_i, b_i d_i \mid 1 \leq i \leq n\} \quad (\text{Fig. 6}).$$

Then  $K_{2n, 2n} = F_1 \cup F_2 \cup \dots \cup F_{n+2}$ . Consequently,  $\ast(K_{2n, 2n}) \leq n+2$ .




Figure 6.  $F_2$  and  $F_5$  of  $K_{8,8}$ .

We write  $d_G(v)$  for the degree of vertex  $v$  in  $G$ . A graph  $G$  is called an  $r$ -regular graph if  $d_G(x) = r$  for all vertices  $x$ .

Theorem 3. Let  $G$  be a  $2r$ -regular graph. Then

$$\ast(G) \geq r+1$$

with the equality if and only if  $G$  can be decomposed into  $r+1$  edge-disjoint  $K_{1,r}$ -factors.

Proof. Since  $a(G) \geq |E(G)| / (|V(G)| - 1) > r$ , we have  $\ast(G) \geq a(G) \geq r+1$ . Suppose  $\ast(G) = r+1$ . Then  $G$  can be decomposed into  $r+1$  star-forests  $H_1, H_2, \dots, H_{r+1}$ . We denote by  $n_i(t)$  the number of components  $K_{1,t}$  in  $H_i$ . Put  $p = |V(G)|$  and  $x_i = |V(G)| - |V(H_i)|$ . Then we have

$$p = x_k + \sum_{j=1}^{2r} (j+1)n_k(j) \quad \text{for all } k, 1 \leq k \leq r+1 \quad (1)$$

$$|E(H_k)| = \sum_{j \geq 1} j n_k(j) = p - (x_k + \sum_{j \geq 1} n_k(j)), \text{ and}$$

$$\begin{aligned} \sum_{x \in V(G)} d_G(x) &= 2pr = 2|E(G)| = 2 \sum_{k=1}^{r+1} |E(H_k)| \\ &= 2p(r+1) - 2 \sum_{k=1}^{r+1} \{x_k + \sum_{j \geq 1} n_k(j)\} \end{aligned}$$

Therefore

$$p = \sum_{k=1}^{r+1} \{x_k + \sum_{j \geq 1} n_k(j)\}. \quad (2)$$

The vertex of  $K_{1,t}$  ( $t \geq 2$ ) with degree  $t$  is called the center of  $K_{1,t}$ . It follows that every vertex  $v$  of  $K_{1,j}$  ( $j < r$ ) in  $H_1$  must be the center of a component  $K_{1,t}$  ( $t \geq 2$ ) in some  $H_k$  ( $k \geq 2$ ), since otherwise  $2r = d_G(v) = d_{H_1}(v) + \sum_{k \geq 2} d_{H_k}(v) < r+r$ , a contradiction.

Similarly, every end vertex of  $K_{1,j}$  ( $j \geq r$ ) in  $H_1$  is contained in the center of a component in some  $H_k$  ( $k \geq 2$ ). Hence

$$x_1 + \sum_{j=1}^{r-1} (j+1)n_1(j) + \sum_{j=r}^{2r} jn_1(j) \leq \sum_{k=2}^{r+1} \left( \sum_{t \geq 2} n_k(t) \right) \quad (3)$$

By substituting (3) into (2), we obtain

$$\begin{aligned} p &= x_1 + \sum_{j \geq 1} n_1(j) + \sum_{k=2}^{r+1} x_k + \sum_{k=2}^{r+1} \left( \sum_{j \geq 2} n_k(j) \right) + \sum_{k=2}^{r+1} n_k(1) \\ &\geq x_1 + \sum_{j \geq 1} n_1(j) + \sum_{k=2}^{r+1} x_k + x_1 + \sum_{j=1}^{r-1} (j+1)n_1(j) + \sum_{j=r}^{2r} jn_1(j) \\ &\quad + \sum_{k=2}^{r+1} n_k(1) \\ &= x_1 + \sum_{j \geq 1} (j+1)n_1(j) + \sum_{j=1}^{r-1} n_1(j) + \sum_{k=2}^{r+1} x_k + x_1 + \sum_{k=2}^{r+1} n_k(1) \\ &= p + \sum_{j=1}^{r-1} n_1(j) + \sum_{k=2}^{r+1} x_k + x_1 + \sum_{k=2}^{r+1} n_k(1). \quad (\text{by (1)}) \end{aligned}$$

Hence  $n_1(j) = 0$  for every  $j$ ,  $1 \leq j \leq r-1$ , and  $x_1 = \dots = x_{r+1} = 0$ .

We can similarly show that  $n_k(j) = 0$  for all  $k$ ,  $j$  ( $k \geq 2$  and  $j \leq r-1$ ).

Therefore, each component of  $H_k$  is  $K_{1,t}$  ( $t \geq r$ ), and  $H_k$  is a spanning subgraph of  $G$ . If  $d_{H_k}(v) \geq r+1$  for some  $k \geq 1$  and  $v \in V(G)$ , then  $2r = d_G(v) = \sum_{t \geq 1} d_{H_t}(v) \geq r+1+r = 2r+1$ , a contradiction.

Consequently, each  $H_k$  has no  $K_{1,t}$  for  $t \geq r+1$ , and we conclude that every  $H_k$  is a  $K_{1,r}$ -factor of  $G$ . Hence the proof is complete.  $\square$

The next theorem can be proved by the same argument in the proof of Theorem 3.

Theorem 4. Let  $G$  be a  $(2r+1)$ -regular graph. Then  $*(G) \geq r+2$ .

By Theorems 3 and 4, we have

$r$	2	3	4	5	6
$*(G)$ of $r$ -regular graph $G$	2	3	3,4	4,5	4,5,6

Note that the existence of a 5-regular graph  $G_1$  with  $*(G_1)=5$  and of a 6-regular graph  $G_2$  with  $*(G_2)=6$  is unknown.

A triangle cluster is a connected graph whose edges partition into disjoint triangles with the property that any two triangles have at most one vertex in common and if such a vertex exists, then it is a cut vertex of the cluster (Fig. 7).




Figure 7. A triangle cluster.

Proposition 2. (Fukuda [5]) Let  $TC$  be a triangle cluster. Then

$$*(TC) = \begin{cases} 2 & \text{if every triangle has a vertex of degree 2} \\ 3 & \text{otherwise.} \end{cases}$$

## References

- [1] J. Akiyama, Factorization and linear arboricity of graphs, Doctor thesis, Science University of Tokyo, (1982).
- [2] J. Akiyama and M. Kano, Path factors of graphs, Graphs and Applications (Proc. of the first Colorado symposium of graph theory, Wiley) to appear.
- [3] Y. Egawa, T. Fukuda, S. Nagoya and M. Urabe, A decomposition of complete bipartite graphs into edge-disjoint subgraphs with star components, to appear.
- [4] H. Enomoto, The linear arboricity of 5-regular graphs, to appear.
- [5] T. Fukuda, 修士論文, 電通大情報数理工学科 (1984),
- [6] C. Nash-William, Decomposition of finite graphs into forests J. London Math. Soc. 39 (1964) 12.
- [7] C. Nash-William, Edge-disjoint spanning trees of finite graphs, J. London Math. Soc. 36 (1961) 445-450.