

The automorphism group of the Klein curve in the mapping class group of genus 3

Atsushi Matsuura
松浦篤司 (東大数理) *

1 The main result and its proof

Let R be a compact Riemann surface of genus $g \geq 2$. Then $\text{Aut}(R)$, the automorphism group of R , can be embedded into the mapping class group (for its definition, see [Bir, Ch. 4]) or the Teichmüller group Γ_g of genus g ;

$$(1.1) \quad \iota: \text{Aut}(R) \hookrightarrow \Gamma_g \simeq \text{Out}^+(\pi_1(R)) = \text{Aut}^+(\pi_1(R))/\text{Int}(\pi_1(R)).$$

Here, $\text{Aut}^+(\pi_1(R))$ consists of the automorphisms of $\pi_1(R)$ inducing the trivial action on $H_2(\pi_1(R), \mathbb{Z}) \simeq \mathbb{Z}$.

Recall the Hurwitz theorem, which states that

$$(1.2) \quad \#\text{Aut}(R) \leq 84(g-1).$$

If the equality holds in (1.2), then R is called a Hurwitz Riemann surface and $\text{Aut}(R)$ is called a Hurwitz group.

Let X be the Klein curve of genus 3 defined by the equation

$$x^3y + y^3z + z^3x = 0.$$

It is well known that X is a Hurwitz Riemann surface; $G := \text{Aut}(X)$ is isomorphic to $PSL_2(\mathbb{F}_7)$ and has order 168.

Now let us forget about the Klein curve, and consider an orientable compact C^∞ surface X of genus 3. We define the canonical generators of $\pi_1(X, b)$ with base point b as in the figure below;

*Department of Mathematical Sciences, University of Tokyo, Japan
E-mail address : matsuura@ms.u-tokyo.ac.jp




Figure 1

They satisfy the fundamental relation

$$(1.3) \quad (a_1 b_1 a_1^{-1} b_1^{-1})(a_2 b_2 a_2^{-1} b_2^{-1})(a_3 b_3 a_3^{-1} b_3^{-1}) = 1.$$

Let $\tilde{\varphi}_2, \tilde{\varphi}_3, \tilde{\varphi}_7$ be the elements of $\text{Aut}^+(\pi_1(X))$ defined by

$$\begin{aligned} \tilde{\varphi}_2(a_1) &= a_2 b_2^{-1} a_2^{-1} a_1^{-1} b_3^{-1} b_2 \\ \tilde{\varphi}_2(a_2) &= b_3^{-1} a_2^{-1} \\ \tilde{\varphi}_2(a_3) &= a_2 b_2^{-1} a_2^{-1} b_1^{-1} a_1^{-1} a_3 a_2^{-1} \\ \tilde{\varphi}_2(b_1) &= b_2^{-1} b_3 b_1^{-1} a_2 b_2 a_2^{-1} \\ \tilde{\varphi}_2(b_2) &= a_2 b_3 b_2^{-1} a_2^{-1} \\ \tilde{\varphi}_2(b_3) &= a_2 b_3 a_2^{-1}, \end{aligned}$$

$$\begin{aligned} \tilde{\varphi}_3(a_1) &= a_2 b_3 a_3^{-1} a_1 a_2 b_2 a_2^{-1} \\ \tilde{\varphi}_3(a_2) &= a_3^{-1} a_1 b_1 a_1^{-1} \\ \tilde{\varphi}_3(a_3) &= a_2 b_2 a_2 b_2^{-1} a_2^{-1} b_1 \\ \tilde{\varphi}_3(b_1) &= a_2 b_2^{-1} a_2^{-1} a_1^{-1} a_3 a_1 a_2 b_2 a_2^{-1} \\ \tilde{\varphi}_3(b_2) &= a_1 b_1^{-1} a_1^{-1} a_3 a_2 b_2^{-1} a_2^{-1} b_1 a_1^{-1} \\ \tilde{\varphi}_3(b_3) &= a_1 b_1^{-1} a_1^{-1} a_3 a_2 b_2^{-1} a_2^{-1} b_1, \end{aligned}$$

$$\begin{aligned} \tilde{\varphi}_7(a_1) &= b_1^{-1} a_1^{-1} a_3 b_3^{-1} a_2^{-1} \\ \tilde{\varphi}_7(a_2) &= a_2 b_2^{-1} a_2^{-1} a_1^{-1} \\ \tilde{\varphi}_7(a_3) &= b_1^{-1} a_2 b_2 a_2^{-1} a_3^{-1} a_1 b_1 a_1^{-1} \\ \tilde{\varphi}_7(b_1) &= a_2 b_3 a_3^{-1} a_1 a_2 b_2 a_2^{-1} b_3^{-1} a_2^{-1} \\ \tilde{\varphi}_7(b_2) &= a_1 a_2 b_2 b_3 a_3^{-1} \\ \tilde{\varphi}_7(b_3) &= a_1 a_2 b_2 a_3^{-1} a_1 b_1 a_1^{-1}. \end{aligned}$$

Then, we have the following:

Theorem 1.1. (1) *The classes φ_i of $\tilde{\varphi}_i$ in $\text{Out}^+(\pi_1(X))$ generate a subgroup H of Γ_3 , which is isomorphic to $PSL_2(\mathbb{F}_7)$.*

(2) *Moreover, if X is the Klein curve, then H is conjugate to the image of ι .*

Proof. (1) First note that $H \neq \{1\}$, because the action of H on the homology group $H_1(X, \mathbb{Z})$ is not trivial. By direct computation using (1.3), we have

$$(1.4) \quad \begin{aligned} \tilde{\varphi}_2^2 = \tilde{\varphi}_3^3 = \tilde{\varphi}_7^7 = 1, \quad \tilde{\varphi}_2 \tilde{\varphi}_3 \tilde{\varphi}_7 = 1, \\ (\tilde{\varphi}_7 \tilde{\varphi}_3 \tilde{\varphi}_2)^4 = [\text{conjugation by } a_2 b_2^{-1} a_2^{-1} b_1]. \end{aligned}$$

For example,

$$\begin{aligned} \tilde{\varphi}_3^2 \cdot b_3 &= (a_2^{-1} b_2 a_2 a_1 a_3 a_1^{-1} a_2^{-1} b_2^{-1} a_2) (a_3 a_1^{-1} b_1^{-1} a_1) \\ &\quad \times (a_1^{-1} b_1 a_1 a_3^{-1} a_2^{-1} b_2 a_2 b_1^{-1} a_1) (a_1^{-1} b_1 a_1 a_3^{-1}) \\ &\quad \times (b_1 a_2^{-1} b_2^{-1} a_2 b_2 a_2) (a_2^{-1} b_3^{-1} a_3 a_1^{-1} a_2^{-1} b_2^{-1} a_2) \\ &\quad \times (a_2^{-1} b_2 a_2 a_1 a_3^{-1} a_1^{-1} a_2^{-1} b_2^{-1} a_2) (a_2^{-1} b_2 a_2 a_1 a_3^{-1} b_3 a_2) \\ &= a_2^{-1} b_2 a_2 (a_1 b_1 a_2^{-1} b_2^{-1} a_2 b_2 b_3^{-1} a_3^{-1} b_3) a_2 \\ &= a_2^{-1} b_2 a_2 b_1 a_1 a_3^{-1} a_2, \end{aligned}$$

hence

$$\begin{aligned} \tilde{\varphi}_3^3 \cdot b_3 &= (a_3 a_1^{-1} b_1^{-1} a_1) (a_1^{-1} b_1 a_2^{-1} b_2^{-1} a_2 a_3 a_1^{-1} b_1^{-1} a_1) \\ &\quad \times (a_1^{-1} b_1 a_1 a_3^{-1}) (a_2^{-1} b_2 a_2 a_1 a_3 a_1^{-1} a_2^{-1} b_2^{-1} a_2) \\ &\quad \times (a_2^{-1} b_2 a_2 a_1 a_3^{-1} b_3 a_2) (a_2^{-1} b_2^{-1} a_2^{-1} b_2 a_2 b_1^{-1}) (a_1^{-1} b_1 a_1 a_3^{-1}) \\ &= b_3. \end{aligned}$$

From (1.4) we obtain

$$(1.5) \quad \varphi_2^2 = \varphi_3^3 = \varphi_7^7 = \varphi_2 \varphi_3 \varphi_7 = (\varphi_7 \varphi_3 \varphi_2)^4 = 1$$

in $\text{Out}^+(\pi_1(X))$. Since (1.5) is the presentation of $PSL_2(\mathbb{F}_7)$ (see [CM, p. 96]), there is a surjective map

$$PSL_2(\mathbb{F}_7) \twoheadrightarrow H.$$

The group $PSL_2(\mathbb{F}_7)$ is simple, and the map is an isomorphism.

(2) To see that H is the automorphism group of a Riemann surface, it is enough to recall the Nielsen realization problem, which was positively solved in [Ker]:

Theorem of Kerckhoff . *For any finite subgroup G of Γ_g , there is a compact Riemann surface R of genus g such that*

$$G \subset \text{Aut}(R) \subset \Gamma_g.$$

□

This theorem shows that there exists a Riemann surface R of genus 3 with $H \subset \text{Aut}(R)$. On the other hand, $\#\text{Aut}(R) \leq 168 = \#H$ by the Hurwitz inequality. Consequently $H = \text{Aut}(R)$. It is classically known that the Klein curve is the unique compact Riemann surface of genus 3 such that $\text{Aut}(R) \simeq PSL_2(\mathbb{F}_7)$. Thus we have proved Theorem 1.1. □

2 $\pi_1(X)$ as a subgroup of the triangle group of type $(2, 3, 7)$

In this section, we give a more elementary proof of Theorem 1.1. The outline is as follows: Let T be the triangle group with angles $\frac{\pi}{2}, \frac{\pi}{3}, \frac{\pi}{7}$ defined below, and N its normal subgroup. Then, T (resp. N) has a fundamental domain Δ (resp. Λ) in the Poincaré unit disk. As was shown in [Kle], the Klein curve X can be realized by gluing the boundaries of Λ . The elements of T act on Λ , hence on X . This action induces an isomorphism $T/N \simeq \text{Aut}(X)$. Moreover, N is isomorphic to $\pi_1(X)$. Because T acts on N by conjugation, T/N can be embedded in $\text{Out}^+(\pi_1(X))$. In this way, we obtain the map ι in (1.1). First, we compute the elements of N corresponding to the generators of $\pi_1(X)$. Using this identification, we show that $\iota(T/N) = H$, which is equivalent to Theorem 1.1.

Let $S = \begin{pmatrix} 0 & -1 \\ 1 & 0 \end{pmatrix}$, $T = \begin{pmatrix} 1 & -1 \\ 0 & 1 \end{pmatrix}$ be the generators of $PSL_2(\mathbb{F}_7)$. Then $S^2 = T^7 = (ST^{-1})^3 = 1$. For the triangle group

$$T := \langle \gamma_2, \gamma_3, \gamma_7 \mid \gamma_2^2 = \gamma_3^3 = \gamma_7^7 = 1, \gamma_2\gamma_3\gamma_7 = 1 \rangle,$$

we define a group homomorphism

$$\varphi: T \rightarrow PSL_2(\mathbb{F}_7)$$

by $\varphi(\gamma_2) = S, \varphi(\gamma_3) = ST^{-1}, \varphi(\gamma_7) = T$. Clearly φ is surjective. The map φ gives an exact sequence

$$(2.1) \quad 1 \rightarrow N \rightarrow T \rightarrow PSL_2(\mathbb{F}_7) \rightarrow 1,$$

where $N := \ker \varphi \simeq \pi_1(X)$ is the kernel of φ . Hence we have $G \simeq T/N$. For any element $\hat{\alpha}$ of N , we shall denote by α the loop with base point b representing $\hat{\alpha}$. First, we give the elements of N corresponding to the canonical generators of $\pi_1(X, b)$. Note that, for two elements $\hat{\alpha}, \hat{\beta} \in N$, their product $\hat{\alpha}\hat{\beta} \in N$ corresponds to the loop $\beta\alpha$.

Proposition 2.1. *Define $\hat{a}_i, \hat{b}_i \in N, i = 1, 2, 3$ by*

$$\begin{aligned}\hat{a}_1 &= \gamma_7 \gamma_3^{-1} \gamma_7^{-3} \gamma_2 \gamma_7^2 (\gamma_3 \gamma_2 \gamma_7)^4 \gamma_7^{-2} \gamma_2 \gamma_7^3 \gamma_3 \gamma_7^{-1} \\ \hat{b}_1 &= \gamma_7 \gamma_3^{-1} \gamma_7^{-3} \gamma_2 \gamma_7^2 (\gamma_2 \gamma_7 \gamma_3)^4 \gamma_7^{-2} \gamma_2 \gamma_7^3 \gamma_3 \gamma_7^{-1} \\ \hat{a}_2 &= \gamma_2 \gamma_7^{-4} \gamma_2 \gamma_7^{-4} \gamma_2 \gamma_7^2 (\gamma_3 \gamma_2 \gamma_7)^4 \gamma_7^{-2} \gamma_2 \gamma_7^4 \gamma_2 \gamma_7^4 \gamma_2 \\ \hat{b}_2 &= \gamma_2 \gamma_7^{-4} \gamma_2 \gamma_7^{-4} \gamma_2 \gamma_7^2 (\gamma_2 \gamma_7 \gamma_3)^4 \gamma_7^{-2} \gamma_2 \gamma_7^4 \gamma_2 \gamma_7^4 \gamma_2 \\ \hat{a}_3 &= \gamma_3 \gamma_7^{-2} \gamma_2 \gamma_7^{-4} \gamma_3 (\gamma_7^{-1} \gamma_2 \gamma_3^{-1})^4 \gamma_3^{-1} \gamma_7^4 \gamma_2 \gamma_7^2 \gamma_3^{-1} \\ \hat{b}_3 &= \gamma_3 \gamma_7^{-2} \gamma_2 \gamma_7^{-4} \gamma_3 (\gamma_2 \gamma_7 \gamma_3)^4 \gamma_3^{-1} \gamma_7^4 \gamma_2 \gamma_7^2 \gamma_3^{-1}.\end{aligned}$$

Set $\hat{a}'_3 = \hat{a}_3^{-1} \hat{b}_3$, $\hat{b}'_3 = \hat{a}_3$. Then the elements $\hat{a}_1, \hat{a}_2, \hat{a}'_3, \hat{b}_1, \hat{b}_2, \hat{b}'_3$ are identified with the canonical generators of $\pi_1(X)$ and they satisfy the equation $[\hat{a}'_3, \hat{b}'_3] [\hat{a}_2, \hat{b}_2] [\hat{a}_1, \hat{b}_1] = 1$. Here $[\alpha, \beta] := \beta^{-1} \alpha^{-1} \beta \alpha$.

Proof. Let Δ (resp. Λ) be the fundamental domain of T (resp. N). Figure 2 below illustrates that Δ is a hyperbolic triangle with angles $\frac{\pi}{3}, \frac{\pi}{3}, \frac{2\pi}{7}$ and Λ is the union of 168 copies of Δ . By tracing paths, we can easily see that the elements $\hat{a}_i, \hat{b}_i$ in the figure can be written as above.

By gluing corresponding edges, we obtain the Riemann surface X . The elements $\hat{a}_i, \hat{b}_i$ are represented by the loops a_i, b_i in Figure 1. We can also check the fundamental relation by computation. $\square$

The conjugation gives the canonical map

$$\tilde{\iota}: T \rightarrow \text{Aut}^+(N).$$


This induces the map ι in (1.1). We take $\tilde{\iota}(\gamma_2), \tilde{\iota}(\gamma_3), \tilde{\iota}(\gamma_7)$ as the generators of $\iota(T/N)$.

The following proposition finishes the direct proof of Theorem 1.1.


Proposition 2.2. *$\tilde{\iota}(\gamma_j)$ can be identified with $\tilde{\varphi}_j$ for $j = 1, 2, 3$.*

Proof. Set $\gamma_j \cdot \alpha := \tilde{i}(\gamma_j)(\alpha) = \gamma_j \alpha \gamma_j^{-1}$ for $\alpha \in N$. Then, for $\hat{a}_i, \hat{b}_i$ in Proposition 2.1, we can describe $\gamma_j \cdot \hat{a}_i, \gamma_j \cdot \hat{b}_i \in \Lambda$ as in the Figure 3.

We shall show that $\tilde{\varphi}_7(a_1)$ represents $\tilde{i}(\gamma_7)(\hat{a}_1)$. By gluing the edges of Λ , we get the following loop ℓ representing $\gamma_7 \cdot \hat{a}_1$.



We can check that ℓ is homotopic to the loop below, which is the loop $b_1^{-1} a_1^{-1} a_3 b_3^{-1} a_2^{-1}$.



The proofs for the other cases are similar and omitted. □

Acknowledgment. I would like to thank Professor Takayuki Oda for suggesting the problem, and for valuable discussions and encouragements. I also

wish to thank Takuya Kon-no for careful reading of the manuscript.

References

- [Bir] J.S. Birman, *Braids, links, and mapping class groups*, Princeton Univ. Press, 1974.
- [CM] H.S.M. Coxeter and W.O.J. Moser, *Generators and relations for discrete groups*, Springer-Verlag, 1972.
- [Ker] S.P. Kerckhoff, *The Nielsen realization problem*, Ann. of Math. **117** (1983), pp. 235–265.
- [Kle] F. Klein, *Über die Transformation siebenter Ordnung der elliptischen Functionen*, Gesammelte Math. Abhandlungen, **Band III**, pp. 90–136, Springer-Verlag, 1923.
- [Mac] A.M. Macbeath, *Generators of the Linear Fractional Groups*, Proc. Symp. Pure Math., vol. 12, A.M.S., 1968, pp. 14–32.
- [Mag] W. Magnus, *Noneuclidean tessellations and their groups*, Academic Press, 1970.







Figure 2: Fundamental domain Λ of N ([Kle, p. 126])

Glue $1=6$, $7=12$, $2=11$, $3=8$, $5=10$, $4=13$, $9=14$ in this order. Each loop is connected to the base point b by dotted path.

Figure 3-(i): Action of γ_2

Figure 3-(ii): Action of γ_3

Figure 3-(iii): Action of γ_7