

A Proof of the Krohn-Rhodes Decomposition Theorem

Z. Ésik*

A. József University
Department of Computer Science
Szeged, Hungary
esik@inf.u-szeged.hu

Abstract

We give a new proof of one part of the Krohn-Rhodes decomposition theorem for automata.

1 Introduction

The Krohn-Rhodes Decomposition Theorem [8] has a number of formulations in terms of automata, transformation semigroups, or semigroups, see [1, 6, 2, 9, 7, 5, 10], or [3], for an extension. The aim of this paper is to give a simple proof of the hard part of the theorem involving automata: Each finite automaton $\mathbf{A}$ is the homomorphic image of a subautomaton of a (generalized) cascade composition of automata $\mathbf{A}_1, \dots, \mathbf{A}_k$, where each $\mathbf{A}_i$ is either the two-state identity-reset automaton $\mathbf{U}$ or a group-type automaton $\mathbf{Aut}(G)$ corresponding to a simple group G which divides the semigroup of $\mathbf{A}$. In addition to the well-known decomposition of permutation-reset automata, the new argument uses a single construction and is based on the following observation. Given the automaton $\mathbf{A}$, there is a sequence

$$\mathbf{B}_0, \mathbf{B}_1, \dots, \mathbf{B}_m$$

of finite automata such that $\mathbf{B}_0$ is trivial, $\mathbf{B}_m$ is the automaton $\mathbf{A}$, and for each integer $1 \leq i \leq m$, either there is a surjective simple regular $\mathcal{G}$ -homomorphism $\mathbf{A}_i \rightarrow \mathbf{A}_{i-1}$, or there is a surjective simple regular $\mathcal{G}$ -homomorphism $\mathbf{A}_{i-1} \rightarrow \mathbf{A}_i$. Here $\mathcal{G}$ denotes the class of simple groups dividing the semigroup of $\mathbf{A}$, and a homomorphism $\mathbf{B} \rightarrow \mathbf{C}$ is termed a simple regular $\mathcal{G}$ -homomorphism if its kernel ρ satisfies the following conditions.

- The non-singleton equivalence classes of ρ , or ρ -blocks, for short, have equal cardinality.
- If C and D are (non-singleton) ρ -blocks and u is an input word of $\mathbf{B}$ with $Cu \subseteq D$, then either $Cu = D$ or Cu is a singleton set.
- For any two non-singleton ρ -blocks C and D there is a word u with $Cu = D$.
- If C is a ρ -block and G is the group of all bijections $C \rightarrow C$ induced by an input word, then any simple group divisor of G belongs to $\mathcal{G}$.

*Partially supported by the National Foundation for Scientific Research of Hungary under grant No. T7383 and by the Japan Society for the Promotion of Science.

We then show that if $h : \mathbf{B} \rightarrow \mathbf{C}$ is a surjective simple regular $\mathcal{G}$ -homomorphism with kernel ρ , then $\mathbf{B}$ is isomorphic to a subautomaton of a cascade composition of $\mathbf{C}$ and a permutation-reset automaton $\mathbf{D}$ such that each simple group divisor of the semigroup of $\mathbf{D}$ is in $\mathcal{G}$.

The proof presented here has been used in [4] to show that the Conway axioms and an identity associated with each finite (simple) group provide a complete axiomatization of iteration theories.

2 Preliminaries

2.1 Automata

Suppose that X is a finite nonempty set. We denote by X^* the free monoid of all words over X including the empty word λ . We set $X^+ = X^* - \{\lambda\}$, so that X^+ is the free semigroup of nonempty words over X .

An X -*automaton* $\mathbf{A}$ is a system (A, X, δ) consisting of the finite nonempty set A of states, the finite nonempty set X of input letters, and the transition function $\delta : A \times X \rightarrow A$ which can be extended to a function $A \times X^* \rightarrow A$ in the usual way. When $a \in A$ and $u \in X^*$, we will usually write au for $\delta(a, u)$, in particular when $\mathbf{A}$ is understood. Suppose that $C \subseteq A$ and $u \in X^*$. We define $Cu = \{cu : c \in C\}$.

Homomorphisms, congruences and subautomata are defined in the usual way.

2.2 Cascade Composition

Suppose that $\mathbf{A}_i = (A_i, X, \delta_i)$ are given automata, for $i \in [k] = \{1, \dots, k\}$, $k \geq 0$. Let X denote a finite nonempty set, and for each $i \in [k]$, let φ_i be a function

$$A_1 \times A_2 \times \dots \times A_{i-1} \times X \rightarrow X_i.$$

The *generalized cascade composition* of the $\mathbf{A}_i$ determined by the set X and functions φ_i is defined to be the automaton $\mathbf{A} = (A, X, \delta)$, where A is the set $A_1 \times \dots \times A_k$, and for each $(a_1, \dots, a_k) \in A$ and $x \in X$,

$$(a_1, \dots, a_k)x = (a_1x_1, \dots, a_kx_k)$$

with

$$x_i = \varphi_i(a_1, \dots, a_{i-1}, x),$$

all $i \in [k]$.

When $X = X_1 = \dots = X_k$ and $\varphi_i(a_1, \dots, a_{i-1}, x) = x$, for each $x \in X$, $a_1 \in A_1, \dots, a_{i-1} \in A_{i-1}$ and $i \in [k]$, the cascade composition becomes the *direct product* $\mathbf{A}_1 \times \dots \times \mathbf{A}_k$.

In the sequel, we will never use a generalized cascade composition of more than two automata at a time. Accordingly, we will write

$$\mathbf{A}_1 \times \mathbf{A}_2(X, \varphi_1, \varphi_2) \tag{1}$$

to denote the generalized cascade composition of $\mathbf{A}_1$ and $\mathbf{A}_2$ determined by the set X and functions φ_i , $i = 1, 2$. When X is the input set of the automaton $\mathbf{A}_1$ and φ_1 is the identity function $X \rightarrow X$, we call the automaton (1) the *cascade composition* of $\mathbf{A}_1$ and $\mathbf{A}_2$ determined by the function φ_2 . Denoting $\varphi = \varphi_2$, we will write

$$\mathbf{A}_1 \times_{\varphi} \mathbf{A}_2 \tag{2}$$

for short.

Suppose that $\mathbf{A} = (A, X, \delta)$ and $\mathbf{B} = (A, Y, \delta')$ are given finite automata with identical state sets. We say that $\mathbf{B}$ is a *renaming* of $\mathbf{A}$ if there is a function $\varphi : Y \rightarrow X$ such that

$$\delta'(a, y) = \delta(a, y\varphi),$$

for all $a \in A$ and $y \in Y$.

Suppose that K is a class of automata. We define:

- $\mathbf{S}(K)$: all subautomata of automata in K ;
- $\mathbf{N}(K)$: all renamings of automata in K ;
- $\mathbf{H}(K)$: all homomorphic images of automata in K ;
- $\mathbf{I}(K)$: all isomorphic images of automata in K ;
- $\mathbf{P}_c(K)$: all generalized cascade compositions of automata in K .

It is known that for any nonempty class K of automata, $\mathbf{V}_c(K) = \mathbf{HSP}_c(K)$ is the smallest class containing K and closed under the operators $\mathbf{H}$, $\mathbf{S}$ and $\mathbf{P}_c$, and also the smallest class containing K and closed under the operators $\mathbf{H}$, $\mathbf{S}$, $\mathbf{N}$ and the cascade composition (2). See [5].

2.3 Semigroups

Except for free semigroups X^+ and free monoids X^* , each semigroup will be assumed to be finite. We will use standard terminology. A submonoid of a semigroup is a subsemigroup which is a monoid. Similarly, a subgroup of a semigroup is a subsemigroup which is a group. Suppose that S and T are semigroups. We say that S *divides* T , denoted $S|T$, if S is a homomorphic image (or quotient) of a subsemigroup of T . It is known that this relation is transitive, see, e.g., [2, 9]. A proof of the following lemma can be found, e.g., in [5].

LEMMA 2.1 *Suppose that $S|T$ and that S is a monoid (group, respectively). Then there is a submonoid T' (subgroup, respectively) of T such that S is a quotient of T' .*

Suppose that $\mathbf{A} = (A, X, \delta)$ is an automaton. Each word $u \in X^*$ induces a function

$$\begin{aligned} u^{\mathbf{A}} : A &\rightarrow A \\ a &\mapsto au. \end{aligned}$$

The functions $u^{\mathbf{A}}$, $u \in X^*$, form a monoid denoted $M(\mathbf{A})$ whose unit is the identity function $\lambda^{\mathbf{A}} : A \rightarrow A$. We will denote by $S(\mathbf{A})$ the subsemigroup of $M(\mathbf{A})$ determined by the functions $u^{\mathbf{A}}$ induced by the nonempty words $u \in X^+$. The group $G(\mathbf{A})$ consists of those functions in $M(\mathbf{A})$ which are permutations.

We may generalize the above concepts. Suppose that C and D are two nonempty subsets of A . We define:

- $M_{\mathbf{A}}(C, D)$: all functions $f : C \rightarrow D$ such that there exists a word $u \in X^*$ with $u^{\mathbf{A}}|_C = f$, where $u^{\mathbf{A}}|_C$ denotes the restriction of $u^{\mathbf{A}}$ to C ;
- $S_{\mathbf{A}}(C, D)$: all functions $f : C \rightarrow D$ such that there exists a word $u \in X^+$ with $u^{\mathbf{A}}|_C = f$;

- $G_{\mathbf{A}}(C, D)$: the bijections in $M_{\mathbf{A}}(C, D)$.

Of course, if $G_{\mathbf{A}}(C, D) \neq \emptyset$, then $|C| = |D|$, i.e., the sets C and D have equal number of elements. We write $M_{\mathbf{A}}(C)$ for $M_{\mathbf{A}}(C, C)$. Note that $M_{\mathbf{A}}(C)$ is a monoid. We define the semigroup $S_{\mathbf{A}}(C)$ and the group $G_{\mathbf{A}}(C)$ in a similar way. Note that $S_{\mathbf{A}}(C)$ may be empty. For a proof of the following lemma, see [5].

LEMMA 2.2 *Suppose that G is a subgroup of $M_{\mathbf{A}}(C)$ or a subgroup of $S_{\mathbf{A}}(C)$. Then there is a nonempty set $D \subseteq C$ such that G is isomorphic to a subgroup of $G_{\mathbf{A}}(D)$. In particular, if G is a subgroup of $M(\mathbf{A})$ or a subgroup $S(\mathbf{A})$, then there is a set $D \subseteq A$ such that G is isomorphic to a subgroup of $G_{\mathbf{A}}(D)$.*

2.4 Permutation-Reset Automata

An X -automaton is a *permutation automaton* if each function $x^{\mathbf{A}}$, $x \in X$, is a permutation. It then follows that the functions $u^{\mathbf{A}}$, $u \in X^*$, are also permutations, so that $M(\mathbf{A}) = S(\mathbf{A}) = G(\mathbf{A})$. Conversely, if $S(\mathbf{A}) = G(\mathbf{A})$, or if $M(\mathbf{A}) = G(\mathbf{A})$, then $\mathbf{A}$ is a permutation automaton. When G is a group, the system $\mathbf{Aut}(G) = (G, G, \delta)$ with $\delta(g, h) = gh$, the product of the group elements g and h , for all $g, h \in G$, is a permutation automaton.

An automaton $\mathbf{A} = (A, X, \delta)$ is a *permutation-reset automaton* if each function $x^{\mathbf{A}}$, $x \in X$, is either a permutation or a constant map. It then follows that each function $u^{\mathbf{A}}$ for $u \in X^*$ is also either a permutation or a constant map. For example, the automaton $\mathbf{U} = ([2], \{x_0, x_1, x_2\}, \delta)$ is a permutation-reset automaton, where $ix_0 = i$ and $ix_j = j$, for $i, j = 1, 2$.

For any automaton $\mathbf{A}$, let $\mathcal{G}(\mathbf{A})$ denote the collection of simple groups G with $G|M(\mathbf{A})$. (Note that for any group G , $G|M(\mathbf{A})$ iff $G|S(\mathbf{A})$.) Moreover, we define $\mathcal{K}_g(\mathbf{A}) = \{\mathbf{Aut}(G) : G \in \mathcal{G}(\mathbf{A})\}$ and $\mathcal{K}(\mathbf{A}) = \mathcal{K}_g(\mathbf{A}) \cup \{\mathbf{U}\}$.

LEMMA 2.3 *Suppose that $\mathbf{A}$ is a permutation-reset automaton. Then*

$$\mathbf{A} \in \mathbf{V}_c(\mathcal{K}(\mathbf{A})).$$

If $\mathbf{A}$ is a permutation automaton such that at least one letter induces a nontrivial permutation, then

$$\mathbf{A} \in \mathbf{V}_c(\mathcal{K}_g(\mathbf{A})).$$

For a proof of Lemma 2.3, see [5], or [9].

3 The Krohn-Rhodes Decomposition Theorem

The Krohn-Rhodes Decomposition Theorem consists of two parts, Theorem 3.1 and Theorem 3.2. Let U denote a semigroup isomorphic to $M(\mathbf{U}) = S(\mathbf{U})$. (The automaton $\mathbf{U}$ was defined above).

THEOREM 3.1 *Suppose that S is either a semigroup dividing U or a simple group. Let $\mathbf{A}$ be an automaton and K a nonempty class of automata with $\mathbf{A} \in \mathbf{V}_c(K)$. If $S|S(\mathbf{A})$ then there is an automaton $\mathbf{B} \in K$ with $S|S(\mathbf{B})$. If $S|M(\mathbf{A})$ then there is an automaton $\mathbf{B} \in K$ with $S|M(\mathbf{B})$.*

THEOREM 3.2 *For each automaton $\mathbf{A}$,*

$$\mathbf{A} \in \mathbf{V}_c(\mathcal{K}(\mathbf{A})).$$

The class $\mathcal{K}(\mathbf{A})$ was defined above.

The rest of the paper is devoted to proving Theorem 3.2. In our argument, we will make use of Lemma 2.3, which is a particular instance of Theorem 3.2.

4 Congruences

In this section we assume that $\mathcal{G}$ is a class of simple groups closed under division. Thus, if G and H are simple groups with $G|H$ and $H \in \mathcal{G}$, then G is also in $\mathcal{G}$. The class $\overline{\mathcal{G}}$ consists of the groups whose simple group divisors are in $\mathcal{G}$. Note that $\overline{\mathcal{G}}$ is closed under the formation of subgroups and homomorphic images. It follows from Theorem 3.1 that $\overline{\mathcal{G}}$ is also closed under semidirect product and thus under direct product.

DEFINITION 4.1 Suppose that $\mathbf{A} = (A, X, \delta)$ is an automaton and that $\rho \subseteq A \times A$ is a congruence relation. We call ρ

- **simple**, if $|C| = |D|$ holds for any two non-singleton ρ -blocks $C, D \in A/\rho$, and if each member of $M_{\mathbf{A}}(C, D)$ is either a bijection or a constant map;
- **regular**, if for each non-singleton ρ -block C , the smallest congruence relation which collapses the states in C is the relation ρ itself;
- **A $\mathcal{G}$ -congruence**, if for each ρ -block C , each subgroup of $M_{\mathbf{A}}(C)$ is in $\overline{\mathcal{G}}$.

Note that ρ is a $\mathcal{G}$ -congruence iff for each ρ -block C , each subgroup of $S_{\mathbf{A}}(C)$ is in $\overline{\mathcal{G}}$, i.e., when $G \in \mathcal{G}$ holds for the simple groups G dividing $S_{\mathbf{A}}(C)$ or $M_{\mathbf{A}}(C)$. Moreover, a simple congruence ρ is a $\mathcal{G}$ -congruence iff $G_{\mathbf{A}}(C) \in \overline{\mathcal{G}}$, for each (non-singleton) ρ -block C . This follows by noting that when ρ is simple, each nontrivial subgroup of $M_{\mathbf{A}}(C)$ is a subgroup of $G_{\mathbf{A}}(C)$.

DEFINITION 4.2 Suppose that $\mathbf{A}$ and $\mathbf{B}$ are X -automata and that h is a homomorphism $\mathbf{A} \rightarrow \mathbf{B}$. We call h a simple, regular, or a $\mathcal{G}$ -homomorphism, if $\ker h$, the kernel of h has the appropriate property.

When $\mathcal{G}$ is empty, a $\mathcal{G}$ -homomorphism will be termed *aperiodic*.

LEMMA 4.3 Suppose that $\mathbf{A}_1, \mathbf{A}_2$ and $\mathbf{A}_3$ are X -automata with homomorphisms $h_1 : \mathbf{A}_1 \rightarrow \mathbf{A}_2$ and $\mathbf{A}_2 \rightarrow \mathbf{A}_3$. If h_1 is surjective and if

$$h = \mathbf{A}_1 \xrightarrow{h_1} \mathbf{A}_2 \xrightarrow{h_2} \mathbf{A}_3$$

is a $\mathcal{G}$ -homomorphism, then so are h_1 and h_2 .

Proof. Denote $\rho_i = \ker h_i$, $i = 1, 2$, and $\rho = \ker h$. Each ρ_1 -block C is included in some ρ -block D . The functions $g \in M_{\mathbf{A}_1}(D)$ with $Cg \subseteq C$ form a submonoid M of $M_{\mathbf{A}_1}(D)$, and the map $g \mapsto g|_C$, $g \in M$ is a surjective homomorphism $M \rightarrow M_{\mathbf{A}_1}(C)$. Thus $M_{\mathbf{A}_1}(C)|M_{\mathbf{A}_1}(D)$, so that any divisor of $M_{\mathbf{A}_1}(C)$ divides $M_{\mathbf{A}_1}(D)$. Since ρ is a $\mathcal{G}$ -congruence, it follows that ρ_1 is also a $\mathcal{G}$ -congruence, hence h_1 is a $\mathcal{G}$ -homomorphism.

Suppose now that C is a ρ_2 -block. Define $D = h_1^{-1}(C)$, so that D is a ρ -block. Since h_1 is surjective, the monoid $M_{\mathbf{A}_2}(C)$ is a quotient of $M_{\mathbf{A}_1}(D)$, a surjective homomorphism $M_{\mathbf{A}_1}(D) \rightarrow M_{\mathbf{A}_2}(C)$ is given by

$$u^{\mathbf{A}_1}|_D \mapsto u^{\mathbf{A}_2}|_C,$$

all $u \in X^*$ with $Du \subseteq D$. Thus any divisor of $M_{A_2}(C)$ divides $M_{A_1}(D)$. It follows that ρ_2 is a $\mathcal{G}$ -congruence and thus h_2 is a $\mathcal{G}$ -homomorphism. $\square$

COROLLARY 4.4 *Suppose that $\rho_1 \leq \rho_2$ are congruence relations of the automaton $\mathbf{A}$. If ρ_2 is a $\mathcal{G}$ -congruence, then so is ρ_1 . Further, ρ_2/ρ_1 is a $\mathcal{G}$ -congruence of the quotient automaton $\mathbf{A}/\rho_1$.*

REMARK 4.5 The assumption that h_1 is surjective was needed only in order to show that h_2 is a $\mathcal{G}$ -congruence.

In order to prove the converse of Lemma 4.3, we need the following fact.

LEMMA 4.6 *Suppose that $\mathbf{A} = (A, X, \delta)$ is a permutation X -automaton. Let ρ be a $\mathcal{G}$ -congruence relation of $\mathbf{A}$ such that $G(\mathbf{A}/\rho) \in \overline{\mathcal{G}}$. Then $G(\mathbf{A})$ is in $\overline{\mathcal{G}}$.*

Proof. Assume first that $\mathbf{A}$ is strongly connected, i.e., for each $a, b \in A$ there is some $u \in X^*$ with $au = b$. Let C_0 be a ρ -block. Define

$$Y = \{y_g : g \in G_{\mathbf{A}}(C_0)\}.$$

We turn C_0 into an Y -automaton $\mathbf{C}_0 = (C_0, Y, \delta_0)$ by defining

$$\delta_0(c, y_g) = cg,$$

for all $c \in C_0$ and $y_g \in Y$. It is known, see, e.g., [6, 2, 7], that $\mathbf{A}$ is isomorphic to a cascade composition of $\mathbf{A}/\rho$ and $\mathbf{C}_0$. See also Remark 6.3. Thus, by Theorem 3.1, each simple group divisor of $G(\mathbf{A})$ divides $G(\mathbf{A}/\rho)$ or $G(\mathbf{C}_0)$. (Note that $\mathbf{C}_0$ is a permutation automaton.) Since ρ is a $\mathcal{G}$ -congruence, $G(\mathbf{C}_0) = G_{\mathbf{A}}(C_0) \in \overline{\mathcal{G}}$. Further, $G(\mathbf{A}/\rho) \in \overline{\mathcal{G}}$, by assumption. It follows that $G(\mathbf{A}) \in \overline{\mathcal{G}}$.

When $\mathbf{A}$ is not strongly connected, then $\mathbf{A}$ is the disjoint sum of its strongly connected components $\mathbf{A}_1 = (A_1, X, \delta_1), \dots, \mathbf{A}_m = (A_m, X, \delta_m)$. Thus each $\mathbf{A}_i$ is a strongly connected permutation automaton, moreover, the sets A_i are pairwise disjoint, $\cup_{i=1}^m A_i = A$, and $\delta(a, x) = \delta_i(a, x)$ for each $a \in A_i$ and $x \in X$ with $i \in [m]$. The group $G(\mathbf{A})$ is isomorphic to a subgroup of the direct product of the groups $G(\mathbf{A}_i)$, in particular

$$G(\mathbf{A}) \mid \prod_{i=1}^m G(\mathbf{A}_i). \quad (3)$$

For each $i \in [m]$, let ρ_i denote the restriction of ρ to A_i . Then each ρ_i is a $\mathcal{G}$ -congruence relation of the strongly connected permutation automaton $\mathbf{A}_i$. But $G(\mathbf{A}_i/\rho_i)$ is a quotient of $G(\mathbf{A}/\rho)$, which is in $\overline{\mathcal{G}}$, by assumption. Thus each group $G(\mathbf{A}_i/\rho_i)$ is in $\overline{\mathcal{G}}$, so that $G(\mathbf{A}_i) \in \overline{\mathcal{G}}$, by the first part of the proof. Since $\overline{\mathcal{G}}$ is closed under direct product, it follows by (3) that $G(\mathbf{A})$ is also in $\overline{\mathcal{G}}$. $\square$

LEMMA 4.7 *Suppose that $\mathbf{A}_1, \mathbf{A}_2$ and $\mathbf{A}_3$ are X -automata and $h_1 : \mathbf{A}_1 \rightarrow \mathbf{A}_2$ and $h_2 : \mathbf{A}_2 \rightarrow \mathbf{A}_3$ are $\mathcal{G}$ -homomorphisms. Then the composite*

$$h = A_1 \xrightarrow{h_1} A_2 \xrightarrow{h_2} A_3$$

is a $\mathcal{G}$ -homomorphism $\mathbf{A}_1 \rightarrow \mathbf{A}_3$.

Proof. Define $\rho_i = \ker h_i$, $i = 1, 2$, and $\rho = \ker h$. Suppose that D is a ρ -block and that G is a subgroup of $M_{\mathbf{A}_1}(D)$. We need to show that $G \in \overline{\mathcal{G}}$. By Lemma 2.2, there exists a nonempty set $D_0 \subseteq D$ such that G is isomorphic to a subgroup of $G_{\mathbf{A}_1}(D_0)$. Let

$$Y = \{y_g : g \in G_{\mathbf{A}_1}(D_0)\}.$$

Defining

$$\delta_0(a, y_g) = ag,$$

D_0 becomes the state set of the permutation Y -automaton $\mathbf{D}_0 = (D_0, Y, \delta_0)$. Since h_1 is a $\mathcal{G}$ -homomorphism, the restriction ρ'_1 of ρ_1 to D_0 is a $\mathcal{G}$ -congruence of $\mathbf{D}_0$. Further, $\mathbf{D}_0/\rho'_1$ is a permutation automaton, and since h_2 is a $\mathcal{G}$ -homomorphism, the group $G(\mathbf{D}_0/\rho'_1)$ is in $\overline{\mathcal{G}}$. Thus, by Lemma 4.6, $G(\mathbf{D}_0) \in \overline{\mathcal{G}}$. But the two groups $G(\mathbf{D}_0)$ and $G_{\mathbf{A}_1}(D_0)$ are isomorphic, so that $G_{\mathbf{A}_1}(D_0)$ is also in $\overline{\mathcal{G}}$. $\square$

COROLLARY 4.8 *Suppose that $\rho_1 \leq \rho_2$ are congruence relations of the automaton $\mathbf{A}$. If ρ_1 is a $\mathcal{G}$ -congruence and if ρ_2/ρ_1 is a $\mathcal{G}$ -congruence of $\mathbf{A}/\rho_1$, then ρ_2 is a $\mathcal{G}$ -congruence.*

LEMMA 4.9 *Suppose that $\mathbf{A}$ and $\mathbf{B}$ are X -automata and that h is a simple homomorphism $\mathbf{A} \rightarrow \mathbf{B}$ which is not injective. Then there is an X -automaton $\mathbf{C}$, a surjective simple regular homomorphism $h_1 : \mathbf{A} \rightarrow \mathbf{C}$ and a simple homomorphism $h_2 : \mathbf{C} \rightarrow \mathbf{B}$ such that h_1 is not injective and*

$$h = \mathbf{A} \xrightarrow{h_1} \mathbf{C} \xrightarrow{h_2} \mathbf{B}.$$

Proof. Let ρ be minimal among those congruence relations of $\mathbf{A}$ which collapse the states in at least one non-singleton congruence class of $\ker h$. Then let $\mathbf{C} = \mathbf{A}/\rho$ and let h_1 be the natural homomorphism $\mathbf{A} \rightarrow \mathbf{A}/\rho$. The definition of h_2 is forced. $\square$

REMARK 4.10 By Lemma 4.3 and Lemma 4.7, h is a $\mathcal{G}$ -homomorphism iff h_1 and h_2 are $\mathcal{G}$ -homomorphisms.

COROLLARY 4.11 *Suppose that $\mathbf{A}$ is an X -automaton and ρ is a simple congruence relation of $\mathbf{A}$ other than the identity relation. Then there is a simple regular congruence relation $\rho' \leq \rho$ which is not the identity relation and such that ρ/ρ' is also simple. Further, ρ is a $\mathcal{G}$ -congruence iff both ρ' and ρ/ρ' are $\mathcal{G}$ -congruences.*

5 Two Relations

Throughout this section $\mathcal{G}$ denotes a given class of simple groups closed under division. We define two relations on automata.

DEFINITION 5.1 *Suppose that $\mathbf{A}$ and $\mathbf{B}$ are X -automata. We define:*

- $\mathbf{A} \geq \mathbf{B}$ if there is a surjective $\mathcal{G}$ -homomorphism $\mathbf{A} \rightarrow \mathbf{B}$;
- $\mathbf{A} \succeq \mathbf{B}$ if there is a surjective simple regular $\mathcal{G}$ -homomorphism $\mathbf{A} \rightarrow \mathbf{B}$.

Thus, if $\mathbf{A} \succeq \mathbf{B}$, then $\mathbf{A} \geq \mathbf{B}$. Moreover, both relations are reflexive, and the relation $\geq$ is transitive, by Lemma 4.7. We let $\equiv$ ($\sim$, respectively) denote the smallest equivalence relation containing the relation $\geq$ ($\succeq$, respectively).

LEMMA 5.2 *Suppose that $\mathbf{A}$ and $\mathbf{B}$ are X -automata with $\mathbf{A} \geq \mathbf{B}$. Then $\mathbf{A} \sim \mathbf{B}$.*

Proof. Suppose that ρ is a $\mathcal{G}$ -congruence of the X -automaton $\mathbf{A} = (A, X, \delta)$. We prove that $\mathbf{A} \sim \mathbf{A}/\rho$. We argue by induction on

$$\#\rho = \max\{|\rho(a)| : a \in A\}.$$

The basis case that $\#\rho = 1$ is obvious. Suppose that $\#\rho > 1$. Define the X -automaton $\mathbf{A}' = (A, X, \delta')$ on the set A as follows. For each $a \in A$ and $x \in X$ with $\rho(a)x \subset \rho(ax)$ and $|\rho(ax)| = \#\rho$, let $\delta'(a, x)$ be some fixed element of $\rho(ax) - \rho(a)x$, depending only on $\rho(a)$ and x . Otherwise define $\delta'(a, x) = ax$. (Here, for any $b \in A$, $\rho(b)$ denotes the ρ -block containing b .) Note that ρ is a congruence relation of $\mathbf{A}'$ and $\mathbf{A}/\rho$ is isomorphic to $\mathbf{A}'/\rho$.

Let R denote the set

$$\{(a, b) \in A \times A : a \rho b \text{ \& } (|\rho(a)| < \#\rho \text{ or } a \neq b)\}.$$

Then R determines a subautomaton of the direct product $\mathbf{A} \times \mathbf{A}'$. To prove this, suppose that $(a, b) \in R$ and $x \in X$. We need to show that $(a, b)x \in R$.

CASE 1 $|\rho(ax)| = \#\rho$ and $\rho(a)x = \rho(ax)$. Then $a \neq b$ and x induces in $\mathbf{A}$ a bijection $\rho(a) \rightarrow \rho(ax)$. Thus $(a, b)x = (ax, bx)$ and $ax \neq bx$, proving $(a, b)x \in R$.

CASE 2 $|\rho(ax)| = \#\rho$ and $\rho(a)x \subset \rho(ax)$. Then $bx \neq ax$, since $bx \notin \rho(a)x$. Thus $(a, b)x \in R$.

CASE 3 $|\rho(ax)| < \#\rho$. Then $(a, b)x \in R$ holds obviously.

As noted above, ρ is a congruence relation of $\mathbf{A}'$. We show that ρ is a $\mathcal{G}$ -congruence. For each ρ -block C , $M_{\mathbf{A}'}(C)$ is a submonoid of $M_{\mathbf{A}}(C)^c$, the semigroup obtained by adding the constant maps $C \rightarrow C$ to $M_{\mathbf{A}}(C)$. But since ρ is a $\mathcal{G}$ -congruence of $\mathbf{A}$, each subgroup of $M_{\mathbf{A}}(C)$ is in $\overline{\mathcal{G}}$, moreover, each nontrivial subgroup of $M_{\mathbf{A}}(C)^c$ is a subgroup of $M_{\mathbf{A}}(C)$. Since ρ is a $\mathcal{G}$ -congruence of $\mathbf{A}$, it follows that ρ is a $\mathcal{G}$ -congruence of $\mathbf{A}'$.

The functions

$$\begin{aligned} \pi : R &\rightarrow A, & (a, b) &\mapsto a \\ \pi' : R &\rightarrow A, & (a, b) &\mapsto b \end{aligned}$$

are surjective homomorphisms $\mathbf{R} \rightarrow \mathbf{A}$ and $\mathbf{R} \rightarrow \mathbf{A}'$, respectively, where $\mathbf{R}$ denotes the subautomaton of $\mathbf{A} \times \mathbf{A}'$ determined by the set R . Define $\theta = \ker \pi$ and $\theta' = \ker \pi'$. Then $\#\theta < \#\rho$ and $\#\theta' < \#\rho$. Thus, if π and π' are $\mathcal{G}$ -homomorphisms, then $\mathbf{A} \sim \mathbf{R}$ and $\mathbf{A}' \sim \mathbf{R}$, by the induction assumption, so that

$$\mathbf{A} \sim \mathbf{A}'. \tag{4}$$

To prove that π is a $\mathcal{G}$ -homomorphism, note that each θ -block C is either of the form

$$\{a\} \times \rho(a)$$

or

$$\{a\} \times (\rho(a) - \{a\}),$$

for some $a \in A$. Thus, writing $D = \rho(a)$ or $D = \rho(a) - \{a\}$, $M_{\mathbf{R}}(C)$ is a quotient of the submonoid of $M_{\mathbf{A}'}(\rho(a))$ determined by the functions $g = u^{\mathbf{A}'}|_D$, $u \in X^*$ with $Dg \subseteq D$ and $au^{\mathbf{A}} = a$. Since ρ is a $\mathcal{G}$ -congruence of $\mathbf{A}'$, it follows that each simple group divisor of $M_{\mathbf{R}}(C)$ is in $\mathcal{G}$. Thus θ is a $\mathcal{G}$ -congruence and π is a $\mathcal{G}$ -homomorphism. The proof of the fact that π' is also a $\mathcal{G}$ -homomorphism is similar. Thus (4) has been established.

By (4) and since $\mathbf{A}/\rho$ and $\mathbf{A}'/\rho$ are isomorphic, to complete the proof we need to show that $\mathbf{A}' \sim \mathbf{A}'/\rho$. Let τ denote the congruence relation of $\mathbf{A}'$ whose non-singleton blocks are those ρ -blocks C with $|C| < \#\rho$. Then $\tau \leq \rho$, so that τ is a $\mathcal{G}$ -congruence of $\mathbf{A}'$, by Corollary 4.4. Moreover, $\#\tau < \#\rho$, and ρ/τ is a simple $\mathcal{G}$ -congruence of $\mathbf{A}'/\tau$. Thus, $\mathbf{A}' \sim \mathbf{A}'/\tau$, by the induction assumption. But by Lemma 5.3 below, $\mathbf{A}'/\tau \sim \mathbf{A}'/\rho$, completing the proof. $\square$

LEMMA 5.3 *Suppose that $\mathbf{A}$ and $\mathbf{B}$ are X -automata and h is a surjective simple $\mathcal{G}$ -homomorphism $\mathbf{A} \rightarrow \mathbf{B}$. Then there is chain*

$$\mathbf{A} \succeq \mathbf{A}_1 \succeq \dots \succeq \mathbf{A}_n \succeq \mathbf{B}.$$

Proof. By Lemma 4.9, there exist X -automata $\mathbf{A}_1, \dots, \mathbf{A}_n$ and surjective simple regular $\mathcal{G}$ -homomorphisms

$$\mathbf{A} \xrightarrow{h_0} \mathbf{A}_1 \xrightarrow{h_1} \dots \xrightarrow{h_{n-1}} \mathbf{A}_n \xrightarrow{h_n} \mathbf{B}. \quad \square$$

COROLLARY 5.4 *For any two X -automata $\mathbf{A}$ and $\mathbf{B}$, $\mathbf{A} \sim \mathbf{B}$ iff $\mathbf{A} \equiv \mathbf{B}$.*

6 Proof of Theorem 3.2

In this section we complete our proof of Theorem 3.2.

LEMMA 6.1 *Suppose that $\mathbf{A} = (A, X, \delta)$ is a given automaton and ρ is a simple regular $\mathcal{G}$ -congruence of $\mathbf{A}$, for some class $\mathcal{G}$ of simple groups closed under division. Let K consist of the automata $\mathbf{A}/\rho$ and $\mathbf{U}$ as well as the automata $\mathbf{Aut}(G)$ for $G \in \mathcal{G}$. Then*

$$\mathbf{A} \in \mathbf{V}_c(K).$$

Proof. We may assume that $\#\rho > 1$. Let $C_1, \dots, C_k$, $k > 0$, denote the ρ -blocks C_i with $|C_i| = \#\rho$, and let $D_1 = \{d_1\}, \dots, D_m = \{d_m\}$ be the singleton ρ -blocks. Since ρ is simple, the sets C_i and D_j are all of the ρ -blocks. For each $i \in [k]$ there exist words $u_i, v_i \in X^*$ with $C_1 u_i = C_i$ and $C_i v_i = C_1$, and such that $u_i v_i$ induces the identity function on C_1 and $v_i u_i$ induces the identity function on C_i , so that $(u_i v_i)^{\mathbf{A}}|_{C_1} = \lambda^{\mathbf{A}}|_{C_1}$ and $(v_i u_i)^{\mathbf{A}}|_{C_i} = \lambda^{\mathbf{A}}|_{C_i}$. (We may assume that $u_1 = v_1 = \lambda$).

Define

$$Y = \{y_a : a \in C_1\} \cup \{y_s : s \in S_{\mathbf{A}}(C_1)\}.$$

We turn C_1 into an Y -automaton $\mathbf{C}_1$ by defining

$$\begin{aligned} c y_a &= a \\ c y_s &= cs, \end{aligned}$$

for all $a, c \in C_1$ and $s \in S_{\mathbf{A}}(C_1)$. Then $\mathbf{A} \in \mathbf{IS}(\{\mathbf{B}\})$ holds for the cascade composition

$$\mathbf{B} = \mathbf{A}/\rho \times_{\varphi} \mathbf{C}_1,$$

where

$$\varphi : A/\rho \times X \rightarrow Y$$

is defined as follows. Let a_0 be a fixed element of C_1 . Then, for each $i \in [k]$ and $x \in X$, define

$$\varphi(C_i, x) = \begin{cases} y_s & \text{if } C_i x \subseteq C_j, \text{ where } s = (u_i x v_j)^{\mathbf{A}}|_{C_1} \text{ and } j \in [k]; \\ y_{a_0} & \text{if } C_i x = D_j \text{ for some } j \in [m]. \end{cases}$$

Moreover, for each $i \in [m]$ and $x \in X$, let

$$\varphi(D_i, x) = \begin{cases} y_a & \text{if } d_i x = b \in C_j, j \in [k], a \in C_i \text{ and } au_j = b; \\ y_{a_0} & \text{if } d_i x = d_j, \text{ for some } j \in [m]. \end{cases}$$

Then the set

$$B_0 = \{(C_i, a) : a \in C_1, i \in [k]\} \cup \{(D_j, a_0) : j \in [m]\}$$

determines a subautomaton $\mathbf{B}_0$ of $\mathbf{B}$. Moreover, the function

$$\begin{aligned} h : B_0 &\rightarrow A \\ (C_i, a) &\mapsto au_i \\ (D_j, a_0) &\mapsto d_j \end{aligned}$$

is an isomorphism $\mathbf{B}_0 \rightarrow \mathbf{A}$, as shown by the following commutative squares corresponding to the 4 cases in the definition of φ :

$$\begin{array}{ccc} (C_i, a) & \xrightarrow{h} & au_i \\ \downarrow x & & \downarrow x \\ (C_j, au_i x v_j) & \xrightarrow{h} & au_i x v_j u_j = au_i x \\ \\ (C_i, a) & \xrightarrow{h} & au_i \\ \downarrow x & & \downarrow x \\ (D_j, a_0) & \xrightarrow{h} & d_j \\ \\ (D_i, a_0) & \xrightarrow{h} & d_i \\ \downarrow x & & \downarrow x \\ (C_j, a) & \xrightarrow{h} & b = au_j \\ \\ (D_i, a_0) & \xrightarrow{h} & d_i \\ \downarrow x & & \downarrow x \\ (D_j, a_0) & \xrightarrow{h} & d_j \end{array}$$

To complete the proof, note that C_1 is a permutation-reset automaton and any simple group dividing $M(C_1)$ is in $\mathcal{G}$, since ρ is a $\mathcal{G}$ -congruence. Thus,

$$C_1 \in V_c(\{U, \text{Aut}(G) : G \in \mathcal{G}\}),$$

by Lemma 2.3. It follows that $A \in V_c(K)$. $\square$

REMARK 6.2 The automaton B_0 is a quotient of B under the homomorphism $h' : B \rightarrow B_0$ defined by:

$$\begin{aligned} (C_i, a) &\mapsto (C_i, a) \\ (D_j, a) &\mapsto (D_j, a_0), \end{aligned}$$

for all $i \in [k]$, $j \in [m]$ and $a \in C_1$. The homomorphism h' is simple and aperiodic, and has the property that each (non-singleton) block of $\ker h'$ contains at most one state which is in the range of the transition function of B . Such homomorphisms are termed *elementary* in [4].

Proof of Theorem 3.2. Let $A = (A, X, \delta)$ be an automaton. Recall that the class $\mathcal{K}(A)$ consists of the automaton U as well as the automata $\text{Aut}(G)$ for simple groups G with $G|M(A)$. We need to show that

$$A \in V_c(\mathcal{K}(A)).$$

Let T denote the trivial one-state X -automaton and let $\mathcal{G}$ denote the class of simple groups G with $G|M(A)$. Then, with respect to this class $\mathcal{G}$, $A \geq T$, so that $A \sim T$, by Corollary 5.4. Thus, there exists a sequence of X -automata $B_0, \dots, B_k$ such that $B_0 = T$, $B_k = A$, and for each $i \in \{0, \dots, k-1\}$ either $B_i \succeq B_{i+1}$ or $B_{i+1} \succeq B_i$. We argue by induction on i to show that $B_i \in V_c(\mathcal{K}(A))$. When $i = 0$, this is obvious. For the induction step, suppose that $i > 0$ and $B_{i-1} \in V_c(\mathcal{K}(A))$. If $B_{i-1} \succeq B_i$, then $B_i \in H(\{B_{i-1}\})$, so that $B_i \in V_c(\mathcal{K}(A))$. Suppose that $B_i \succeq B_{i-1}$. Then there is a surjective simple regular $\mathcal{G}$ -homomorphism $h : B_i \rightarrow B_{i-1}$. Thus, by Lemma 6.1,

$$B_i \in V_c(\mathcal{K}(A) \cup B_{i-1}).$$

It follows from the induction assumption that $B_i \in V_c(\mathcal{K}(A))$. $\square$

REMARK 6.3 When A is a permutation automaton and $\#\rho > 1$, there is no singleton ρ -block. We may define $Y = \{y_s : s \in G_A(C_1)\}$, so that C_1 becomes the Y -automaton with $cy_s = cs$, for all $c \in C_1$ and $s \in G_A(C_1)$. Then C_1 is a permutation automaton and $G(C_1)$ is in $\bar{\mathcal{G}}$. Moreover, A is isomorphic to a cascade composition of A/ρ with C_1 .

COROLLARY 6.4 Suppose that $\mathcal{G}$ is a class of simple groups closed under division. Let K consist of U and the automata $\text{Aut}(G)$ for $G \in \mathcal{G}$. Then the following conditions are equivalent for an automaton A :

1. Each simple group divisor of $S(A)$ is in $\mathcal{G}$.
2. There is a sequence of automata $A_0, \dots, A_n$ such that A_0 is trivial, A_n is A , and for each $i \in [n]$, either A_i is a quotient of A_{i-1} under a simple regular $\mathcal{G}$ -homomorphism, or A_{i-1} is a quotient of A_i under a simple regular $\mathcal{G}$ -homomorphism.
3. $A \in V_c(K)$.

4. *A is in the least class of automata containing K and closed under subautomata, simple regular G -homomorphic images, renaming and cascade composition.*
5. *A is in the least class of automata containing K and closed under subautomata, G -homomorphic images, renaming and cascade composition.*

Note This paper was submitted to an editor of Theoretical Computer Science in December 1995. Unfortunately the author has not received any referee report since then.

References

- [1] M.A. Arbib, ed., *Algebraic Theory of Machines, Languages, and Semigroups*, Academic Press, 1968.
- [2] S. Eilenberg, *Automata, Languages, and Machines*, vol. B., Academic Press, 1976.
- [3] Z. Ésik, Results on homomorphic realization of automata by α_0 -products, *Theoretical Computer Science*, 87(1991), 229–249.
- [4] Z. Ésik, Group axioms for iteration, *Information and Computation*, 148(1999), 131–180.
- [5] F. Gécseg, *Products of Automata*, Springer-Verlag, 1986.
- [6] A. Ginzburg, *Algebraic Theory of Automata*, Academic Press, 1968.
- [7] W.M.L. Holcombe, *Algebraic Automata Theory*, Cambridge University Press, 1982.
- [8] K. Krohn and J. Rhodes, The algebraic theory of machines I, *Trans. Amer. Math. Soc.*, 116(1965), 450–464.
- [9] G. Lallement, *Semigroups and Combinatorial Applications*. Wiley-Interscience, 1979.
- [10] H. Straubing, *Finite Automata, Formal Logic, and Circuit Complexity*, Birkhäuser, 1994.