

UNIFORMLY SHADOWING PROPERTY

都立大理 花井 博史 (Hirofumi Hanai)

東京理大理工 岡 正俊 (Masatoshi Oka)

Let M be a closed C^∞ manifold and $C^r(M)$ be the set of all C^r -differentiable maps endowed with the C^r -topology ($r \geq 1$). $D_x f$ is the derivative of f at x . Denote as $\tilde{M}$ the topological product space $\prod_{-\infty}^{\infty} M$ and define a compatible metric $\tilde{d}$ on $\tilde{M}$ by $\tilde{d}((x_n), (y_n)) = \sum_{-\infty}^{\infty} d(x_n, y_n)/2^{|n|}$ for $(x_n), (y_n) \in \tilde{M}$, where d is a metric on M induced by a Riemannian metric. We define a continuous map $\tilde{f} : \tilde{M} \rightarrow \tilde{M}$ by

$$\tilde{f}((x_n)) = (f(x_n)).$$

Then the projection $P^0 : \tilde{M} \rightarrow M$ defined by $P^0((x_n)) = x_0$ satisfies $P^0 \circ \tilde{f} = f \circ P^0$. For a subset Λ an $\tilde{f}$ -invariant set Λ_f is defined by

$$\Lambda_f = \{(x_n) \in \tilde{M} : x_n \in \Lambda, f(x_n) = x_{n+1}, n \in \mathbb{Z}\}.$$

If $\Lambda_f \neq \emptyset$ then $\tilde{f}|_{\Lambda_f} : \Lambda_f \rightarrow \Lambda_f$ is a surjective homeomorphism. Remark that $\Lambda_f = M_f \neq \emptyset$ when $\Lambda = M$. We say that each element of M_f is an orbit of f .

For $\delta \geq 0$ a sequence $\{x^i\}_{i \in \mathbb{Z}} \subset M$ is called a δ -pseudo-orbit of f if $d(f(x^i), x^{i+1}) \leq \delta$ for every $i \in \mathbb{Z}$. A sequence $\{x^i\}_{i \in \mathbb{Z}} \subset M$ is said to be ε -traced by an orbit $(y_i) \in M_f$ if $d(x^i, y_i) < \varepsilon$ for every $i \in \mathbb{Z}$. We say that f has the *shadowing property* if for every $\varepsilon > 0$ there exists $\delta > 0$ such that every δ -pseudo-orbit of f can be ε -traced by an orbit of f .

A sequence $\{\tilde{x}^i\}_{i \in \mathbb{Z}} \subset M_f$ is called an orbit of $\tilde{f}$ if $\tilde{f}(\tilde{x}^i) = \tilde{x}^{i+1}$. For $\delta \geq 0$ a sequence $\{\tilde{x}^i\}_{i \in \mathbb{Z}} \subset M_f$ is a δ -pseudo-orbit of $\tilde{f}$ if $\tilde{d}(\tilde{f}(\tilde{x}^i), \tilde{x}^{i+1}) \leq \delta$ for every $i \in \mathbb{Z}$. A sequence $\{\tilde{x}^i\}_{i \in \mathbb{Z}} \subset \tilde{M}$ is said to be ε -traced by an orbit $(\tilde{y}^i)$ of $\tilde{f}$ if $\tilde{d}(\tilde{x}^i, \tilde{y}^i) < \varepsilon$ for every $i \in \mathbb{Z}$. We say that $\tilde{f}$ satisfies the *shadowing property* if for every $\varepsilon > 0$ there exists $\delta > 0$ such that every δ -pseudo-orbit of $\tilde{f}$ can be ε -traced by an orbit of $\tilde{f}$.

We say that $\tilde{f}$ satisfies C^r *uniformly shadowing property* if there is a neighborhood $\mathcal{U}(f)$ of f in $C^r(M)$ with the property that for $\varepsilon > 0$ there is $\delta > 0$ such that for $g \in \mathcal{U}(f)$ every δ -pseudo-orbit of $\tilde{g}$ is ε -traced by an orbit of $\tilde{g}$.

Let $\pi : TM \rightarrow M$ be a tangent bundle of M and $\|\cdot\|$ be a Riemannian metric on TM . Define a subset of the product topological space $\tilde{M} \times TM$ by

$$T\tilde{M} = \{(\tilde{x}, v) \in \tilde{M} \times TM : P^0(\tilde{x}) = \pi(v)\}$$

and define a Finsler $\|\cdot\|$ on $T\tilde{M}$ by $\|(\tilde{x}, v)\| = \|v\|$. Then $\tilde{\pi} : T\tilde{M} \rightarrow \tilde{M}$ defined by $\tilde{\pi}(\tilde{x}, v) = \tilde{x}$ is a C^0 -vector bundle over $\tilde{M}$. Define the projection $\bar{P}^0 : T\tilde{M} \rightarrow TM$ by $\bar{P}^0(\tilde{x}, v) = v$. Then,

$$\bar{P}^0|_{T_{\tilde{x}}\tilde{M}} : T_{\tilde{x}}\tilde{M} \rightarrow T_{P^0(\tilde{x})}M$$

is a linear isomorphism where $T_{\tilde{x}}\tilde{M} = \tilde{\pi}^{-1}(\tilde{x})$. A linear bundle map $D\tilde{f} : T\tilde{M} \rightarrow T\tilde{M}$ covering $\tilde{f}$ is defined by

$$D\tilde{f}(\tilde{x}, v) = (\tilde{f}(\tilde{x}), D_{P^0(\tilde{x})}f(v)).$$

Then we have $D\tilde{f}(T_{\tilde{x}}\tilde{M}) \subset T_{\tilde{f}(\tilde{x})}\tilde{M}$ and $\bar{P}^0 \circ D\tilde{f} = Df \circ \bar{P}^0$. To simplify the notation we write $D_{\tilde{x}}\tilde{f} = D\tilde{f}|_{T_{\tilde{x}}\tilde{M}}$. For a subset $\tilde{\Lambda}$ define

$$T\tilde{M}|_{\tilde{\Lambda}} = \bigcup_{\tilde{x} \in \tilde{\Lambda}} T_{\tilde{x}}\tilde{M}.$$

A closed f -invariant set Λ ($f(\Lambda) = \Lambda$) is said to be *hyperbolic* if $T\tilde{M}|_{\Lambda_f}$ splits into the Whitney sum $T\tilde{M}|_{\Lambda_f} = \mathbb{E}^s \oplus \mathbb{E}^u$ of subbundles $\mathbb{E}^s$ and $\mathbb{E}^u$, and there are $C > 0$ and $0 < \lambda < 1$ such that

- (i) $D\tilde{f}(\mathbb{E}^s) \subset \mathbb{E}^s$ and $D\tilde{f}(\mathbb{E}^u) = \mathbb{E}^u$,
- (ii) $D\tilde{f}|_{\mathbb{E}^u} : \mathbb{E}^u \rightarrow \mathbb{E}^u$ is invertible,
- (iii) $\|D\tilde{f}^n|_{\mathbb{E}^s}\| \leq C\lambda^n$ and $\|(D\tilde{f}|_{\mathbb{E}^u})^{-n}\| \leq C\lambda^n$ for $n \geq 0$,

where $\|T\|$ denotes the supremum norm of a linear bundle map T . The number λ is called the *skewness* of the hyperbolic set Λ . For $\varepsilon > 0$ and

$\tilde{x} \in M_f$ the *local stable* and the *local unstable manifolds* are defined by

$$W_\varepsilon^s(\tilde{x}, f) = \{y \in M : d(x_n, f^n(y)) \leq \varepsilon \text{ for } n \geq 0\},$$

$$W_\varepsilon^u(\tilde{x}, f) = \left\{ y \in M \left| \begin{array}{l} \text{there exists } \tilde{y} \in M_f \text{ such that } y_0 = y \\ \text{and } d(x_{-n}, y_{-n}) \leq \varepsilon \text{ for } n \geq 0 \end{array} \right. \right\}.$$

Then, $W_\varepsilon^s(\tilde{x}, f) = W_\varepsilon^s(\tilde{y}, f)$ for $\tilde{x}, \tilde{y} \in M_f$ with $x_0 = y_0$.

For $\tilde{x} \in M_f$ the *stable* and the *unstable sets* are defined by

$$W^s(\tilde{x}, f) = \{y \in M : \lim_{n \rightarrow \infty} d(x_n, f^n(y)) = 0\},$$

$$W^u(\tilde{x}, f) = \left\{ y \in M \left| \begin{array}{l} \text{there is } \tilde{y} \in M_f \text{ satisfying } y_0 = y \\ \text{and } \lim_{n \rightarrow \infty} d(x_{-n}, y_{-n}) = 0 \end{array} \right. \right\}.$$

Then, $W^s(\tilde{x}, f) = W^s(\tilde{y}, f)$ for $\tilde{x}, \tilde{y} \in M_f$ with $x_0 = y_0$. If Λ is a hyperbolic set, for $\tilde{x} \in \Lambda_f$ we have

$$W^s(\tilde{x}, f) = \bigcup_{n=0}^{\infty} f^{-n}(W_\varepsilon^s(\tilde{f}^n(\tilde{x}), f)), \quad W^u(\tilde{x}, f) = \bigcup_{n=0}^{\infty} f^n(W_\varepsilon^u(\tilde{f}^{-n}(\tilde{x}), f)).$$

Remark that $W^\sigma(\tilde{x}, f)$ ($\sigma = s, u$) are not always the manifolds like the stable and unstable manifolds given by diffeomorphisms. However we can define the transversality condition between $W^s(\tilde{x}, f)$ and $W^u(\tilde{y}, f)$ as follows.

Let $\tilde{y}$ and $\tilde{z}$ be points in Λ_f . We say that $W^s(\tilde{y}, f)$ is *transversal* to $W^u(\tilde{z}, f)$ if $f^{n+m} | W_\varepsilon^u(\tilde{f}^{-m}(\tilde{z}), f)$ is transversal to $W_\varepsilon^s(\tilde{f}^n(\tilde{y}), f)$ for $\varepsilon > 0$ small enough and $n, m \geq 0$.

The non-wandering set $\Omega(f)$ is defined by

$$\Omega(f) = \left\{ x \in M \left| \begin{array}{l} \text{for any neighborhood } U \text{ of } x \text{ there is } n > 0 \\ \text{satisfying } U \cap f^n(U) \neq \emptyset \end{array} \right. \right\}.$$

Obviously, $\Omega(f)$ is closed and satisfies that $f(\Omega(f)) \subset \Omega(f)$ and $Per(f) \subset \Omega(f)$, where $Per(f)$ denotes the set of all periodic points of f . A differentiable map f is said to satisfy *Axiom A* if

- (i) $Per(f)$ is dense in $\Omega(f)$,
- (ii) $\Omega(f)$ is hyperbolic.

We say that an Axiom A differentiable map f satisfies the *strong transversality* if $W^s(\tilde{y}, f)$ is transversal to $W^u(\tilde{z}, f)$ for $\tilde{y}, \tilde{z} \in \Omega(f)_f$.

Theorem. If C^1 -differentiable map f satisfies both Axiom A and the strong transversality, then $\tilde{f}$ satisfies C^1 uniformly shadowing property.

This result was proved by Sakai for the class of C^1 -diffeomorphisms. The full proof of our theorem will appear elsewhere.

REFERENCES

- [1] N.Aoki, K.Moriyasu and M.Oka, Differentiable maps having hyperbolic sets, *Topology Appl.* 82(1998) 15 – 48.
- [2] N.Aoki, K.Moriyasu and N.Sumii, C^1 maps having hyperbolic periodic points, to appear in *Fund. Math.*
- [3] J.Guckenheimer, J.Moser and S.Newhouse, *Dynamical Systems* (Birkhäuser, Boston, 1980).

- [4] M.Hirsch, J.Palis, C.Pugh and M.Shub, Neighborhoods of hyperbolic sets, *Inv. Math.* 9 (1970) 121–134.
- [5] M.Hirsch, C.Pugh and M.Shub, Invariant manifolds, *Lecture Note in Math.* 583 (Springer, Berlin, 1977).
- [6] K.Moriyasu, The topological stability of diffeomorphisms, *Nagoya Math. J.* 123 (1991) 91–102.
- [7] K.Moriyasu, Axiom A endomorphisms having no cycle, in: N.Aoki, K.Shiraiwa, Y.Takahashi, ed., *Proceedings of the International Conference on Dynamical Systems and Chaos 1* (World Scientific, Singapore, 1995) 182–186.
- [8] K.Moriyasu and M.Oka, The creation of homoclinic points of C^1 -maps, *Topology Appl.* 54 (1993) 47–64.
- [9] K.Moriyasu and M.Oka, Pseudo orbit tracing property and strong transversality of C^1 maps, in: N.Aoki, K.Shiraiwa, Y.Takahashi, ed., *Proceedings of the International Conference on Dynamical Systems and Chaos 1* (World Scientific, Singapore, 1995) 187–191.
- [10] F.Przytycki, Anosov endomorphisms, *Studia Math.* 58 (1976) 249–285.
- [11] F.Przytycki, On Ω -stability and structural stability of endomorphisms satisfying Axiom A, *Studia Math.* 60 (1977) 61–77.
- [12] C.Robinson, Stability theorem and hyperbolicity in dynamical systems, *Rocky Mountain J. Math.* 7 (1977) 425–437.
- [13] C.Robinson, Structural stability of C^1 diffeomorphisms, *J. Diff. Equat.* 22 (1976) 28–73.
- [14] K.Sakai, The C^1 uniform pseudo-orbit tracing property, *Tokyo J. Math.* 15 (1992) 99–109.
- [15] K.Sakai, C^r Axiom A maps having strong transversality, *Topology Appl.* 54 (1993) 173–182.
- [16] K.Sakai, Pseudo orbit tracing property and strong transversality, *Osaka J. Math.* 31 (1994) 373–386.