
 ABSTRACTS (MASTER THESIS)

**Effects of copper on oxalate biosynthesis
in the brown-rot fungus *Fomitopsis palustris***

(Graduate School of Agriculture, Laboratory of Metabolic Science of Forest Plants and Microorganisms, RISH, Kyoto University)

Hiromichi Hisamori

Wood-rotting basidiomycetes cause severe damage on wooden structures. To protect the wooden structures from the wood-rotting fungi, copper-containing wood preservatives have been used. However, many species of wood-rotting brown-rot fungi can degrade even the wood materials treated with the copper-containing preservatives, by which these wood-rotting fungi are called copper-tolerant fungi. The copper-tolerant ability has been recognized as being associated primarily with oxalic acid excretion, in which oxalic acid produced by copper-tolerant fungi reacts with copper in wood to form insoluble, bio-unavailable inert forms.

From a viewpoint of wood protection from the fungal degradation, the copper-tolerance of these wood-rotting fungi should be diminished. At the same time, the copper-tolerant wood-rotting fungi are promising to be used for bioremediation of wastes of woods treated with copper-containing preservatives [1]. Therefore, it is important to elucidate effects of copper on oxalate biosynthesis to develop bioremediation of copper from wood waste containing wood preservatives including copper.

Brown-rot fungus *Fomitopsis palustris* possesses two metabolic pathways for oxalate biosynthesis: one is hydrolysis of oxaloacetate catalyzed by oxaloacetate acetylhydrolase (*Fomitopsis palustris* oxaloacetate acetylhydrolase, FpOAH) in cytosol and the other is dehydrogenation of glyoxylate catalyzed by cytochrome *c* dependent glyoxylate dehydrogenase (*Fomitopsis palustris* glyoxylate dehydrogenase, FpGLOXDH) in peroxisome [2-4].

This author investigated effects of Cu^{2+} on oxalate biosynthesis including expressions of FpOAH and FpGLOXDH. In the absence of Cu^{2+} , amounts of FpOAH transcripts were 22 – 140 times greater than those of FpGLOXDH. The results suggest that FpOAH plays the more significant role than FpGLOXDH and the pathway including FpOAH as a key enzyme is a major pathway for oxalate biosynthesis, supporting our proposed idea that oxalate is biosynthesized mainly in the cytosol by FpOAH but not in the peroxisome by FpGLOXDH [2-4]. On the other hand, under the condition in the presence of Cu^{2+} an amount of FpOAH transcript was 19.4 times greater on day 4 (minimal magnitude) and 151.1 times greater on day 9 (maximal magnitude) than those of FpGLOXDH. The results suggest that FpOAH also plays a major role in oxalate biosynthesis regardless of absence or presence of Cu^{2+} .

However, the amounts of FpOAH and FpGLOXDH transcripts increased 2.2 and 7 times as their maximal rates by the presence of Cu^{2+} . The results indicate that Cu^{2+} increased expressions of FpOAH and FpGLOXDH, but the rates of the increments are suggested to be just several times.

References

- [1] Kartal, S., and Imamura, Y. (2003) “Chemical and biological remediation of CCA-treated waste wood”, *Wood Research* 90, 111-115.
- [2] Munir, E., Yoon, J. J., Tokimatsu, T., Hattori, T., and Shimada, M. (2001) “A physiological role for oxalic acid biosynthesis in the wood-rotting basidiomycete *Fomitopsis palustris*”, *PNAS* 98:11126-11130.
- [3] Hattori, T., Okawa, K., Fujimura, M., Mizoguchi, M., Watanabe, T., Tokimatsu, T., Inui, H., Baba, K., Suzuki, S., Umezawa, T., and Shimada, M. (2007) “Subcellular localization of the oxalic acid-producing enzyme, cytochrome *c*-dependent glyoxylate dehydrogenase in brown-rot fungus *Fomitopsis palustris*”, *Cellulose Chemistry and Technology*, 41: 545-553.
- [4] Sakai, S., Nishide, T., Munir, E., Baba, K., Inui, H., Nakano, Y., Hattori, T. and Shimada, M. (2006) “Subcellular localization of glyoxylate cycle key enzymes involved in oxalate biosynthesis of wood-destroying basidiomycete *Fomitopsis palustris* grown on glucose”, *Microbiology*, 152: 1857-1866.