

DMRG Studies for Itinerant Ferromagnetism in a Two-leg Optical Lattice System

M. Okumura^{1,2}, S. Yamada^{3,2}, M. Machida^{3,2}, and H. Aoki⁴

¹ CCMP, RIKEN, Wako, Saitama 351-0198, JAPAN

² CREST(JST), Kawaguchi, Saitama 332-0012, JAPAN

³ CCSE, Japan Atomic Energy Agency, Taito-ku, Tokyo 110-0015, JAPAN

⁴ Department of Physics, The University of Tokyo, Hongo, Tokyo 113-0033, Japan

Density matrix renormalization group (DMRG) is one of the most powerful methods for investigating strongly correlated systems. Recently, three of the present authors have developed a parallel DMRG code [1] that enables us to calculate the ground state of n -leg lattice models efficiently. We can thus explore multi-leg ladder systems to look for intriguing physics that is specific to ladder structures, e.g., ferromagnetism in the Hubbard model.

It has been well-known that the lattice structure in addition to strong on-site repulsion is responsible for non-trivial ground state. A typical example is the itinerant ferromagnetism, which still deserves full understanding. Besides electron systems, cold Fermionic atom systems is an ideal test-bench for such studies due to its flexibility and controllability on lattice structures. In fact, three itinerant ferromagnetism, i.e., Stoner's, Nagaoka's, and flat-band mechanisms have been so far discussed in ultra-cold atom experiments, and the first one has been reported very recently [2].

With this background, here we propose that a two-leg ladder optical lattice system should accommodate a stable itinerant ferromagnetism for cold fermionic atoms with finite hole densities [3], which may be regarded as an extended Nagaoka's ferromagnetism. A special interest in the cold atom system is that, on top of the optical lattice potential, we have a trapping potential. We predict that this causes an interesting phase separation into magnetic and nonmagnetic phases in real space. We show the phase-separated magnetism specially in spin-imbalanced situations, and the phase diagram obtained here indicates that the required strength of correlation is realistic.

References

- [1] S. Yamada, M. Okumura, and M. Machida, J. Phys. Soc. Jpn. **78**, 094004 (2009).
- [2] G.-B. Jo *et al.*, Science **325**, 1521 (2009).
- [3] M. Okumura, S. Yamada, M. Machida, and H. Aoki, arXiv:1008.3055.