

An approximation theorem to a solution to an equilibrium problem in complete CAT(1) spaces

完備 CAT(1) 空間における均衡問題の解の近似定理

東邦大学・理学部 木村泰紀

Yasunori Kimura

Department of Information Science

Toho University

東邦大学・大学院理学研究科 大口智輝

Tomoki Oguchi

Department of Information Science

Toho University

1 Introduction

Let K be a nonempty set and $f: K \times K \rightarrow \mathbb{R}$ a function. Equilibrium problems are defined as follows: Find $z_0 \in K$ such that

$$f(z_0, y) \geq 0$$

for all $y \in K$.

In 1994, Blum and Oettli defined a mapping called resolvent for equilibrium problems on Banach spaces. In 2005, Combettes and Hirstoaga showed many important properties about resolvents of equilibrium problems in Hilbert spaces and made contribution to the development of the approximation methods for equilibrium problem. The following is one of the most important theorems.

Theorem 1 (Combettes–Hirstoaga [?]). *Let H be a Hilbert space, and K a nonempty, closed convex subset of H . Suppose that $f: K \times K \rightarrow \mathbb{R}$ satisfies the conditions (E1)–(E4).*

(E1) $f(x, x) = 0$ for all $x \in K$;

(E2) $f(x, y) + f(y, x) \leq 0$ for all $x, y \in K$;

(E3) $f(x, \cdot): K \rightarrow \mathbb{R}$ is lower semicontinuous and convex for all $x \in K$;

(E4) $f(\cdot, y): K \rightarrow \mathbb{R}$ is upper hemicontinuous for all $y \in K$.

The set of solutions to the equilibrium problem is denoted by $\text{Equil } f$, that is,

$$\text{Equil } f = \left\{ x \in K \mid \inf_{y \in K} f(x, y) \geq 0 \right\}.$$

Define the resolvent J_f by

$$J_f x = \left\{ z \in K \mid \inf_{y \in K} (f(z, y) + \langle z - x, y - z \rangle) \geq 0 \right\}$$

for $x \in H$. Then J_f has the following properties:

1. The domain of J_f is H ;
2. J_f is single-valued and firmly nonexpansive;
3. the set of all fixed points of J_f coincides with $\text{Equil } f$ and it is closed and convex.

A basic approach to solving fixed point problems in general normed spaces is Mann's type iteration.

Further in 2018, Kimura and Kishi proposed the resolvent of $\text{CAT}(0)$ space. On the other hand, in 2012, He, Fang, López and Li showed a convergence theorem to a fixed point of a nonexpansive mapping in a $\text{CAT}(\kappa)$ spaces with Mann's type iteration.

In this paper, we obtain a convergence theorem of an iterative sequence to a solution of an equilibrium problem on a $\text{CAT}(1)$ space. We use Mann's type iterative method to generate the approximate sequence.

2 Preliminaries

Let $x, y \in X$ and γ a mapping of $[0, d(x, y)]$ into X . A mapping γ is called a geodesic with endpoints x and y if $\gamma(0) = x$, $\gamma(d(x, y)) = y$, $d(\gamma(u), \gamma(v)) = |u - v|$ for all $u, v \in [0, d(x, y)]$. X is called a uniquely π -geodesic space if for any $x, y \in X$ with $d(x, y) < \pi$, a geodesic with endpoints x and y exists uniquely. For $x, y \in X$ and $t \in [0, 1]$, there exists $z \in [x, y]$ such that $d(x, z) = (1 - t)d(x, y)$ and $d(z, y) = td(x, y)$ which is denoted by $tx \oplus (1 - t)y$. X is called a $\text{CAT}(1)$ space if $\cos d(tx \oplus (1 - t)y, z) \sin d(x, y) \geq \cos d(x, z) \sin td(x, y) + \cos d(y, z) \sin(1 - t)d(x, y)$ for all $x, y, z \in X$ and $t \in [0, 1]$. A $\text{CAT}(1)$ space X is said to be admissible if $d(u, v) < \pi/2$ for any $u, v \in X$.

Let X be a metric space and $T: X \rightarrow X$. We call $x \in X$ a fixed point of T if $x = Tx$, and denote the set of all fixed points of T by $F(T)$. An admissible complete $\text{CAT}(1)$ space X has the convex hull finite property if every continuous selfmapping on $\text{clco } E$ has a fixed point for every finite subset E of X , where $\text{clco } E$ is the closure of the convex hull of X .

Theorem 2.1 (Kimura [?]). *Let X be an admissible complete $\text{CAT}(1)$ space having the convex hull finite property and $K \subset X$ a nonempty closed convex set. Suppose that $f: K \times K \rightarrow \mathbb{R}$ satisfies the conditions (E1)–(E4). Define the resolvent R_f of f*

by

$$R_fx = \left\{ z \in K \mid \inf_{y \in K} (f(z, y) - \log \cos d(x, y) + \log \cos d(x, z)) \geq 0 \right\}$$

for all $x \in X$. Then the following hold:

1. $R_f: X \rightarrow K$ is defined as a single-valued mapping;
2. R_f satisfies the following inequality for any $x, y \in X$:

$$\frac{\cos d(x, R_f y)}{\cos d(x, R_f x)} + \frac{\cos d(y, R_f x)}{\cos d(y, R_f y)} \leq 2 \cos d(R_f x, R_f y);$$

3. $F(R_f) = \text{Equil } f$ and it is closed and convex.

Let X be $\text{CAT}(1)$ space and $\{x_n\} \subset X$ a sequence. An asymptotic center $\text{AC}(\{x_n\})$ of $\{x_n\}$ is defined by

$$\text{AC}(\{x_n\}) = \left\{ z \in X \mid \inf_{x \in X} \limsup_{n \rightarrow \infty} d(x, x_n) = \limsup_{n \rightarrow \infty} d(z, x_n) \right\}.$$

A sequence $\{x_n\} \subset X$ is said to be Δ -convergent to $x_0 \in X$ if $\text{AC}(\{x_{n_i}\}) = \{x_0\}$ for all subsequences $\{x_{n_i}\}$ of $\{x_n\}$. It is denoted by $x_n \xrightarrow{\Delta} x_0$.

3 Main result

We begin with the basic result of the resolvent on $\text{CAT}(1)$ spaces.

Theorem 3.1 (Kimura [?]). *Let X be an admissible complete $\text{CAT}(1)$ space having the convex hull finite property. Let K is nonempty closed convex subset of X , suppose that $f: K \times K \rightarrow \mathbb{R}$ satisfies (E1)–(E4). Let $R_f: X \rightarrow K$ be the resolvent of f . Then the following the inequality*

$$\lambda \frac{\cos d(v, R_{\lambda f} u)}{\cos d(v, R_{\mu f} v)} + \mu \frac{\cos d(u, R_{\mu f} v)}{\cos d(u, R_{\lambda f} u)} \leq (\lambda + \mu) \cos d(R_{\lambda f} u, R_{\mu f} v)$$

holds for all $\lambda, \mu \in]0, \infty[$ and $u, v \in X$.

Lemma 3.1. *Let X be an admissible complete $\text{CAT}(1)$ space having the convex hull finite property. Let K is nonempty closed convex subset of X , suppose that $f: K \times K \rightarrow \mathbb{R}$ satisfies (E1)–(E4). Let $\{\lambda_n\} \subset]0, \infty[$ and $\{x_n\} \subset X$ be sequences satisfying $\inf_n \lambda_n > 0$, $x_n \xrightarrow{\Delta} x_0$ and $d(x_n, R_{\lambda_n f} x_n) \rightarrow 0$. Then, $x_0 \in \text{Equil } f$.*

Proof. Let $\{n_i\} \subset \mathbb{N}$ be an increasing sequence and $z \in X$. Since

$$\begin{aligned} d(z, x_{n_i}) &\leq d(z, R_{\lambda_{n_i} f} x_{n_i}) + d(R_{\lambda_{n_i} f} x_{n_i}) \\ &\leq d(z, x_{n_i}) + 2d(R_{\lambda_{n_i} f} x_{n_i}), \end{aligned}$$

we get

$$\limsup_{i \rightarrow \infty} d(z, x_{n_i}) = \limsup_{i \rightarrow \infty} d(z, R_{\lambda_{n_i}} f x_{n_i}).$$

We first suppose that $\sup_{n \in \mathbb{N}} \lambda_n < \infty$. There exists $\{\lambda_{n_j}\} \subset \{\lambda_n\}$ such that $\lambda_{n_j} \rightarrow \lambda_0 \in [\inf_{n \in \mathbb{N}} \lambda_n, \sup_{n \in \mathbb{N}} \lambda_n]$. Fix $j \in \mathbb{N}$. Then,

$$\lambda_{n_j} \cos d(x_0, R_{\lambda_{n_j}} f x_{n_j}) + \cos d(x_{n_j}, R_f x_0) \leq (\lambda_{n_j} + 1) \cos d(R_{\lambda_{n_j}} f x_{n_j}, R_f x_0).$$

Letting $j \rightarrow \infty$, we have

$$\begin{aligned} \lambda_0 \liminf_{j \rightarrow \infty} \cos d(x_0, R_{\lambda_{n_j}} f x_{n_j}) + \liminf_{j \rightarrow \infty} \cos d(x_{n_j}, R_f x_0) \\ \leq \liminf_{j \rightarrow \infty} (\lambda_{n_j} + 1) \cos d(R_{\lambda_{n_j}} f x_{n_j}, R_f x_0) \end{aligned}$$

and hence

$$\begin{aligned} \lambda_0 \cos \limsup_{j \rightarrow \infty} d(x_0, R_{\lambda_{n_j}} f x_{n_j}) + \cos \limsup_{j \rightarrow \infty} d(x_{n_j}, R_f x_0) \\ \leq (\lambda_0 + 1) \cos \limsup_{j \rightarrow \infty} d(R_{\lambda_{n_j}} f x_{n_j}, R_f x_0). \end{aligned}$$

Thus,

$$\begin{aligned} \lambda_0 \cos \limsup_{j \rightarrow \infty} d(x_0, x_{n_j}) + \cos \limsup_{j \rightarrow \infty} d(x_{n_j}, R_f x_0) \\ \leq (\lambda_0 + 1) \cos \limsup_{j \rightarrow \infty} d(x_{n_j}, R_f x_0). \end{aligned}$$

and therefore

$$\cos \limsup_{j \rightarrow \infty} d(x_0, x_{n_j}) \leq \cos \limsup_{j \rightarrow \infty} d(x_{n_j}, R_f x_0).$$

Consequently, we have

$$\limsup_{j \rightarrow \infty} d(x_0, x_{n_j}) \geq \limsup_{j \rightarrow \infty} d(x_{n_j}, R_f x_0).$$

Since $\{x_0\} = \text{AC}(\{x_{n_j}\})$, we have $x_0 = R_f x_0$. It means that $x_0 \in \text{Equil } f$.

We next suppose that $\sup_{n \in \mathbb{N}} \lambda_n = \infty$. Then, there exists $\{\lambda_{n_j}\} \subset \{\lambda_n\}$ such that $\lambda_{n_j} \rightarrow \infty$. Fix $j \in \mathbb{N}$. Then,

$$\lambda_{n_j} \cos d(x_0, R_{\lambda_{n_j}} f x_{n_j}) + \cos d(x_{n_j}, R_f x_0) \leq (\lambda_{n_j} + 1) \cos d(R_{\lambda_{n_j}} f x_{n_j}, R_f x_0).$$

and thus

$$\cos d(x_0, R_{\lambda_{n_j}} f x_{n_j}) + \frac{\cos d(x_{n_j}, R_f x_0)}{\lambda_{n_j}} \leq \left(1 + \frac{1}{\lambda_{n_j}}\right) \cos d(R_{\lambda_{n_j}} f x_{n_j}, R_f x_0).$$

Letting $j \rightarrow \infty$, we get

$$\liminf_{j \rightarrow \infty} d \cos d(x_0, R_{\lambda_{n_j}} f x_{n_j}) \leq \liminf_{j \rightarrow \infty} d \cos d(R_f x_0, R_{\lambda_{n_j}} f x_{n_j})$$

and hence

$$\limsup_{j \rightarrow \infty} d(x_0, R_{\lambda_{n_j}} f x_{n_j}) \geq \limsup_{j \rightarrow \infty} d(R_f x_0, R_{\lambda_{n_j}} f x_{n_j}).$$

It implies that

$$\limsup_{j \rightarrow \infty} d(x_0, x_{n_j}) \geq \limsup_{j \rightarrow \infty} d(R_f x_0, x_{n_j}).$$

Since $\{x_0\} = \text{AC}(\{X_{n_j}\})$, we have $x_0 = R_f x_0$. Therefore $x_0 \in \text{Equil } f$. $\square$

Theorem 3.2. *Let X be an admissible complete CAT(1) space having the convex hull finite property and $K \subset X$ a nonempty closed convex set. Suppose that $f: K \times K \rightarrow \mathbb{R}$ satisfies the conditions (E1)–(E4), and $\text{Equil } f \neq \emptyset$.*

Let $\{\lambda_n\} \subset]0, \infty[$ and $\{\alpha_n\} \subset [0, 1[$ be sequences satisfying $\inf_n \lambda_n > 0$ and $\sup_n \alpha_n < 1$. Let $R_{\lambda_n} f: X \rightarrow K$ be the resolvent of $\lambda_n f$ for each $n \in \mathbb{N}$. Let $\{x_n\}$ be a sequence defined by $x_1 \in X$, and

$$x_{n+1} = \alpha_n x_n \oplus (1 - \alpha_n) R_{\lambda_n} f x_n$$

for each $n \in \mathbb{N}$. Then, $x_n \xrightarrow{\Delta} x_0 \in \text{Equil } f$.

Proof. Since $\text{Equil } f \neq \emptyset$, let $u \in \text{Equil } f$. From Theorem ??,

$$\frac{\cos d(u, R_{\lambda_n} f x_n)}{\cos d(u, R_{\lambda_n} f u)} + \frac{\cos d(x_n, R_{\lambda_n} f u)}{\cos d(x_n, R_{\lambda_n} f x_n)} \leq 2 \cos d(R_{\lambda_n} f u, R_{\lambda_n} f x_n).$$

Since $u \in \text{Equil } f$, we have $d(u, R_{\lambda_n} f u) = 0$ and thus

$$\cos d(u, R_{\lambda_n} f x_n) + \frac{\cos d(x_n, u)}{\cos d(x_n, R_{\lambda_n} f x_n)} \leq 2 \cos d(u, R_{\lambda_n} f x_n).$$

It implies that $\cos d(u, R_{\lambda_n} f x_n) \cos d(x_n, R_{\lambda_n} f x_n) \geq \cos d(x_n, u)$ and hence $d(u, R_{\lambda_n} f x_n) \leq d(u, x_n)$. From the parallelogram law, we get

$$\cos d(u, x_{n+1}) \geq \alpha_n \cos d(u, x_n) + (1 - \alpha_n) \cos d(u, R_{\lambda_n} f x_n) \geq \cos d(u, x_n)$$

and hence

$$d(u, x_{n+1}) \leq d(u, x_n) \leq d(u, x_1) < \frac{\pi}{2}.$$

Therefore $\{x_n\}$ is spherically bounded and $d(u, x_n) \rightarrow c \in [0, \frac{\pi}{2}[$.

Further, we have

$$\begin{aligned}
 \cos d(u, x_{n+1}) &\geq \alpha_n \cos d(u, x_n) + (1 - \alpha_n) \cos d(u, R_{\lambda_n f} x_n) \\
 &\geq \alpha_n \cos d(u, x_n) + (1 - \alpha_n) \frac{\cos d(u, x_n)}{\cos d(R_{\lambda_n f} x_n, x_n)} \\
 &= \cos d(u, x_n) + (1 - \alpha_n) \cos d(u, x_n) \left(\frac{1}{\cos d(R_{\lambda_n f} x_n, x_n)} - 1 \right),
 \end{aligned}$$

which implies that

$$0 \leq (1 - \alpha_n) \left(\frac{1}{\cos d(R_{\lambda_n f} x_n, x_n)} - 1 \right) \leq \frac{\cos d(u, x_{n+1})}{\cos d(u, x_n)} - 1.$$

Since $d(u, x_n) \rightarrow c \in [0, \frac{\pi}{2}[$ and $\sup_{n \in \mathbb{N}} \alpha_n < 1$, we know $d(R_{\lambda_n f} x_n, x_n) \rightarrow 0$. Since $\{x_n\}$ is spherically bounded, any subsequence $\{x_{n_i}\}$ is spherically bounded. Let $\{x_0\} = \text{AC}(\{x_n\})$ and $\{w_0\} = \text{AC}(\{x_{n_i}\})$. We can take a subsequence $\{x_{n_{i_j}}\}$ such that $x_{n_{i_j}} \xrightarrow{\Delta} z_0 \in X$. From Lemma ?? and since $d(R_{\lambda_n f} x_{n_{i_j}}, x_{n_{i_j}}) \rightarrow 0$, we obtain $z_0 \in \text{Equil } f$. Further, we have

$$\begin{aligned}
 \limsup_{n \rightarrow \infty} d(x_n, z_0) &= \limsup_{j \rightarrow \infty} d(x_{n_{i_j}}, z_0) \\
 &\leq \limsup_{j \rightarrow \infty} d(x_{n_{i_j}}, w_0) \\
 &\leq \limsup_{i \rightarrow \infty} d(x_{n_i}, w_0) \\
 &\leq \limsup_{i \rightarrow \infty} d(x_{n_i}, x_0) \\
 &\leq \limsup_{n \rightarrow \infty} d(x_n, x_0) \leq \limsup_{n \rightarrow \infty} d(x_n, z_0).
 \end{aligned}$$

Therefore $x_0 = w_0 = z_0$ and thus we get $\{x_0\} = \text{AC}(\{x_{n_i}\})$ for all $\{x_{n_i}\} \subset \{x_n\}$. Consequently, $x_n \xrightarrow{\Delta} x_0 \in \text{Equil } f$. $\square$

Acknowledgement. This work was partially supported by JSPS KAKENHI Grant Number JP21K03316.

References

- [1] P. L. Combettes and S. A. Hirstoaga, *Equilibrium programming in Hilbert spaces*, J. Nonlinear Convex Anal. **6** (2005), 117–136.
- [2] R. Espínola and Frenández-León, *CAT(k)-spaces, weak convergence and fixed points*, J. Math. Anal. Appl. **353** (2009), 410–427.
- [3] J. S. He, D. H. Fang, G. López and C. Li, *Mann's algorithm for nonexpansive mappings in CAT(κ) spaces*, Nonlinear Anal. **75** (2012), 445–452.

- [4] Y. Kimura, *Resolvents of equilibrium problems on a complete geodesic space with curvature bounded above*, Carpathian J. Math. **37** (2021), 463–476.
- [5] Y. Kimura, *Note on the resolvent for equilibrium problems in a complete geodesic space*, preprint.
- [6] Y. Kimura and Y. Kishi, *Equilibrium problems and their resolvents in Hadamard spaces*, J. Nonlinear Convex Anal. **19** (2018), 1503–1513.