

Lifting Galois representations over arbitrary number fields: A resume

By

Yoshiyuki TOMIYAMA*

Let $\mathbf{k}$ be a finite field of characteristic $p > 5$. Let K be a number field of finite degree over $\mathbb{Q}$ and G_K its absolute Galois group $\text{Gal}(\bar{K}/K)$. Let $\bar{\rho}: G_K \rightarrow \text{GL}_2(\mathbf{k})$ be a continuous representation and let $W(\mathbf{k})$ be the ring of Witt vectors of $\mathbf{k}$. We consider the following question:

Question 0.1. Is there a continuous representation $\rho: G_K \rightarrow \text{GL}_2(W(\mathbf{k}))$ satisfying $\bar{\rho} = \rho \pmod{p}$?

This question has been motivated by a conjecture of Serre ([S]) saying that all odd absolutely irreducible continuous representations $\bar{\rho}: G_{\mathbb{Q}} \rightarrow \text{GL}_2(\mathbf{k})$ are modular. This implies the existence of a lift to characteristic zero. This conjecture was proved by Khare and Wintenberger in [KW1, KW2].

In [K], Khare proved the existence of a lift to $W(\mathbf{k})$ for any $\bar{\rho}: G_K \rightarrow \text{GL}_2(\mathbf{k})$ which is reducible. Thus we may assume that $\bar{\rho}$ is irreducible.

For a place v of K , let K_v be the completion of K at v , and let G_v be its absolute Galois group $\text{Gal}(\bar{K}_v/K_v)$. Let $\text{Ad}^0 \bar{\rho}$ be the $\mathbf{k}$ -vector space of all trace zero two-by-two matrices over $\mathbf{k}$ on which G_K acts by conjugation. Our main result is the following:

Theorem 0.2. Assume that $H^2(G_v, \text{Ad}^0 \bar{\rho}) = 0$ for each place $v \mid p$. Then $\bar{\rho}$ lifts to a continuous representation $\rho: G_K \rightarrow \text{GL}_2(W(\mathbf{k}))$ which is unramified outside a finite set of places of K .

For $K = \mathbb{Q}$, Ramakrishna proved under very general conditions on $\bar{\rho}$ that there exist lifts to $W(\mathbf{k})$ in [R1, R2]. Gee ([G]) and Manoharmayum ([M]) proved, independently, that there exist lifts to $W(\mathbf{k})$ for K satisfying $[K(\mu_p) : K] > 2$, where μ_p is the group of p -th roots of unity.

Our method used in the proof of the Theorem is essentially that of Ramakrishna [R1, R2], but we follow the more axiomatic treatment presented in [Ta]. We denote by S

Received March 23, 2009. Revised June 1, 2009.

2000 Mathematics Subject Classification(s): 11F80

*Graduate School of Mathematics, Kyushu University, Fukuoka 812-8581, Japan.

e-mail: tomiyama@math.kyushu-u.ac.jp

a finite set of places of K containing the places above p , the infinite places and the places at which $\bar{\rho}$ is ramified. Let K_S denote the maximal algebraic extension of K unramified outside S and put $G_{K,S} = \text{Gal}(K_S/K)$. Thus $\bar{\rho}$ factors through $G_{K,S}$. The existence of a lifting of $\bar{\rho}$ follows from the triviality of a certain class of $H^2(G_{K,S}, \text{Ad}^0 \bar{\rho})$. Since it is difficult to calculate it directly, we reduce the calculations of global obstructions to those of local obstructions, which are much better understood. By Taylor [Ta], it boils down to showing the vanishing of a certain Selmer group. We can prove the triviality of this group by extending S by a suitably chosen finite set Q . With the choice of Q of Ramakrishna and Gee, they could not prove the triviality of the Selmer group for an arbitrary number field. Our choice of Q is different from Ramakrishna's and Gee's. For more details, see our preprint [To].

References

- [G] T. Gee, *Companion forms over totally real fields, II*, Duke Math. J. **136** (2007), 275–284.
- [K] C. Khare, *Base Change, Lifting and Serre's Conjecture*, J. Number Theory **63** (1997), 387–395.
- [KW1] C. Khare and J.-P. Wintenberger, *Serre's modularity conjecture (I)*, preprint
- [KW2] C. Khare and J.-P. Wintenberger, *Serre's modularity conjecture (II)*, preprint
- [M] J. Manoharmayum, *Lifting Galois representations of number fields*, preprint
- [R1] R. Ramakrishna, *Lifting Galois representations*, Invent. Math. **138** (1999), 537–562.
- [R2] R. Ramakrishna, *Deforming Galois representations and the conjectures of Serre and Fontaine-Mazur*, Ann. of Math. **156** (2002), 115–154.
- [S] J.-P. Serre, *Sur les représentations modulaires de degré 2 de $\text{Gal}(\bar{\mathbb{Q}}/\mathbb{Q})$* , Duke Math. J. **54** (1987), 179–230.
- [Ta] R. Taylor, *On icosahedral Artin representations, II*, Amer. J. Math. **125** (2003), 549–566.
- [To] Y. Tomiyama, *Lifting Galois representations over arbitrary number fields*, preprint available online at <http://arxiv.org/abs/0809.3076>.