

On uniformly-type Sakaguchi functions

Junichi Nishiwaki and Shigeyoshi Owa

Abstract

Let $\mathcal{A}$ be the class of analytic functions $f(z)$ in the open unit disc $\mathbb{U}$. Furthermore, let $\mathcal{US}_s$ and $\mathcal{US}_s(\alpha, \beta)$ be the subclasses of $\mathcal{A}$ consisting of functions $f(z)$ related to uniformly convex and Sakaguchi functions. The object of the present paper is trying to guess inclusive relations between uniform convexity, $\mathcal{S}_p$ and $\mathcal{US}_s$, and considering coefficient inequalities for $f(z)$ belonging to the class $\mathcal{US}_s(\alpha, \beta)$.

1 Introduction

Let $\mathcal{A}$ be the class of functions $f(z)$ of the form

$$f(z) = z + \sum_{n=2}^{\infty} a_n z^n$$

which are analytic in the open unit disc $\mathbb{U} = \{z \in \mathbb{C} : |z| < 1\}$. A function $f(z) \in \mathcal{A}$ is said to be starlike with respect to symmetrical points in $\mathbb{U}$ if it satisfies

$$\operatorname{Re} \left(\frac{zf'(z)}{f(z) - f(-z)} \right) > 0 \quad (z \in \mathbb{U}).$$

This class is introduced by Sakaguchi [7]. A function $f(z) \in \mathcal{A}$ is said to be in the class of uniformly convex (or starlike) functions denoted by $\mathcal{UCV}$ (or $\mathcal{UST}$) if $f(z)$ is convex (or starlike) in $\mathbb{U}$ and maps every circle or circular arc in $\mathbb{U}$ with center at ζ in $\mathbb{U}$ onto the convex arc (or the starlike arc with respect to $f(\zeta)$). These classes are introduced by Goodman [1] (see also [2]). For the class $\mathcal{UCV}$, it is defined as the one variable characterization by Rønning [5] and [6], that is, a function $f(z) \in \mathcal{A}$ is said to be in the class $\mathcal{UCV}$ if it satisfies

$$\operatorname{Re} \left\{ 1 + \frac{zf''(z)}{f'(z)} \right\} > \left| \frac{zf''(z)}{f'(z)} \right| \quad (z \in \mathbb{U}).$$

It is independently studied by Ma and Minda [3]. Further, a function $f(z) \in \mathcal{A}$ is said to be the corresponding class denoted by $\mathcal{S}_p$ if it satisfies

2010 Mathematics Subject Classification: Primary 30C45

Keywords and Phrases: Analytic function, Sakaguchi function, uniformly starlike, uniformly convex, parabolic starlike.

$$\operatorname{Re} \left\{ \frac{zf'(z)}{f(z)} \right\} > \left| \frac{zf'(z)}{f(z)} - 1 \right| \quad (z \in \mathbb{U}).$$

This class $\mathcal{S}_p$ was introduced by Rønning [5]. We easily know that the relation $f(z) \in \mathcal{UCV}$ if and only if $zf'(z) \in \mathcal{S}_p$. By virtue of these classes, we define the subclass $\mathcal{US}_s(\alpha, \beta)$ of $\mathcal{A}$ consisting of functions $f(z)$ which satisfy

$$\operatorname{Re} \left\{ \frac{zf'(z)}{f(z) - f(-z)} \right\} > \alpha \left| \frac{zf'(z)}{f(z) - f(-z)} - \frac{1}{2} \right| + \beta \quad (z \in \mathbb{U})$$

for some α ($\alpha \geq 0$) and β ($0 \leq \beta < \frac{1}{2}$). We denote $\mathcal{US}_s(1, 0) \equiv \mathcal{US}_s$.

2 Some examples of relation between the class $\mathcal{S}_p$, $\mathcal{UCV}$ and $\mathcal{US}_s$

We don't have the inclusion relation between the class $\mathcal{S}_p$, $\mathcal{UCV}$ and $\mathcal{US}_s$. However, we give two examples to consider some relations between these classes.

Example 2.1. Let us consider the function $f(z) \in \mathcal{A}$ as given by

$$f(z) = z + \frac{1}{5}z^3.$$

Then we obtain

$$\begin{aligned} \operatorname{Re} \left(\frac{zf'(z)}{f(z) - f(-z)} \right) - \left| \frac{zf'(z)}{f(z) - f(-z)} - \frac{1}{2} \right| & \quad (z = re^{i\theta}) \\ &= \frac{25 + 20r^2 \cos 2\theta + 3r^4}{50 + 20r^2 \cos 2\theta + 2r^4} - \frac{r^2}{\sqrt{25 + 10r^2 \cos 2\theta + r^4}} \\ &\geq \frac{(1-r)(5+2r)}{2(5-2r)} > 0 \end{aligned}$$

which shows that $f(z) \in \mathcal{US}_s$. On the other hand, choosing $z = \frac{2}{3}e^{\frac{\pi}{2}i}$, we get

$$\operatorname{Re} \left(1 + \frac{zf''(z)}{f'(z)} \right) - \left| \frac{zf''(z)}{f'(z)} \right| = -\frac{5}{11}$$

which shows that $f(z) \notin \mathcal{UCV}$.

Example 2.2. Let us consider the function $f(z) \in \mathcal{A}$ as given by

$$f(z) = z + \frac{1}{7}z^4.$$

Then we obtain

$$\operatorname{Re} \left(\frac{zf'(z)}{f(z)} \right) - \left| \frac{zf'(z)}{f(z)} - 1 \right| \quad (z = re^{i\theta})$$

$$\begin{aligned}
&= \frac{49 + 35r^3 \cos 2\theta + 4r^6}{49 + 14r^3 \cos 2\theta + r^6} - \frac{3r^3}{\sqrt{49 + 14r^3 \cos 2\theta + r^6}} \\
&\geq \frac{7(1 - r^3)}{7 - r^3} > 0
\end{aligned}$$

which shows that $f(z) \in \mathcal{S}_p$. On the other hand, choosing $z = \frac{23}{24}e^{\frac{\pi}{3}i}$, we get

$$\operatorname{Re} \left(\frac{zf'(z)}{f(z) - f(-z)} \right) - \left| \frac{zf'(z)}{f(z) - f(-z)} - \frac{1}{2} \right| = -\frac{71}{24192}$$

which shows that $f(z) \notin \mathcal{US}_s$.

3 Coefficient inequalities for the class $\mathcal{US}_s(\alpha, \beta)$

Our aim of this section is to discuss some coefficient inequalities for function $f(z)$ to be in the class $\mathcal{US}_s(\alpha, \beta)$.

Theorem 3.1. *If $f(z) \in \mathcal{A}$ satisfies*

$$(3.1) \quad \sum_{n=2}^{\infty} [2(n-1)(1+\alpha)|a_{2n-2}| + \{2n-1-2\beta+2\alpha(n-1)\}|a_{2n-1}|] \leq 1-2\beta$$

for some α ($\alpha \geq 0$) and β ($0 \leq \beta < \frac{1}{2}$), then $f(z) \in \mathcal{US}_s(\alpha, \beta)$.

Let us consider an example for Theorem 3.1.

Example 3.1. *supposing that the function $f(z) \in \mathcal{A}$ as given by*

$$f(z) = z + \sum_{n=2}^{\infty} \frac{(1-2\beta)t\delta_{2n-2}}{2n(n-1)^2(1+\alpha)} z^{2n-2} + \sum_{n=2}^{\infty} \frac{(1-2\beta)(1-t)\delta_{2n-1}}{\{2n-1-2\beta+2\alpha(n-1)\}n(n-1)} z^{2n-1}$$

for some α ($\alpha \geq 0$), β ($0 \leq \beta < \frac{1}{2}$), t ($0 \leq t \leq 1$) and $|\delta_{2n-2}| = |\delta_{2n-1}| = 1$. Then the coefficient (3.1) yields

$$\begin{aligned}
&\sum_{n=2}^{\infty} [2(n-1)(1+\alpha)|a_{2n-2}| + \{2n-1-2\beta+2\alpha(n-1)\}|a_{2n-1}|] \\
&= \sum_{n=2}^{\infty} \left\{ \frac{(1-2\beta)t}{n(n-1)} + \frac{(1-2\beta)(1-t)}{n(n-1)} \right\} \\
&\leq 1-2\beta.
\end{aligned}$$

This implies that $f(z) \in \mathcal{US}_s(\alpha, \beta)$.

Theorem 3.2. *If $f(z) \in \mathcal{US}_s(\alpha, \beta)$, then*

$$|a_2| \leq \frac{1-2\beta}{|1-\alpha|},$$

$$|a_3| \leq \frac{1-2\beta}{|1-\alpha|},$$

$$|a_{2n}| \leq \frac{1-2\beta}{n|1-\alpha|} \prod_{j=1}^{n-1} \left(1 + \frac{1-2\beta}{j|1-\alpha|}\right) \quad (n = 2, 3, 4, \dots)$$

and

$$|a_{2n+1}| \leq \frac{1-2\beta}{n|1-\alpha|} \prod_{j=1}^{n-1} \left(1 + \frac{1-2\beta}{j|1-\alpha|}\right) \quad (n = 2, 3, 4, \dots).$$

References

- [1] A. W. Goodman, *On uniformly convex functions*, Annal. Polon. Math. **56**(1991), 87 – 92.
- [2] A. W. Goodman, *On uniformly starlike functions*, J. Math. Anal. Appl. **155**(1991), 364 – 370.
- [3] W. Ma and D. Minda, *Uniformly convex functions*, Annal. Polon. Math. **57**(1992), 165 – 175.
- [4] S. Owa, T. Sekine, R. Yamakawa, *Notes on Sakaguchi functions*, Austral. J. Math. Appl. **3** (12) (2005), 1 – 7.
- [5] F. Rønning, *Uniformly convex functions and a corresponding class of starlike functions*, Proc. Amer. Math. Soc. **118**(1993), 189 – 196.
- [6] F. Rønning, *On uniform starlikeness and related properties of univalent functions*, Complex Variables **24**(1994), 233 – 239.
- [7] K. Sakaguchi, *On a certain univalent mapping*, J. Math. Soc. Japan **11**(1959), 72 – 75.
- [8] S. Shams, S. R. Kulkarni, and J. M. Jahangiri, *Classes of uniformly starlike and convex functions*, Internat. J. Math. Math. Sci. **55**(2004), 2959 – 2961.

Junichi Nishiwaki
 Department of Mathematics and Physics
 Setsunan University
 Neyagawa, Osaka 572-8508 Japan
 email:jerjun2002@yahoo.co.jp

Shigeyoshi Owa
Department of Mathematics
Faculty of Education
Yamato University Katayama 2-5-1, Suita, Osaka 564-0082
Japan
email: *owa.shigeyoshi@yamato-u.ac.jp*