

On the parity of poly-Euler numbers

By

YASUO OHNO* and YOSHITAKA SASAKI**

Abstract

Poly-Euler numbers are introduced in [9] via special values of an L -function as a generalization of the Euler numbers. In this article, poly-Euler numbers with negative index are mainly treated, and the parity of them is shown as the main theorem. Furthermore the divisibility of poly-Euler numbers are also discussed.

§ 1. Introduction

For every integer k , we define *poly-Euler numbers* $E_n^{(k)}$ ($n = 0, 1, 2, \dots$), which is introduced as a generalization of the Euler number, by

$$(1.1) \quad \frac{\operatorname{Li}_k(1 - e^{-4t})}{4t(\cosh t)} = \sum_{n=0}^{\infty} \frac{E_n^{(k)}}{n!} t^n.$$

Here,

$$\operatorname{Li}_k(x) := \sum_{n=1}^{\infty} \frac{x^n}{n^k} \quad (|x| < 1, \ k \in \mathbb{Z})$$

is the k -th polylogarithm. When $k = 1$, $E_n^{(1)}$ is the Euler number defined by

$$(1.2) \quad \frac{1}{\cosh t} = \sum_{n=0}^{\infty} \frac{E_n^{(1)}}{n!} t^n.$$

Received March 31, 2011. Revised December 8, 2011.

2000 Mathematics Subject Classification(s): 11B68, 11M41.

Key Words: Poly-Euler number, poly-Bernoulli number

*Department of Mathematics, Kinki University,

Kowakae 3-4-1, Higashi-Osaka, Osaka 577-8502, Japan.

e-mail: ohno@math.kindai.ac.jp

**Interdisciplinary Graduate School of Science and Engineering, Kinki University,

Kowakae 3-4-1, Higashi-Osaka, Osaka 577-8502 Japan.

e-mail: sasaki@alice.math.kindai.ac.jp

© 2012 Research Institute for Mathematical Sciences, Kyoto University. All rights reserved.

The reason why we refer to $E_n^{(k)}$'s as “poly-Euler numbers” will be mentioned in the next section from the point of view of the relation between the poly-Bernoulli number and Arakawa-Kaneko's zeta-function. In this article, we treat some number theoretical properties of poly-Euler numbers with negative index ($k \leq 0$). Our main theorem is Theorem 3.1 described in Section 3, which mentions the parity of poly-Euler numbers can be determined definitely. In Section 4, we discuss the divisibility of poly-Euler numbers via congruence relations of them. Tables 1 and 2 cited at the end of this article are the lists of numerical values of poly-Euler numbers. General properties of poly-Euler numbers including the case of positive index are treated in [8].

§ 2. The poly-Bernoulli numbers and Arakawa-Kaneko's zeta-function

For every integer k , the poly-Bernoulli numbers $\mathbb{B}_n^{(k)}$ and the modified poly-Bernoulli numbers $C_n^{(k)}$ introduced by Kaneko [4] are defined by

$$(2.1) \quad \frac{\text{Li}_k(1 - e^{-t})}{1 - e^{-t}} = \sum_{n=0}^{\infty} \frac{\mathbb{B}_n^{(k)}}{n!} t^n \quad \text{and} \quad \frac{\text{Li}_k(1 - e^{-t})}{e^t - 1} = \sum_{n=0}^{\infty} \frac{C_n^{(k)}}{n!} t^n,$$

respectively. When $k = 1$, the above generating functions become

$$\frac{te^t}{e^t - 1} \quad \text{and} \quad \frac{t}{e^t - 1},$$

respectively. Therefore $\mathbb{B}_n^{(k)}$ and $C_n^{(k)}$ are generalizations of the classical Bernoulli numbers. Some number theoretic properties of the poly-Bernoulli number were given by Kaneko [4], Arakawa and Kaneko [2] and others. Furthermore the combinatorial interpretations of $\mathbb{B}_n^{(-k)}$ were given by Brewbaker [3] and Launois [6]. Recently, Shikata [10] gives the alternative proof of the result of Brewbaker.

It is known that the poly-Bernoulli numbers are special values of Arakawa-Kaneko's zeta-function. Arakawa and Kaneko [1] introduced a zeta-function:

$$\xi_k(s) := \frac{1}{\Gamma(s)} \int_0^\infty t^{s-1} \frac{\text{Li}_k(1 - e^{-t})}{e^t - 1} dt \quad (k \geq 1).$$

We refer to the above function as *Arakawa-Kaneko's zeta-function*. Arakawa-Kaneko's zeta-function satisfies $\xi_1(s) = s\zeta(s+1)$, and

$$\xi_k(-n) = \sum_{l=0}^n (-1)^l \binom{n}{l} \mathbb{B}_l^{(k)} = (-1)^n C_n^{(k)}$$

for any non-positive integer n , where $\zeta(s)$ is the Riemann zeta-function. Hence Arakawa-Kaneko's zeta-function is a kind of generalization of the Riemann zeta-function. Furthermore, we should mention that Arakawa-Kaneko's zeta-function is applied to research

on multiple zeta values. For example, Kaneko and Ohno [5] showed a duality property of multiple zeta-star values by using this property.

From the point of view mentioned above, poly-Euler numbers should be defined as special values of an L -function generalized by using the method of Arakawa and Kaneko. Moreover it can be reasonably expected that such L -function has nice properties and applications similar to Arakawa-Kaneko's zeta-function.

The Euler number E_n is the generalized Bernoulli number associated with the Dirichlet character of conductor 4, and the L -function is

$$(2.2) \quad L(s) := \frac{1}{\Gamma(s)} \int_0^\infty t^{s-1} \frac{1}{e^t + e^{-t}} dt$$

satisfies $L(-n) = E_n/2$ for any non-negative integer n . The second author gave in [9] a general method for defining L -functions that have similar properties to Arakawa-Kaneko's zeta-function. By using the method, a generalization of $L(s)$ is given by

$$(2.3) \quad L_k(s) = \frac{1}{\Gamma(s)} \int_0^\infty t^{s-1} \frac{\text{Li}_k(1 - e^{-4t})}{4(e^t + e^{-t})} dt \quad (k \geq 1).$$

It is natural to define poly-Euler numbers as special values at non-positive integers of the above L -function. Therefore, we define poly-Euler numbers as above (1.1).

We should mention that the above generalized L -function is applicable to research on multiple L values. In fact, the second author [9] treated the generalization of Dirichlet L -functions for general Dirichlet characters and showed such generalized L -functions can be written in terms of multiple L -functions.

Remark. Although $L_1(s)$ does not reduce to $L(s)$ ($L_1(s) = sL(s+1)$), the above definition (2.3) is optimal as an analogue of Arakawa-Kaneko's zeta-function. In fact, we have $\xi_1(s) = s\zeta(s+1)$.

§ 3. The parity of poly-Euler numbers

In this section, we determine the parity of poly-Euler numbers $(n+1)E_n^{(-k)}$.

Theorem 3.1. *For any non-negative integer k , $(n+1)E_n^{(-k)}$ is even (odd, respectively) integer, when n is odd (even, respectively).*

Proof of Theorem 3.1. We first review an explicit formula:

Lemma 3.2 (Ohno-Sasaki [8]). *For any non-negative integers k and n , we have*

$$(3.1) \quad (n+1)E_n^{(-k)} = (-1)^k \sum_{l=0}^k (-1)^l l! \left\{ \begin{matrix} k \\ l \end{matrix} \right\} \sum_{\substack{m=1, \\ m:\text{odd}}}^{n+1} \binom{n+1}{m} (4l+2)^{n+1-m},$$

where the symbol $\left\{ \begin{smallmatrix} k \\ l \end{smallmatrix} \right\}$ is the Stirling number of the second kind defined by the recurrence relation

$$(3.2) \quad \left\{ \begin{smallmatrix} k+1 \\ l \end{smallmatrix} \right\} = \left\{ \begin{smallmatrix} k \\ l-1 \end{smallmatrix} \right\} + l \left\{ \begin{smallmatrix} k \\ l \end{smallmatrix} \right\}$$

with

$$\left\{ \begin{smallmatrix} 0 \\ 0 \end{smallmatrix} \right\} = 1, \quad \left\{ \begin{smallmatrix} k \\ 0 \end{smallmatrix} \right\} = \left\{ \begin{smallmatrix} 0 \\ l \end{smallmatrix} \right\} = 0 \quad (k, l \neq 0)$$

for any integers k and l .

See [8] for the detailed proof of Lemma 3.2.

Remark. We easily see that the right-hand side of (3.1) is an integer, since the Stirling numbers are integers. Furthermore this fact indicates that the denominator of poly-Euler number $E_n^{(-k)}$ is at most $n+1$.

We prove Theorem 3.1 by using the above lemma. Note that

$$\sum_{\substack{m=1, \\ m:\text{odd}}}^{n+1} \binom{n+1}{m} (4l+2)^{n+1-m} \equiv \begin{cases} 1 & (\text{mod } 2) \quad n : \text{even}, \\ 0 & (\text{mod } 2) \quad n : \text{odd} \end{cases}$$

for any non-negative integer l . Hence, when n is odd, we have Theorem 3.1 immediately from Lemma 3.2.

On the other hand, when n is even, we have

$$(n+1)E_n^{(-k)} \equiv \sum_{l=0}^k l! \left\{ \begin{smallmatrix} k \\ l \end{smallmatrix} \right\} \equiv \left\{ \begin{smallmatrix} k \\ 0 \end{smallmatrix} \right\} + \left\{ \begin{smallmatrix} k \\ 1 \end{smallmatrix} \right\} = \left\{ \begin{smallmatrix} k+1 \\ 1 \end{smallmatrix} \right\} = 1 \pmod{2}$$

for any non-negative integer k . Here, we have used the recurrence relation (3.2). Thus the proof of Theorem 3.1 is completed. $\square$

§ 4. Congruence relations of poly-Euler numbers

In the previous section, we definitely determined the parity of poly-Euler numbers. In this section, we treat the divisibility of poly-Euler numbers via congruence relations of them. In particular, we consider the case of

$$(n+1)E_n^{(-k)} \pmod{n+1},$$

which allows us to evaluate whether $E_n^{(-k)}$ is an integer. In [8], we treat the case when $n+1$ is an odd prime. Hence we discuss the composite cases here.

Theorem 4.1. *For any non-negative integer k , we have*

$$6E_5^{(-k)} \equiv \begin{cases} 4 \pmod{6} & \text{if } k \text{ is even,} \\ 0 \pmod{6} & \text{if } k \text{ is odd.} \end{cases}$$

Proof of Theorem 4.1. From Lemma 3.2, we have

$$(4.1) \quad 6E_5^{(-k)} = (-1)^k \sum_{l=0}^k (-1)^l l! \left\{ \begin{matrix} k \\ l \end{matrix} \right\} \sum_{j=0}^2 \binom{6}{2j+1} (4l+2)^{5-2j}.$$

We see that $l! \equiv 0 \pmod{6}$ for $l \geq 3$ and

$$\begin{aligned} \sum_{j=0}^2 \binom{6}{2j+1} (4l+2)^{5-2j} &\equiv 2(4l+2)^3 \pmod{6} \\ &\equiv \begin{cases} 4 \pmod{6} & \text{if } l \equiv 0 \pmod{3}, \\ 0 \pmod{6} & \text{if } l \equiv 1 \pmod{3}, \\ 2 \pmod{6} & \text{if } l \equiv 2 \pmod{3}. \end{cases} \end{aligned}$$

Thus (4.1) becomes

$$\begin{aligned} 6E_5^{(-k)} &\equiv (-1)^k 4 \left(\left\{ \begin{matrix} k \\ 0 \end{matrix} \right\} + \left\{ \begin{matrix} k \\ 2 \end{matrix} \right\} \right) \pmod{6} \\ &\equiv \begin{cases} 4 \pmod{6} & \text{if } k = 0, \\ 0 \pmod{6} & \text{if } k = 1, \\ (-1)^k \left\{ \begin{matrix} k \\ 2 \end{matrix} \right\} 4 \pmod{6} & \text{if } k \geq 2. \end{cases} \end{aligned}$$

Therefore, we claim

$$(4.2) \quad (-1)^k \left\{ \begin{matrix} k \\ 2 \end{matrix} \right\} \equiv \begin{cases} 1 \pmod{6} & \text{if } k \text{ is even,} \\ 3 \pmod{6} & \text{if } k \text{ is odd.} \end{cases}$$

Using the expression $\left\{ \begin{matrix} k \\ 2 \end{matrix} \right\} = 2^{k-1} - 1$ ($k \geq 2$), we have

$$\begin{aligned} \left\{ \begin{matrix} k \\ 2 \end{matrix} \right\} &= 2^{k-1} - 1 = \sum_{j=0}^{k-2} 2^j = 1 + \sum_{\substack{j=1, \\ j:\text{odd}}}^{k-2} 2^j + \sum_{\substack{j=2, \\ j:\text{even}}}^{k-2} 2^j \\ &\equiv 1 + 2[(k-1)/2] + 4[(k-2)/2] \pmod{6}, \end{aligned}$$

which gives (4.2). Thus we have Theorem 4.1. □

By the same way as above, we can also show the following theorem:

Theorem 4.2. *For any non-negative integer k , we have*

$$12E_{11}^{(-k)} \equiv \begin{cases} 4 \pmod{12} & \text{if } k \text{ is even,} \\ 0 \pmod{12} & \text{if } k \text{ is odd.} \end{cases}$$

In general, to understand the prime factors of the numerator of $E_n^{(-k)}$ is proper. In [8], we prove a congruence relation

$$(n+1)E_n^{(-k)} \equiv 0 \pmod{p}$$

holds for any odd prime p , odd positive integer n and non-positive integer k satisfying $k \equiv p-2 \pmod{p-1}$. The second assertion of Theorem 4.1 is also given by combining the above congruence relation with Theorem 3.1.

Acknowledgements

The authors would like to express their sincere gratitude to Professors Masanari Kida, Noriyuki Suwa and Shinichi Kobayashi for giving us the opportunity to talk in the conference “Algebraic Number Theory and Related Topics 2010”. Thanks are also due to the referee for his/her helpful suggestion. The first author is supported by Grant-in-Aid for Scientific Research (C) No. 20540033. The second author is supported by Grants-in-Aid for Research Activity Start-up No. 21840053.

References

- [1] T. Arakawa and M. Kaneko, *Multiple zeta values, poly-Bernoulli numbers, and related zeta functions*, Nagoya Math. J. **153** (1999), 189–209.
- [2] T. Arakawa and M. Kaneko, *On poly-Bernoulli numbers*, Comment. Math. Univ. St. Paul. **48** (1999), 159–167.
- [3] C. Brewbaker, *Lonesum $(0, 1)$ -matrices and poly-Bernoulli numbers of negative index*, Master’s thesis, Iowa State University, 2005.
- [4] M. Kaneko, *Poly-Bernoulli numbers*, J. Th. Nombre Bordeaux, **9** (1997), 199–206.
- [5] M. Kaneko and Y. Ohno, *On a kind of duality of multiple zeta-star values*, Int. J. Number Theory, **6** (2010), 1927–1932.
- [6] S. Launois, *Rank t $\mathcal{H}$ -primes in quantum matrices*, Comm. Algebra **33** (2005), 837–854.
- [7] Y. Ohno, *A generalization of the duality and sum formulas on the multiple zeta values*, J. Number Theory **74** (1999), 39–43.
- [8] Y. Ohno and Y. Sasaki, *On poly-Euler numbers*, preprint.
- [9] Y. Sasaki, *On generalized poly-Bernoulli numbers and related L -functions*, J. Number Theory **132** (2012), 156–170.
- [10] M. Shikata, *Lonesum matrices and poly-Bernoulli numbers*, Kyoto Sangyo University essays. Natural science series, **40** (2011), 1–12.

Table 1. $E_n^{(k)}$ (positive index)

$n \backslash k$	1	2	3	4
0	1	1	1	1
1	0	-1	$-\frac{3}{2}$	$-\frac{7}{4}$
2	-1	$-\frac{1}{9}$	$\frac{41}{27}$	$\frac{221}{81}$
3	0	3	$\frac{7}{6}$	$-\frac{85}{36}$
4	5	$-\frac{51}{25}$	$-\frac{3493}{375}$	$-\frac{31079}{5625}$
5	0	-25	$\frac{521}{90}$	$\frac{77071}{2700}$
6	-61	$\frac{33221}{735}$	$\frac{8169601}{77175}$	$-\frac{41535229}{8103375}$
7	0	427	$-\frac{18313}{70}$	$-\frac{19160833}{44100}$
8	1385	$-\frac{1288391}{945}$	$-\frac{70339397}{33075}$	$\frac{33076559267}{31255875}$
9	0	-12465	$\frac{11557561}{1050}$	$\frac{748381847}{73500}$
10	-50521	$\frac{252042789}{4235}$	$\frac{986047910537}{14674275}$	$-\frac{3065162516767009}{50846362875}$
11	0	555731	$-\frac{4213240267}{6930}$	$-\frac{5414464148791}{16008300}$
12	2702765	$-\frac{706698126353}{195195}$	$-\frac{2983263446051093}{976950975}$	$\frac{530921430493573134689}{132020270006625}$
13	0	-35135945	$\frac{624220472627}{14014}$	$\frac{89775517476597787}{6312606300}$
14	-199360981	$\frac{66380136099899}{225225}$	$\frac{631679911607350337}{3381753375}$	$-\frac{1910856585406763406371}{5641891880625}$

Table 2. $E_n^{(-k)}$ (negative index)

$n \backslash -k$	0	-1	-2	-3
0	1	1	1	1
1	2	6	14	30
2	$\frac{13}{3}$	$\frac{109}{3}$	$\frac{493}{3}$	$\frac{1837}{3}$
3	10	222	1798	10710
4	$\frac{121}{5}$	$\frac{6841}{5}$	$\frac{95161}{5}$	$\frac{865081}{5}$
5	$\frac{182}{3}$	8502	$\frac{594554}{3}$	2670350
6	$\frac{1093}{7}$	$\frac{372709}{7}$	$\frac{14331493}{7}$	$\frac{280592677}{7}$
7	410	335886	21078134	591278790
8	$\frac{9841}{9}$	$\frac{19200241}{9}$	$\frac{1951326961}{9}$	$\frac{77624198641}{9}$
9	$\frac{14762}{5}$	$\frac{68177406}{5}$	$\frac{11157142694}{5}$	124916013054
10	$\frac{88573}{11}$	$\frac{964249309}{11}$	$\frac{252966361693}{11}$	$\frac{19811958812317}{11}$
11	$\frac{66430}{3}$	566547774	$\frac{712300066738}{3}$	25898630029750
12	$\frac{797161}{13}$	$\frac{47834153641}{13}$	$\frac{31933012161961}{13}$	$\frac{4834065180508201}{13}$
13	$\frac{1195742}{7}$	$\frac{168029889306}{7}$	25466927239262	$\frac{37348182927733890}{7}$
14	$\frac{7174453}{15}$	$\frac{2358521965909}{15}$	$\frac{3968998515355093}{15}$	$\frac{1148212279510308757}{15}$