

Singular domains in higher dimensional complex dynamics

Yûsuke Okuyama

Department of Comprehensive Sciences,
Kyoto Institute of Technology, Kyoto 606-8585 JAPAN
email; okuyama@kit.ac.jp

This article aims to extend the fundamental Cremer theorem from the iteration theory of one complex variable to the setting of higher-dimensional dynamics over more general valued fields, not necessarily $\mathbb{C}$. This article is an announcement of the preprint [Oku2].

Projective spaces over valued fields. Let K be a commutative algebraically closed field which is complete and nondiscrete with respect to a non-trivial absolute value (or valuation) $|\cdot|$. This $|\cdot|$ is said to be *non-Archimedean* if $\forall z, \forall w \in K, |z - w| \leq \max\{|z|, |w|\}$. Otherwise, $|\cdot|$ is said to be *Archimedean* and K is then topologically isomorphic to $\mathbb{C}$ (with Hermitian norm). We extend $|\cdot|$ to K^ℓ ($\ell \in \mathbb{N}$) as the maximum norm $|Z| = |Z|_\ell = \max_{j=1, \dots, \ell} |z_j|$ for $Z = (z_1, \dots, z_\ell)$. Let $\pi : K^{n+1} \setminus \{O\} \rightarrow \mathbb{P}^n(K)$ be the canonical projection and set $\ell(n) \in \mathbb{N}$ so that $\bigwedge^2 K^{n+1} \cong K^{\ell(n)}$. The *chordal distance* $[\cdot, \cdot]$ on $\mathbb{P}^n(K)$ is defined as

$$[z, w] := \frac{|Z \wedge W|_{\ell(n)}}{|Z|_{n+1} |W|_{n+1}},$$

where $Z \in \pi^{-1}(z), W \in \pi^{-1}(w)$ (cf. [KS]). For $z_0 \in \mathbb{P}^n(K)$ and $r > 0$, we consider the ball

$$\overline{B}(z_0, r) := \{z \in \mathbb{P}^n(K); [z, z_0] \leq r\}.$$

Nonlinearity of morphisms. Let $f : \mathbb{P}^n(K) \rightarrow \mathbb{P}^n(K)$ be a (finite) *morphism*, i.e., there is a homogeneous polynomial map $F : K^{n+1} \rightarrow K^{n+1}$ over K , which is called a *lift* of f , such that $F^{-1}(O) = \{O\}$ and satisfies

$$\pi \circ F = f \circ \pi.$$

The degree $d = \deg f$ is that of F as homogeneous polynomial map. As in the case of $K = \mathbb{C}$, the *Fatou set* $F(f)$ is the largest open set at each point of which the family $\{f^k; k \in \mathbb{N}\}$ is equicontinuous.

The *Julia set* $J(f)$ is defined by $\mathbb{P}^n(K) \setminus F(f)$. In non-Archimedean case, $J(f)$ may be empty even if $d \geq 2$. One of the main results is

Theorem 1 (nonlinearity of morphisms). *Let $f : \mathbb{P}^n(K) \rightarrow \mathbb{P}^n(K)$ be a morphism of degree $d \geq 1$. If there are a ball $\overline{B}(z_0, r) \subset \mathbb{P}^n(K)$ and a morphism $g : \mathbb{P}^n(K) \rightarrow \mathbb{P}^n(K)$ such that*

$$\liminf_{k \rightarrow \infty} \frac{1}{d^k} \log \sup_{\overline{B}(z_0, r)} [f^k, g] = -\infty,$$

then either f is linear or $J(f) = \emptyset$.

We give a few applications of Theorem 1.

Analytic linearization over a field K . Consider the K -algebra

$$\mathcal{O}_\ell \cong K\{X_1, \dots, X_\ell\} = \{f = \sum c_I X^I; \limsup_{|I| \rightarrow \infty} |c_I|^{1/|I|} =: r_f^{-1} < \infty\}$$

of all germs of analytic functions at the origin $O \in K^\ell$. Here $I = (i_1, \dots, i_\ell) \in \mathbb{Z}_{\geq 0}^\ell$ is a multi-index, $X_1^{i_1} \cdots X_\ell^{i_\ell}$ is denoted by X^I and we put $|I| := i_1 + \dots + i_\ell$. For germ of analytic map $\phi = (f_1, \dots, f_n) \in (\mathcal{O}_n)^n$, we identify the linear part of $\phi - \phi(O)$ at O with

$$A_\phi := \left(\frac{\partial f_i}{\partial X_j}(O) \right)_{i,j=1,\dots,n} \in M(n, K) \cong \text{End}(K^n).$$

We also denote the operator norm on $M(n, K)$ by $|\cdot|$.

A germ $\phi = (f_1, \dots, f_n) \in (\mathcal{O}_n)^n$ fixing O is (analytically) *linearizable* if there is $H \in (\mathcal{O}_n)^n$ fixing O such that $A_H = I_n$ (unit matrix) and H satisfies the *Schröder* (or *Poincaré*) equation

$$\phi \circ H = H \circ A_\phi.$$

From Siegel and Sternberg ([Sie], [Ste]) and its non-Archimedean version by Herman-Yoccoz [HY], ϕ is linearizable if A_ϕ is diagonalizable and its eigenvalues $\lambda_1, \dots, \lambda_n$ satisfy the *Diophantine* condition: there exist $C > 0$ and $\beta \geq 0$ such that for every $I \in \mathbb{Z}_{\geq 0}^n$ (multi-index) with $|I| \geq 1$,

$$|(\lambda_1, \dots, \lambda_n)^I - 1| \geq \frac{C}{|I|^\beta}.$$

On the other hand, consider an inverse of a coordinate chart

$$\sigma : K^n \ni (z_1, \dots, z_n) \mapsto (1 : z_1 : \dots : z_n) \in \mathbb{P}^n(K).$$

When a morphism $f : \mathbb{P}^n(K) \rightarrow \mathbb{P}^n(K)$ fixes a point $z_0 \in \mathbb{P}^n(K)$, assuming that $z_0 = \sigma(O)$ without loss of generality, we say f to be *linearizable* at z_0 if the germ $\phi_f \in (\mathcal{O}_n)^n$ of the analytic map $\sigma^{-1} \circ f \circ \sigma : \overline{P}^n(O, r) \rightarrow K^n$ is linearizable. The following is regarded as a higher dimensional version of the Cremer condition [Cre, p. 157].

Theorem 2 (nonresonance). *Let $f : \mathbb{P}^n(K) \rightarrow \mathbb{P}^n(K)$ be a morphism of degree $d \geq 2$ which fixes $z_0 \in \mathbb{P}^n(K)$, and suppose that $J(f) \neq \emptyset$. If f is linearizable at z_0 and $|A_{\phi_f}| \leq 1$, then*

$$\liminf_{k \rightarrow \infty} \frac{1}{d^k} \log |(A_{\phi_f})^k - I_n| > -\infty.$$

If in addition A_{ϕ_f} is diagonalizable, then its eigenvalues $\lambda_1, \dots, \lambda_n$ satisfy

$$\liminf_{k \rightarrow \infty} \frac{1}{d^k} \log \max_{j=1, \dots, n} |\lambda_j^k - 1| > -\infty.$$

Singular domain over the field $\mathbb{C}$. Let $f : \mathbb{P}^n = \mathbb{P}^n(\mathbb{C}) \rightarrow \mathbb{P}^n$ be a morphism, which is now holomorphic, of degree $d \geq 2$.

Each component D of $F(f)$, which is called a *Fatou component* of f , is Stein and Kobayashi hyperbolic [Ued1]. In particular, D is holomorphically separable and the biholomorphic automorphisms $\text{Aut}(D)$ is a Lie group. When there is a sequence $(f^{k_j}) \subset \{f^k\}$ which converges to Id_D locally uniformly on D , we have $f^p(D) = D$ for some $p \in \mathbb{N}$ and moreover $f^p|_D \in \text{Aut}(D)$. Following Fatou [Fat, §28], we call such D a *singular domain* (un domaine *singulier*) of f . A singular domain is also called a *Siegel domain* or *rotation domain*. When $n = 1$, a singular domain D is either a Siegel disk or an Herman ring. When $n \geq 2$, a partial analogue is known: let G be the closed subgroup generated by $f^p|_D$ in $\text{Aut}(D)$, and G_0 the component of G containing Id_D . Then there is a Lie group isomorphism $G_0 \rightarrow \mathbb{T}^s$ for some $s \in [1, n]$, which maps $f^q|_D$ for some $q \in \mathbb{N}$ to $(e^{2i\pi\alpha_1}, \dots, e^{2i\pi\alpha_s})$ for some $\alpha_1, \dots, \alpha_s \in \mathbb{R} \setminus \mathbb{Q}$ (see [FS1], [Ued2], [Mih]). In the maximal case of $s = n$, we say the singular domain D to be of *maximal type*.

A singular domain D of maximal type is exactly a generalization of one-dimensional Siegel disks and Herman rings: setting $\lambda_j := e^{2i\pi\alpha_j}$ ($j = 1, \dots, n$), we have by [BBD, Theorem 1] a biholomorphic homeomorphism Φ from a Reinhardt domain $U \subset \mathbb{C}^n$ to D such that the Schröder equation

$$f^q(\Phi(w_1, \dots, w_n)) = \Phi(\lambda_1 w_1, \dots, \lambda_n w_n) \quad \text{on } U$$

holds.

Theorem 3 (a priori bound). *Let $f : \mathbb{P}^n \rightarrow \mathbb{P}^n$ be a holomorphic map of degree $d \geq 2$. If a singular domain D of f is of maximal type, then under the same notation as in the above, D satisfies*

$$\lim_{k \rightarrow \infty} \frac{1}{d^{qk}} \log \max_{j=1, \dots, n} |\lambda_j^k - 1| = 0.$$

In the case of $n = 1$, every singular domain of f is of maximal type. In this case, Theorem 3 is essentially proved in [FS2, p. 169] by pluripotential theory, and in [Oku1, Main Theorem 3] by a Nevanlinna theoretical argument. Both proofs contain some one-dimensional arguments which are not easily extended to higher dimensions. Our proof of Theorem 3 is based on a proof of Theorem 1, which dispenses with pluripotential theory.

Finally, we give a *vanishing* result on the Valiron deficiency

$$\delta_V(\text{Id}_{\mathbb{P}^n}, (f^k)) := \limsup_{k \rightarrow \infty} \frac{1}{d^k} \int_{\mathbb{P}^n} \log \frac{1}{[f^k, \text{Id}]} d\omega_{FS}^n$$

(cf. [DO]). Here ω_{FS} denotes the Fubini-Study Kähler form on $\mathbb{P}^n$.

Theorem 4 (a vanishing theorem). *Let $f : \mathbb{P}^n \rightarrow \mathbb{P}^n$ be a holomorphic map of degree ≥ 2 . If every singular domain of f is of maximal type, then*

$$\delta_V(\text{Id}_{\mathbb{P}^n}, (f^k)) = 0.$$

We expect that the assertion of Theorem 4 still remains true with no maximality assumption on singular domains.

References

- [BBD] David E. Barrett, Eric Bedford, and Jiri Dadok. T^n -actions on holomorphically separable complex manifolds. *Math. Z.* **202**(1989), 65–82.
- [Cre] Hubert Cremer. Zum Zentrumproblem. *Math. Ann.* **98**(1928), 151–163.
- [DO] David Drasin and Yûsuke Okuyama. Equidistribution and Nevanlinna theory. *Bull. Lond. Math. Soc.* **39**(2007), 603–613.
- [Fat] Pierre Fatou. Sur les équations fonctionnelles. *Bull. Soc. Math. France* **48**(1920), 33–94.

- [FS1] John Erik Fornæss and Nessim Sibony. Classification of recurrent domains for some holomorphic maps. *Math. Ann.* **301**(1995), 813–820.
- [FS2] John Erik Fornæss and Nessim Sibony. Complex dynamics in higher dimension. II. In *Modern methods in complex analysis (Princeton, NJ, 1992)*, volume 137 of *Ann. of Math. Stud.*, pages 135–182. Princeton Univ. Press, Princeton, NJ, 1995.
- [HY] M. Herman and J.-C. Yoccoz. Generalizations of some theorems of small divisors to non-Archimedean fields. In *Geometric dynamics (Rio de Janeiro, 1981)*, volume 1007 of *Lecture Notes in Math.*, pages 408–447. Springer, Berlin, 1983.
- [KS] Shu Kawaguchi and Joseph H. Silverman. Nonarchimedean Green functions and dynamics on projective space. *Math. Z.* **262**(2009), 173–197.
- [Mih] Eugen Mihailescu. Periodic points for actions of tori in Stein manifolds. *Math. Ann.* **314**(1999), 39–52.
- [Oku1] Yûsuke Okuyama. Nevanlinna, Siegel, and Cremer. *Indiana Univ. Math. J.* **53**(2004), 755–763.
- [Oku2] Yûsuke Okuyama. Nonlinearity of morphisms in non-Archimedean and complex dynamics. *preprint*.
- [Sie] Carl Ludwig Siegel. Iteration of analytic functions. *Ann. of Math. (2)* **43**(1942), 607–612.
- [Ste] Shlomo Sternberg. Infinite Lie groups and the formal aspects of dynamical systems. *J. Math. Mech.* **10**(1961), 451–474.
- [Ued1] Tetsuo Ueda. Fatou sets in complex dynamics on projective spaces. *J. Math. Soc. Japan* **46**(1994), 545–555.
- [Ued2] Tetsuo Ueda. Critical orbits of holomorphic maps on projective spaces. *J. Geom. Anal.* **8**(1998), 319–334.