

Game theoretic properties of the club filter on $\mathcal{P}_\kappa\lambda$

MASAHIRO SHIOYA

塩谷真弘

Let κ be a regular cardinal $> \omega$, λ a cardinal $\geq \kappa$ and F a filter on $\mathcal{P}_\kappa\lambda$. Jech and Prikry [JP] called F precipitous when the poset F^+ ordered by inclusion forces that the generic ultrapower is well-founded. Refining an ingenious idea of Foreman, Magidor and Shelah [FMS], Goldring [G] and Shelah independently established that the club filter $\mathcal{C}_{\kappa\lambda}$ on $\mathcal{P}_\kappa\lambda$ can be made precipitous by Levy collapsing a Woodin cardinal to the successor of λ when λ is regular. In [MS] we showed that $\mathcal{C}_{\kappa\lambda}$ cannot be precipitous e.g. when λ is strong limit and of cofinality $< \kappa$. On the other hand Galvin, Jech and Magidor [GJM] observed that precipitousness is also definable in terms of games (see below). This led us to investigate game theoretic properties of $\mathcal{C}_{\kappa\lambda}$ in this paper.

We refer to Kanamori [K] for basic material. We use μ to denote a regular cardinal. By S_κ^ω and $\lim X$ for $X \subset \kappa$ we denote the set $\{\alpha < \kappa : \text{cf } \alpha = \omega\}$ and $\{\alpha < \kappa : \sup(X \cap \alpha) = \alpha > 0\}$ respectively. For $\alpha < \lambda$ we fix a bijection $\pi_\alpha : |\alpha| \rightarrow \alpha$. Suppose that two players I and II take in turn $S_n \in F^+$ and $T_n \in F^+$ respectively so that $S_n \supset T_n \supset S_{n+1}$. We define the game $\mathcal{G}_0(F)$ by: I wins iff $\bigcap_{n < \omega} S_n = \emptyset$. Recall from [GJM] that F is precipitous iff I has no winning

1991 *Mathematics Subject Classification*. 03E05, 03E55.

Part of this work was done during the author's stay at Boston University as one of the Japanese Overseas Research Fellows. He gratefully acknowledges Prof. Kanamori's hospitality. He also thanks Mr. Yoshinobu for pointing out Proposition 3.

strategy in $\mathcal{G}_0(F)$. Also the game $\mathcal{G}_1(F)$ (resp. $\mathcal{G}_*(F)$) is defined by: I wins iff $|\bigcap_{n<\omega} S_n| \leq 1$ (resp. $\bigcap_{n<\omega} S_n \in F^*$). As observed by Jech [J], F^+ is ω_1 -Baire iff I has no winning strategy in $\mathcal{G}_*(F)$. A winning strategy for I in $\mathcal{G}_1(\mathcal{C}_\kappa)$ from [GJM] yields the following

Proposition 1. $\mathcal{C}_{\kappa\lambda}^+$ is not ω_1 -Baire.

Proof. We give a winning strategy for I in $\mathcal{G}_*(\mathcal{C}_{\kappa\lambda})$. We let $S_0 = \{x \in \mathcal{P}_{\kappa\lambda} : x \cap \kappa \in S_\kappa^\omega\}$ be the first move of I, and for $x \in S_0$ fix an unbounded set $\{\alpha_n^x : n < \omega\} \subset x \cap \kappa$. Next suppose that T is an n -th move of II. Then by the regressive function lemma we have $\gamma_n < \kappa$ such that $\{x \in T : \alpha_n^x = \gamma_n\}$ is stationary, which we let be the next move of I.

Let $\langle S_n : n < \omega \rangle$ be a play of I following the above strategy with $\langle \gamma_n : n < \omega \rangle$ as above. Then $\bigcap_{n<\omega} S_n \subset \{x \in \mathcal{P}_{\kappa\lambda} : x \cap \kappa = \sup_{n<\omega} \gamma_n\} \in \mathcal{C}_{\kappa\lambda}^*$, as desired. $\square$

The following lemma is proved similarly to the original version in [GJM].

Lemma 1. Let $S \subset \{x \in \mathcal{P}_{\kappa\lambda} : x \cap \kappa \in S_\kappa^\omega\}$ be stationary and $\{\alpha_n^x : n < \omega\}$ unbounded in $x \cap \kappa$ for $x \in S$. Then for some $m < \omega$ $\{\gamma < \kappa : \{x \in S : \alpha_m^x = \gamma\} \text{ is stationary}\}$ is unbounded.

Now we have a $\mathcal{P}_{\kappa\lambda}$ generalization of the Galvin-Jech-Magidor result [GJM], which shows that Goldring's result is sharp from a game theoretic point of view:

Proposition 2. II has no winning strategy in $\mathcal{G}_0(\mathcal{C}_{\kappa\lambda})$.

Proof. Let τ be a strategy for II. We construct a play in which II follows τ and I wins. Build inductively $\langle S_t : t \in \mathcal{T} \rangle$ with $\mathcal{T}$ a subtree of $\kappa^{<\omega}$ such that for any $t \in \mathcal{T}$ $\text{succ}_{\mathcal{T}}(t) = \{\gamma < \kappa : t * \langle \gamma \rangle \in \mathcal{T}\}$ is unbounded and $\langle S_{t|i} : i < |t| \rangle$ is a partial

play of I with which II follows τ as follows: First set $S_\emptyset = \{x \in \mathcal{P}_\kappa \lambda : x \cap \kappa \in S\}$, where $S \subset S_\kappa^\omega$ and $S_\kappa^\omega - S$ are both stationary, and for $x \in S_\emptyset$ fix an unbounded set $\{\alpha_n^x : n < \omega\} \subset x \cap \kappa$. Next suppose that $\langle S_t : t \in \mathcal{T} \cap \kappa^n \rangle$ is defined. Fix $t \in \mathcal{T} \cap \kappa^n$. Let T be the n -th move of II following τ when I plays $\langle S_{t|_i} : i < n \rangle$. Then by the above lemma take $m < \omega$ such that $\{\gamma < \kappa : \{x \in T : \alpha_m^x = \gamma\} \text{ is stationary}\}$ is unbounded, which we let be $\text{suc}_T(t)$. For $\gamma \in \text{suc}_T(t)$ set $S_{t* \langle \gamma \rangle} = \{x \in T : \alpha_m^x = \gamma\}$. Fix $\gamma \in S_\kappa^\omega - S$ with $\gamma \in \lim \text{suc}_T(t)$ for any $t \in \mathcal{T} \cap \gamma^{<\omega}$, and an unbounded set $\{\gamma_n : n < \omega\} \subset \gamma$. Build inductively a branch $b : \omega \rightarrow \gamma$ through $\mathcal{T}$ with $\gamma_n \leq b(n)$ for any $n < \omega$. Then $\bigcap_{n < \omega} S_{b|_n} \subset \{x \in S_\emptyset : x \cap \kappa = \gamma\} = \emptyset$, as desired. $\square$

For the rest of the paper we give some cases where I has a winning strategy in the game $\mathcal{G}_1(\mathcal{C}_{\kappa\lambda})$, and show optimality of Goldring's result in a different way. Let us begin by the following observation, which should be folklore like non- ω_1 -Baireness of F^+ for a fine filter F on $\mathcal{P}_{\omega_1} \lambda$:

Proposition 3. *Let F be a normal filter on $\mathcal{P}_{\omega_1} \lambda$. Then I has a winning strategy in $\mathcal{G}_1(F)$.*

Proof. We let $\mathcal{P}_{\omega_1} \lambda$ be the first move of I, and for $x \in \mathcal{P}_{\omega_1} \lambda$ fix a list $\{\alpha_n^x : n < \omega\}$ of x . Next suppose that $T \in F^+$ is an n -th move of II. Then by the normality of F we have $\gamma_n < \lambda$ with $\{x \in T : \alpha_n^x = \gamma_n\} \in F^+$, which we let be the next move of I.

Let $\langle S_n : n < \omega \rangle$ be a play of I following the above strategy with $\langle \gamma_n : n < \omega \rangle$ as above. Then $\bigcap_{n < \omega} S_n = \{\gamma_n\}$, as desired. $\square$

Our first result, Proposition 4 follows from Proposition 3 and our Theorem below.

We nevertheless present it here because of the following lemma, which refines a fact

from [BT] and seems of independent interest.

Lemma 2. *Let $\lambda < \kappa^{+\omega}$. Then there is a stationary set $S \subset \{x \in \mathcal{P}_\kappa \lambda : \kappa \leq \forall \mu \leq \lambda (\text{cf } \sup(x \cap \mu) = \omega)\}$ on which the map $x \mapsto \langle \sup(x \cap \mu) : \kappa \leq \mu \leq \lambda \rangle$ is injective.*

Proof. We go by induction on $\lambda < \kappa^{+\omega}$. For $\lambda = \kappa$ just note that S_κ^ω is stationary in $\mathcal{P}_\kappa \kappa$. Next suppose that $S \subset \mathcal{P}_\kappa \lambda$ is as above. Then for $\gamma \in S_{\lambda^+}^\omega - \lambda$ $S_\gamma = \{x \in \mathcal{P}_\kappa \gamma : \pi_\gamma^{-1} x = x \cap \lambda \in S \wedge \sup x = \gamma\}$ is stationary, and for any $x, y \in S_\gamma$ with $\langle \sup(x \cap \mu) : \kappa \leq \mu \leq \lambda \rangle = \langle \sup(y \cap \mu) : \kappa \leq \mu \leq \lambda \rangle$ we have $x = \pi_\gamma''(x \cap \lambda) = \pi_\gamma''(y \cap \lambda) = y$, since $x \cap \lambda = y \cap \lambda \in S$. We claim that $\bigcup \{S_\gamma : \gamma \in S_{\lambda^+}^\omega - \lambda\}$ is stationary in $\mathcal{P}_\kappa \lambda^+$. Fix $f : (\lambda^+)^{<\omega} \rightarrow \lambda^+$. Take $\gamma \in S_{\lambda^+}^\omega - \lambda$ with $f''\gamma^{<\omega} \subset \gamma$. Then by stationarity of S_γ some $x \in S_\gamma$ with $x \cap \kappa \in \kappa$ is closed under f , as desired. $\square$

Now the rest of the proof goes as in Proposition 1:

Proposition 4. *I has a winning strategy in $\mathcal{G}_1(\mathcal{C}_{\kappa\lambda})$ when $\lambda < \kappa^{+\omega}$.*

Proof. We let the stationary set as in Lemma 2 be the first move S_0 of I, and for $x \in S_0$ and a cardinal $\kappa \leq \mu \leq \lambda$ fix an unbounded set $\{\alpha_{\mu,n}^x : n < \omega\} \subset x \cap \mu$. Next suppose that T is an n -th move of II. Then by finite applications of the regressive function lemma we have $\langle \gamma_{\mu,n} : \kappa \leq \mu \leq \lambda \rangle$ such that $\{x \in T : \kappa \leq \forall \mu \leq \lambda (\alpha_{\mu,n}^x = \gamma_{\mu,n})\}$ is stationary, which we let be the next move of I.

Let $\langle S_n : n < \omega \rangle$ be a play of I following the above strategy with $\langle \gamma_{\mu,n} : n < \omega \rangle$ as above. Then $\bigcap_{n < \omega} S_n \subset \{x \in S_0 : \kappa \leq \forall \mu \leq \lambda (\sup(x \cap \mu) = \sup_{n < \omega} \gamma_{\mu,n})\}$, which is at most a singleton, as desired. $\square$

For our main result below, we are indebted to Shelah's notion [S] of internally

approachable submodels of H_θ (see also [BM]), as well as Baumgartner's style [B] of purely combinatorial proofs.

Theorem. *I has a winning strategy in $\mathcal{G}_1(\mathcal{C}_{\kappa\lambda})$ when $\kappa > \omega_1$ and $\lambda < \kappa^{+\omega_1}$.*

Proof. Let $\{\mu_i : i < \omega\}$ list (not necessarily injectively) the set of regular cardinals between κ and λ . First we claim that $S_0 = \{\bigcup_{n < \omega} x_n : \forall n < \omega (x_n \in \mathcal{P}_\kappa \lambda \wedge x_n \cap \kappa \in \kappa \wedge x_n \cup \{\sup(x_n \cap \mu_i) : i < \omega\} \subset x_{n+1} \wedge \forall \alpha \in x_n (x_n \text{ is closed under } \pi_\alpha \text{ and } \pi_\alpha^{-1}) \wedge \forall i < \omega (\text{cf } \sup(x_n \cap \mu_i) = \omega_1 \wedge x_n \cap \mu_i \text{ contains a club subset of } \sup(x_n \cap \mu_i)))\}$ is stationary in $\mathcal{P}_\kappa \lambda$.

Fix $f : \lambda^{<\omega} \rightarrow \lambda$. We build inductively $\langle x_n : n < \omega \rangle$ as above with $\bigcup_{n < \omega} x_n$ closed under f as follows: First set $x_0 = \bigcup_{\xi < \omega_1} x_{0,\xi}$, where $x_{0,\xi}$ is defined inductively by $x_{0,0} = \{\mu_i : i < \omega\}$, $x_{0,\xi+1} = x_{0,\xi} \cup \sup(x_{0,\xi} \cap \kappa) \cup \bigcup \{\pi_\alpha^{-1}(x_{0,\xi} \cap \alpha) : \alpha \in x_{0,\xi}\} \cup \bigcup \{\pi_\alpha^{-1}(x_{0,\xi} \cap \alpha) : \alpha \in x_{0,\xi}\} \cup \{\sup(x_{0,\xi} \cap \mu_i) : i < \omega\}$ and $x_\xi = \bigcup_{\zeta < \xi} x_{0,\zeta}$ for a limit ξ . Next set $x_{n+1} = \bigcup_{\xi < \omega_1} x_{n+1,\xi}$, where $x_{n+1,\xi}$ is defined inductively by $x_{n+1,0} = x_n \cup f^{\leq \omega}(x_n \cap \{\sup(x_n \cap \mu_i) : i < \omega\})$, $x_{n+1,\xi+1} = x_{n+1,\xi} \cup \sup(x_{n+1,\xi} \cap \kappa) \cup \bigcup \{\pi_\alpha^{-1}(x_{n+1,\xi} \cap \alpha) : \alpha \in x_{n+1,\xi}\} \cup \bigcup \{\pi_\alpha^{-1}(x_{n+1,\xi} \cap \alpha) : \alpha \in x_{n+1,\xi}\} \cup \{\sup(x_{n+1,\xi} \cap \mu_i) : i < \omega\}$ and $x_\xi = \bigcup_{\zeta < \xi} x_{n+1,\zeta}$ for a limit ξ . Then $\{\sup(x_{n,\xi} \cap \mu_i) : \xi < \omega_1\} \subset x_n$ is club in $\sup(x_n \cap \mu_i)$ for any $n, i < \omega$.

We let S_0 be the first move of I, and for $x \in S_0$ fix $\langle x_n : n < \omega \rangle$ as above with $x = \bigcup_{n < \omega} x_n$. Next suppose that T is an n -th move of II. Then by $n+1$ applications of the regressive function lemma we have $\langle \gamma_{n,i} : i \leq n \rangle$ such that $\{x \in T : \forall i \leq n (\sup(x_n \cap \mu_i) = \gamma_{n,i})\}$ is stationary, which we let be the next move of I.

Let $\langle S_n : n < \omega \rangle$ be a play of I following the above strategy with $\langle \gamma_{n,i} : i \leq n < \omega \rangle$

as above. To see that $|\bigcap_{n<\omega} S_n| \leq 1$, fix $x, y \in \bigcap_{n<\omega} S_n$. By induction on cardinals $\kappa \leq \mu \leq \lambda$ we show that $x \cap \mu = y \cap \mu$.

First take $i < \omega$ with $\mu_i = \kappa$. Then $x \cap \kappa = \bigcup_{i \leq n < \omega} x_n \cap \kappa = \bigcup_{i \leq n < \omega} \gamma_{n,i} = \bigcup_{i \leq n < \omega} y_n \cap \kappa = y \cap \kappa$, as desired.

Next suppose that $x \cap \mu = y \cap \mu$ with $\mu < \lambda$. Take $i < \omega$ with $\mu_i = \mu^+$. Fix $i \leq n < \omega$. Then $x_n \cap y_n \cap \mu^+$ is unbounded in $\sup(x_n \cap \mu^+) = \sup(y_n \cap \mu^+) = \gamma_{n,i}$. Thus $x \cap \gamma_{n,i} = y \cap \gamma_{n,i}$, since $x \cap \alpha = \bigcup_{n \leq j < \omega} x_j \cap \alpha = \bigcup_{n \leq j < \omega} \pi_\alpha''(x_j \cap \mu) = \pi_\alpha''(x \cap \mu) = \pi_\alpha''(y \cap \mu) = \bigcup_{n \leq j < \omega} \pi_\alpha''(y_j \cap \mu) = \bigcup_{n \leq j < \omega} y_j \cap \alpha = y \cap \alpha$ for any $\alpha \in x_n \cap y_n \cap (\mu^+ - \mu)$. Now we have $x \cap \mu^+ = \bigcup_{i \leq n < \omega} x \cap \gamma_{n,i} = \bigcup_{i \leq n < \omega} y \cap \gamma_{n,i} = y \cap \mu^+$ as desired, since $\sup(x \cap \mu^+) = \sup(y \cap \mu^+) = \sup_{i \leq n < \omega} \gamma_{n,i}$.

The limit case is clear from the inductive hypothesis. $\square$

REFERENCES

- [B] J. Baumgartner, *On the size of the closed unbounded sets*, Ann. Pure Appl. Logic **54** (1991), 195–227.
- [BT] J. Baumgartner and A. Taylor, *Saturation properties of ideals in generic extensions. I*, Trans. Amer. Math. Soc. **270** (1982), 557–574.
- [BM] M. Burke and M. Magidor, *Shelah's pcf theory and its applications*, Ann. Pure Appl. Logic **50** (1990), 207–254.
- [FMS] M. Foreman, M. Magidor and S. Shelah, *Martin's Maximum, saturated ideals and non-regular ultrafilters. Part I*, Ann. Math. **127** (1988), 1–47.
- [GJM] F. Galvin, T. Jech and M. Magidor, *An ideal game*, J. Symbolic Logic **43** (1978), 284–292.
- [G] N. Goldring, *The entire NS ideal on $\mathcal{P}_{\gamma\mu}$ can be precipitous*, J. Symbolic Logic **62** (1997), 1161–1172.
- [J] T. Jech, *A game theoretic property of Boolean algebras*, Logic Colloquium '77, North Holland, Amsterdam, 1978, pp. 135–144.
- [JP] T. Jech and K. Prikry, *Ideals over uncountable sets: Application of almost disjoint functions and generic ultrapowers*, Mem. Amer. Math. Soc. **214** (1979).
- [K] A. Kanamori, *The Higher Infinite*, Springer, Berlin, 1994.
- [MS] Y. Matsubara and M. Shioya, *Nowhere precipitousness of some ideals*, J. Symbolic Logic **63** (1998), 1003–1006.
- [S] S. Shelah, *Cardinal Arithmetic*, Oxford University Press, Oxford, 1994.