

ある作用素平均族のべき単調性

Power monotonicity for a class of operator means

大阪教育大学・教育協働学科 藤井 淳一

Jun Ichi Fujii

Department of Educational collaboration

Osaka Kyoiku University

1 Introduction.

The theory of operator means is established by Kubo and Ando [4]: An *operator mean* $A \mathfrak{m} B$ for positive invertible operators A, B is defined by a positive normalized operator monotone function f on $(0, \infty)$ by

$$A \mathfrak{m} B = A^{\frac{1}{2}} f \left(A^{-\frac{1}{2}} B A^{-\frac{1}{2}} \right) A^{\frac{1}{2}}.$$

Here the *normalization* is $f(1) = 1$. One of the result of the Kubo-Ando theory is to give the bijection between the operator means and the positive normalized operator monotone functions on $(0, \infty)$ as above. In this bijection, f is often called the *representing function* of an operator mean $f(x) = 1 \mathfrak{m} x$.

Related to this, we gave a path of positive function on $(0, \infty)$ in [2];

$$F_r(x) = \frac{3r - 1}{3r + 1} \frac{x^{\frac{3r+1}{2}} - 1}{x^{\frac{3r-1}{2}} - 1}.$$

It is monotone increasing for $r \in \mathbb{R}$, which is the result of Takahasi, Tsukada, Tanahashi and Ogiwara [5]. For $r \in [-1, 1]$, they are positive normalized operator monotone functions. Moreover they are symmetric:

$$A \mathfrak{m}_f B = B \mathfrak{m}_f A, \quad \text{that is,} \quad f(x) = x f \left(\frac{1}{x} \right).$$

Typical means are listed below as numerical ones:

r	-1	$-1/3$	0	$1/3$	1
$a \mathfrak{m}_r b = a F_r(b/a)$	$\frac{2ab}{a+b}$	$L^*(a, b) = \frac{b \log \frac{a}{b}}{a-b}$	$\sqrt{ab}$	$L(a, b) = \frac{a-b}{\log \frac{a}{b}}$	$\frac{a+b}{2}$

Transformed by $s = \frac{3r-1}{2}$, it is equivalent to the following path:

$$\tilde{F}_s(x) = \frac{s}{s+1} \frac{x^{s+1} - 1}{x^s - 1},$$

which is also discussed in [3].

Recently Wada [6] introduced the power monotonicity of representing functions f and showed the relation to the Ando-Hial inequality [7, 1]: f is called *PMI* (resp., *PMD*) if f satisfies

$$f(x^p) \leq f(x)^p \quad (\text{resp., } f(x^p) \geq f(x)^p) \quad \text{for all } p > 1.$$

In this note, we show the power monotonicity for F_r . Incidentally we see the role of the terminal means in these inequalities.

2 Main result.

Theorem 1. *The function F_r is PMI for $r \geq 0$ and PMD for $r \leq 0$.*

For $\tilde{F}_s$, this result is equivalent to:

Theorem F. *The function $\tilde{F}_s$ is PMI for $s \geq -\frac{1}{2}$ and PMD for $s \leq -\frac{1}{2}$.*

In fact, we show Theorem 1' since $\tilde{F}_s$ has simple parameters. To show this, we need two lemmas due to Takahasi-Tsukada-Tanahashi-Hagiwara [5]. For completeness, we give each proof. First, we see the following property:

Lemma 2. *The function $J(t) = \begin{cases} \frac{e^t}{e^t-1} - \frac{1}{t} & (t \neq 0) \\ \frac{1}{2} & (t = 0) \end{cases}$ is monotone increasing.*

It is easy to see that

$$\text{For } x > 1, G_x(s) = \begin{cases} \log \frac{x^s - 1}{s} & (s \neq 0) \\ \log(\log x) & (s = 0) \end{cases} \text{ is monotone increasing.} \quad (*)$$

Combining these, we have:

Corollary 3. *For $x > 1$, $G'_x(s)$ is monotone increasing.*

Incidentally, these results show the known property: $\tilde{F}_s$ is monotone increasing for s , which is the required result in [5]. In fact, by $\log \tilde{F}_s(x) = G_x(s+1) - G_x(s)$,

$$\frac{\partial \log \tilde{F}_s(x)}{\partial s} = G'_x(s+1) - G'_x(s) \geq 0.$$

Thus $\log \tilde{F}_s$ is monotone increasing for s when $x > 1$. As for the case $0 < x < 1$, we have

$$\tilde{F}_s(x) = \frac{s}{s+1} \frac{x^{s+1} - 1}{x^s - 1} = \frac{s}{s+1} \frac{\left(\frac{1}{x}\right)^{s+1} - 1}{\left(\frac{1}{x}\right)^s - 1} \times x,$$

which also shows the monotonicity for s . Thus it holds for all $x > 0$.

Remark. The representing function $S_\alpha(x)$ of the *Stolarsky mean* is defined by

$$s_\alpha(t) = \left(\frac{t^\alpha - 1}{\alpha(t - 1)} \right)^{\frac{1}{\alpha-1}}.$$

S.Wada [6, Prop.3.2] showed that s_α is PMD on $[-2, -1]$ and PMI on $[-1, 2]$. Putting $\alpha = \frac{s+1}{s}$, $t = x^s$, we have

$$\tilde{S}_s(x) = \left(\frac{s}{s+1} \frac{x^{s+1} - 1}{x^s - 1} \right)^s,$$

which is closely related to our path $\tilde{F}_s$. In fact, for $s > 0$, we have $\tilde{F}_s$ is PMD (resp., PMI) on $\mathcal{I}$ if and only if $\tilde{S}_s$ is PMD (resp., PMI) on $\mathcal{I}$. For the negative case, these concepts are exchanged. Then we show Wada's result directly implies that F_r is PMD for $r \in (-\frac{1}{3}, 0)$ and that it is PMI for $r \in (0, \frac{1}{9}) \cup (1, \infty)$. Thus Wada's result does not imply all our results in the above theorem.

In fact, consider the PMD case:

$$-2 < \alpha = \frac{s+1}{s} < -1 \iff -\frac{1}{2} < s = \frac{3r-1}{2} < -\frac{1}{3} \iff 0 < r < \frac{1}{9},$$

which shows F_r is PMI for $r \in (0, \frac{1}{9})$ by $s < 0$.

Next consider the PMI case, which is divided into the negative case and the positive one: For $(-1 <) s < 0$, we have

$$-1 < \alpha = \frac{s+1}{s} < 0 \iff -1 < s = \frac{3r-1}{2} < -\frac{1}{2} \iff -\frac{1}{3} < r < 0,$$

which shows F_r is PMD for $r \in (-\frac{1}{3}, 0)$. Lastly, for $s > 0$ we have

$$0 < \alpha = \frac{s+1}{s} < 2 \iff s > 1 \iff r > 1,$$

which shows F_r is PMI for $r \in (1, \infty)$.

3 Relation to the terminal means

Restricting ourselves to the case $p = n$, integers and $|r| \geq \frac{1}{3}$. Then, we show the following partial result of Theorem 1 via the arithmetic or harmonic means:

Theorem 4. For any integer n and $r \geq \frac{1}{3}$,

$$F_r(x^n) - F_r(x)^n \geq F_r(x) \left(\left(\frac{x+1}{2} \right)^{n-1} - F_r(x)^{n-1} \right) \geq 0.$$

For $r \leq -\frac{1}{3}$,

$$F_r(x)^n - F_r(x^n) \geq F_r(x) \left(F_r(x)^{n-1} - \left(\frac{2x}{1+x} \right)^{n-1} \right) \geq 0.$$

To see this, we give a lemma:

Lemma 5. A function $g_n(r) = \frac{\sum_{\ell=0}^{n-1} x^{\frac{\ell(3r+1)}{2}}}{\sum_{k=0}^{n-1} x^{\frac{k(3r-1)}{2}}} = \frac{F_r(x^n)}{F_r(x)}$ is monotone increasing.

4 Concluding remark

Very recently, Yamazaki extend Theorem 4 to the following:

Theorem (Yamazaki). For $p, q \in [-1, 1]$, the represenming function

$$F_{p,q}(x) = \left(\frac{p}{p+q} \frac{x^{p+q} - 1}{x^p - 1} \right)^{\frac{1}{q}}$$

is PMI for $2p + q \geq 0$, and PMD for $2p + q \leq 0$.

This theorem is shown by the following integral representation:

$$F_{p,q}(x) = \left(\int_0^1 (1-t+tx^p)^{\frac{q}{p}} dt \right)^{\frac{1}{q}} \quad \text{Consider:}$$

Acknowledgement. This study is partially supported by the Ministry of Education, Science, Sports and Culture, Grant-in-Aid for Scientific Research (C), JSPS KAKENHI Grant Number JP 16K05253.

References

- [1] T.Ando and F.Hiai, Log majorization and complementary Golden-Thompson type inequality, Linear Algebra Appl., **197**(1994), 113–131.
- [2] J.I.Fujii and Y.Seo, On parametrized operator means dominated by power ones, Sci. Math., **1**(1998), 301–306.
- [3] F.Hiai and H.Kosaki, Means for matrices and comparison of their norms, Indiana Univ. Math. J., **48** (1999), 899–936.
- [4] F.Kubo and T.Ando, Means of positive linear operators, Math. Ann., **246**(1980), 205–224.
- [5] S.-E.Takahasi, M.Tsukada, K.Tanahashi and T.Ogiwara, An inverse type of Jensen's inequality, Math. Japon., **50**(1999), 85–92.
- [6] S.Wada, Some ways of constructing Furuta-type inequalities, Linear Algebra Appl., **457**(2014), 276–286.
- [7] S.Wada, When does Ando-Hiai inequality hold?, Linear Algebra Appl., **540**(2018), 234–243.

Department of Educational collaboration
 Osaka Kyoiku University
 Osaka 582-8582
 JAPAN
 Email address: fujii@cc.osaka-kyoiku.ac.jp