

**Hydrothermal fluid flow system around the Iheya North Knoll in the
mid-Okinawa Trough based on seismic reflection data**

Takeshi Tsuji^{1, 2}, Ken Takai², Hisashi Oiwane³, Yasuyuki Nakamura³, Yuka Masaki^{2, 4},
Hidenori Kumagai², Masataka Kinoshita², Fujio Yamamoto², Tadashi Okano², and
Shin'ichi Kuramoto²

¹Graduate school of Engineering, Kyoto University

²Japan Agency for Marine Earth Science and Technology (JAMSTEC)

³Ocean Research Institute, University of Tokyo

⁴Department of Applied Science, Kochi University

Research Highlights (4 bullet points)

(1) Possible fluid flow paths in the Iheya North Knoll hydrothermal field

(2) Widespread permeable volcanic deposits produced by silicic arc volcanism

(3) Layered sequence focuses migration of fluids derived from trough-fill sediments

(4) Fluid alteration accelerated by interactions with sediments

24 **Abstract**

25 **Seismic reflection data around the Iheya North Knoll hydrothermal field provide**
26 **insights into geological structures that control subseafloor hydrothermal fluid flow**
27 **in the sediment-covered continental backarc basin of the mid-Okinawa Trough.**
28 **We identified the seismic expression of widespread porous volcanoclastic**
29 **pumiceous deposits and intrusions as a result of silicic arc volcanism. The porous**
30 **and permeable volcanic deposits distribute in an area extending updip from the**
31 **thick succession of the deep trough to the seafloor at the hydrothermal field. Their**
32 **regional structure focuses the flow of hydrothermal fluids derived from the**
33 **surrounding trough-fill sediments and directs them to the vents of the**
34 **hydrothermal field. The high concentrations of CH₄ and NH₄ in the fluids of the**
35 **hydrothermal field are likely derived from the interaction of migrating fluids with**
36 **trough-fill sediments.**

37

38 **Keywords:** Hydrothermal fluid paths, Okinawa Trough, seismic reflection data,
39 continental backarc basin, silicic volcanism

40

41

1. Introduction

The circulation of hydrothermal fluids enhances chemical and thermal exchange and the interaction between the crust and ocean (e.g., Stein and Stein, 1992). Because of the considerable influence of hydrothermal fluid circulation within oceanic lithosphere on the physical, chemical, and biological evolution of the crust and ocean, it has been well studied at spreading axes and on their flanks (e.g., Fisher et al., 1997; Fisher, 1998; Johnson et al., 2000; Fisher et al., 2003; Hutnak et al., 2008; Fisher et al., 2008; Waldhauser and Tolstoy, 2011; Fisher et al., 2011). In contrast, there have been few studies of hydrothermal fluid circulation in subduction zones, such as at island arc submarine volcanoes and in continental backarc basins. Because backarc basins at continental margins are usually covered by thick sequences of terrigenous sediments, an understanding of fluid recharge and migration paths cannot be obtained only from near-seafloor observations.

The hydrothermal field on Iheya North Knoll, about 150 km NNW of Okinawa Island, was discovered in 1995 (Fig. 1; Momma et al., 1996) and has since been investigated during more than 40 dives of manned deep submergence vehicles (DSVs) and remotely operated vehicles (ROVs). These investigations have demonstrated that prosperous subseafloor microbial communities associated with the hydrothermal fluid

chemistry are hosted along the entire hydrothermal fluid flow path, from sedimentary pore water to hydrothermal discharge zones (Nakagawa et al., 2005; Takai et al., 2006). The chemistry of the hydrothermal fluids in the Okinawa Trough (a backarc basin) is unusual among deep-sea hydrothermal fluids studied to date (Ishibashi and Urabe, 1995; Gamo et al., 2006; Takai et al., 2006; Takai and Nakamura, 2010; Kawagucci et al., 2011). High concentrations of CO₂ and CH₄ are common in fluids in hydrothermal systems underlying sedimentary sequences, such as at the Juan de Fuca Ridge (Lilley et al., 1993; Shanks et al., 1995) and in the Guaymas Basin (Welhan and Lupton, 1987; Pearson et al., 2005). However, the concentrations of CO₂ and other gaseous carbon compounds in hydrothermal fluids from the Okinawa Trough are among the highest in the world (Gamo, 1995; Konno et al., 2006; Takai and Nakamura, 2010; Kawagucci et al., 2011). Most of these volatiles are likely derived from subseafloor interactions between sediments and infiltrated seawater (Kawagucci et al., 2011). In addition, vent fluids from Okinawa Trough hydrothermal systems have high concentrations of Ba, provided by hydrothermal reactions between felsic volcanic rocks and hydrothermal fluids (Gamo et al., 2006). Despite the unusual fluid chemistry of the Okinawa Trough hydrothermal systems, there has been little investigation of their geological and hydrogeological setting. Understanding the hydrogeology of hydrothermal systems is of

fundamental importance in studying the interactions among geological, geochemical, and microbial processes in the deep-sea hydrothermal environment of a continental backarc basin. In this study, we investigated the structure of the hydrothermal fluid flow system of the Iheya North Knoll hydrothermal field on the basis of our interpretation of a dense grid of seismic reflection data covering the Iheya North Knoll.

2. Geologic setting

The Okinawa Trough lies in the eastern East China Sea, between the Ryukyu Arc – Trench system and the Eurasian continent (Fig. 1a). In this area, the Okinawa Trough is considered to be in the initial stage of continental rifting, although there is some argument on the commencement and order of the Okinawa Trough rifting (e.g., Letouzey and Kimura, 1985). Most previous studies suggest that rifting commenced in the southern part of the trough (e.g., Sibuet et al., 1995; Hsu et al., 2001) and that it started at ~2 Ma. The Yangtze River is the main source of the thick terrigenous sediments that cover the crust of the mid-Okinawa Trough (Narita et al., 1990; Huh and Su, 1999), where the sedimentation rate has been estimated to be 1–2 m ka⁻¹ (Tsugaru et al., 1991).

The Iheya North Knoll is on the northern termination of the depression of the

96 Okinawa Trough (27.8°N, 126.9°E; Fig. 1) where water depths are greater than 1000 m.
97 The volcanic body of the Iheya North Knoll penetrates through a thick sedimentary
98 sequence and rises about 500 m above the seafloor. The hydrothermal field lies on the
99 eastern flank of the western peak of the knoll, facing the central valley (Fig. 1c). Several
100 hydrothermal mounds have been recognized during previous surveys (e.g., Takai et al.,
101 2006; Takai and Nakamura, 2010). They are clearly shown in high-resolution
102 bathymetric data (Fig. 2) obtained by autonomous underwater vehicle (AUV) *Urashima*
103 equipped with acoustic sounding equipment (SeaBat7125, 400 Hz source) during cruise
104 YK07-07. The hydrothermal mounds are aligned roughly N–S (Fig. 2).

105 The highest temperatures (maximum 311 °C) and highest hydrothermal flow rates
106 have been recorded at the vent known as North Big Chimney (NBC; Fig. 2) over a
107 period of more than ten years (e.g., Kawagucci et al., 2011), indicating that the
108 hydrothermal fluid vented from NBC comes from the main hydrothermal flow path in
109 this region. The high temperature of discharged fluid (~300 °C) at NBC suggests a
110 magma-driven system. Both temperatures and flow rates of other vents in the area
111 decrease with increasing distance from NBC (Masaki et al., 2011). Furthermore,
112 intensive hydrothermal plumes have been imaged by side-scan sonar in seawater only
113 around the hydrothermal field (Kumagai et al., 2010).

Recent Integrated Ocean Drilling Program (IODP), conducting a series of drill holes in the central valley of the Iheya North Knoll, showed that the sediments there are mainly pelagic and hemipelagic mud and volcanoclastic deposits (pumiceous gravels), and that they have been variably altered by hydrothermal processes (Expedition 331 Scientists, 2010). Drilling data revealed an alternating sequence of hard low-porosity layers and porous layers that were hydrothermally altered at high temperatures. The hard layers, presumably of low permeability, may provide the cap rock above hydrothermal fluid flow paths. The alignment of the active hydrothermal vents along the strike of the sedimentary layers (N–S; Fig. 2) suggests that the hydrothermal fluid flow paths are controlled by the layered sedimentary structures.

3. Seismic data acquisition and processing

We used 2D seismic reflection data acquired along 54 survey lines recorded in a dense orthogonal grid during four cruises (Table 1, Figs. 1b and 1c). Multi-channel seismic reflection data were recorded during cruises KY02-11, KY07-03, and KR10-02, and single-channel data during cruise YK06-09. During cruise KR10-02, long-offset (~4.5 km) multi-channel data were acquired in order to determine the velocity structure required to calculate depths of reflection events. A larger volume airgun was also used

during cruise KR10-02 (Table 1).

The seismic data from cruises KY02-11 and KY07-03 were processed with a conventional processing sequence that included spherical divergence correction, deconvolution, common midpoint (CMP) sorting, normal moveout (NMO) correction, multiple suppression with a Radon filter, CMP stacking, migration, and bandpass filtering (Yilmaz, 2001). The short streamer deployed during cruises KY02-11 and KY07-03 (Table 1) did not allow estimation of accurate seismic interval velocities for deep reflectors. However, shallow geological structures were clearly imaged on these profiles (KY02-11 and KY07-03) and the single-channel data (KR06-09), because of the high resolution (high frequency) of the source signals used (Fig. 5).

To obtain accurate depths of subsurface geological structures for data from cruise KR10-02, we applied prestack depth migration incorporating an iterative horizon-based tomographic analysis of seismic velocity (see Figs. 3 and 4). We determined seismic velocities at six horizons, including the sediment–crust interface. Because the dominant frequency of the cruise KR10-02 data was lower than that of the other cruises, we used these data mainly for interpretation of the deep geological structure (Figs. 3 and 4).

Although 2D seismic reflection data do not unequivocally resolve structures in 3D (French, 1974), the dense grid of orthogonal profiles we used allowed us to identify the

3D subseafloor structures with reasonable confidence. Furthermore, by comparing orthogonal profiles we were able to identify sideswipe reflections from nearby areas of elevated seafloor. Because the seismic interval velocities necessary for depth conversion could not be accurately determined within the knoll, we used 3D fence diagrams constructed from several time-domain reflection profiles (Fig. 6) to develop a representation of both subseafloor geological structures and possible hydrothermal fluid paths.

To interpret and characterize the subseafloor structures (e.g., volcanic deposits), we calculated seismic attributes (e.g., amplitude envelope; Fig. 7) (Taner et al., 1979; Tsuji et al., 2005). Magmatic intrusions can be imaged as blocks on the envelope profile (Fig. 7) mainly because the envelope ignores phase information; the intrusions disturb the original structure of sedimentary sequence (Tsuji et al., 2007).

4. Volcanic activity and pumiceous deposits

The seismic profiles across the Iheya North Knoll reveal the subseafloor volcanic and sedimentary structures in the area under and surrounding the knoll. The subseafloor rocks of the Iheya North Knoll volcano are characterized by chaotic reflections that extrude through trough sediments of >1500 m thickness (Figs. 3 and 4). The depth of

the seafloor around the knoll is greater on the southwestern side of the knoll (Figs. 3 and 4) because of ongoing tectonic rifting in the Okinawa Trough. The seismic reflection profiles did not allow us to identify a heat source beneath the knoll because the complicated subsurface structure and steep seafloor terrain of the knoll made it difficult to resolve deep structures. The magma reservoir probably lies below the knoll at a depth beyond the resolution of our seismic data. However, southwest of the knoll we identified strong low-frequency reflectors beneath the trough sediments (~4 km below seafloor; red arrows in Fig. 4) that possibly represent a magmatic body. Numerous normal faults are developed in the sediments overlying the possible magmatic body and dip toward it (Fig. 4).

Some of the peaks of the Iheya North Knoll show intervals of low-amplitude reflections at and beneath the seafloor (Fig. 7a), including the peak east of the central valley (Fig. 5a). This seismic transparency indicates a lack of acoustic impedance contrast over these intervals, indicating that they represent intervals of homogeneous and porous rock. Piston cores obtained at the peak above the transparent interval during cruise KY08-01 (labeled A in Figs. 1c and 7a) are composed of volcanoclastic pumiceous deposits interbedded with silt (Oiwane et al., 2008), which would be expected to be seismically transparent. Thus, the transparent intervals likely represent

volcaniclastic pumiceous deposits.

Pumiceous breccias are commonly observed within sediment cores obtained in and around the Iheya North Knoll (Expedition 331 Scientists, 2010). In the central valley of the Iheya North Knoll (the depression east of the hydrothermal field; Fig. 2), piston cores revealed several pumice layers of ~10 cm thickness within 3 m below the seafloor (labeled B in Figs. 1c and 5b) (Oiwane et al., 2008). On the eastern side of the knoll, a volcaniclastic layer of about 40 cm thickness is interbedded within an approximately 5-m-thick sequence of mud-dominated trough sediments (labeled C in Fig. 1c). Our seismic data cannot resolve such thin pumiceous layers (e.g., Rayleigh's criterion; Sheriff, 2002). However, we noted high-amplitude reflectors extending horizontally away from the Iheya North Knoll (white dashed lines in Fig. 7b), some of which are discordant with the layered sedimentary sequence in the trough. These may represent thick submarine volcaniclastic deposits, lava flows, or sills. Furthermore, a low-frequency, high-amplitude reflection at the seafloor trace of a normal fault (gray dashed line in Fig. 7c) may represent a magmatic body extruded on the seafloor after upwelling via the normal fault. This observation also explains frequent intrusions (complex network of volcanic deposits) within trough sediment.

5. Reservoir formation and hydrothermal fluid pathways

5.1 Small-scale fluid circulation

In the central valley of the Iheya North Knoll, there are repetitive east-dipping seismic reflectors within the sediments immediately below the seafloor (Fig. 5b), suggesting the existence of a layered sedimentary structure. Core samples demonstrate that the layered structure within this sedimentary sequence includes high-permeability pumiceous flow deposits (Oiwane et al., 2008; Expedition 331 Scientists, 2010). The permeability of the pumiceous gravels (Nakamura et al., 2008) obtained during drilling (Expedition 331 Scientists, 2010) was considerably higher than that of the mud-dominated pelagic sediments. Therefore, these highly permeable volcanic sequences might provide lateral migration pathways for pore fluids. Furthermore, the lateral distribution of high-permeability (pumiceous) layers overlain by low-permeability (pelagic and hemipelagic mud) layers may trap the fluid and provide reservoirs for subseafloor hydrothermal fluids. Whereas, this sedimentary sequence is also intersected by numerous steep normal faults, which might provide pathways for vertical (rather than lateral) flow of hydrothermal fluids.

Pyroclastic flow deposits may exist below the 200-m-thick layered sedimentary sequence that includes low-permeability mud, in the sequence between the black and

gray diamonds in Fig. 5b. This porous, high-permeability sequence dips basinward from the volcanic body near the hydrothermal field, and may provide a path for hydrothermal fluids to migrate laterally up the slope. Furthermore, the N–S trend of the vents of the hydrothermal field at the western edge of the low-permeability central valley sediments (Fig. 2) corresponds to the trend of the seafloor outcrop of the pyroclastic deposits (Fig. 5b), which supports the view that they may provide a conduit for hydrothermal fluid flow.

As one of the possible recharge entries around the central valley, a linear volcanic ridge located at ~2 km east of the hydrothermal vent sites (~30 m high and 500 m wide; Figs. 2, 5b and 8) is suggested on the basis of the heat flow measurements (*Masaki et al.*, 2011). Numerous heat flow data in the region of the knoll have shown temperature increasing with depth, but heat flow data from the sediments on the volcanic ridge reveal an inverse gradient (*Masaki et al.*, 2011). Furthermore, the heat flow values increase gradually with distance from the ridge, reaching a maximum at the hydrothermal vents. The weak seafloor reflection of the volcanic ridge (Fig. 5b) suggests a porous volcanoclastic pumiceous seafloor there. Therefore, the ridge may provide an entry for recharge of hydrothermal fluids in the pyroclastic flow deposits beneath the central valley.

If the volcanic ridge provides the primary hydrothermal fluid source at Iheya North Knoll, the hydrothermal fluid circulation cell is small, with only a few kilometers separating the main recharge and discharge areas (Fig. 5b). A circulation cell of this scale is comparable to those characteristic of fast-spreading mid-ocean ridges, where fluid residence times on the order of several to ten years have been estimated. A circulation cell of this scale would not sustain the steady hydrothermal fluid discharge, its chemistry, and the associating enormous chemosynthetic microbial and faunal communities of the Iheya North Knoll hydrothermal field (Nakagawa et al., 2005; Takai et al., 2006; Takai and Nakamura, 2010; Kawagucci et al., 2011). In particular, the carbon mass balance between hydrothermal fluid discharges and inorganic and organic carbon sources in the volcanic deposits and sediments within the Iheya Knoll structure clearly indicate that the hypothesized hydrothermal fluid paths could not support the CH₄ mass balance and stable isotope composition of the discharging hydrothermal fluids (Kawagucci et al., 2011). Thus, there must be other sources and pathways of hydrothermal fluids discharging at the Iheya North Knoll hydrothermal field.

5.2 Large-scale fluid circulation

On E–W seismic profile KY0703-18, we identified a strong west-dipping reflector

258 beneath Iheya North Knoll, the eastern end of which approaches the seafloor near the
259 hydrothermal vent sites (black circles in Fig. 5b). On the intersecting N–S seismic
260 profile (KY0703-E), this strong reflector dips to the north away from the hydrothermal
261 field (black circles in Fig. 5e). The high reflection amplitude (Fig. 5b) of this continuous
262 reflector may represent an important boundary that controls hydrothermal fluid flow
263 paths. The 3D structure of this reflector determined from our fence diagram of
264 orthogonal seismic profiles (Fig. 6) indicates that it extends away from the knoll to the
265 north and west, and that the shallowest part of the horizon converges on the
266 hydrothermal field. Furthermore, we noted several reflectors below and parallel to the
267 strong reflection, indicating a layered structures (gray circles in Figs. 5b and 5e). The
268 polarity of the strong west-dipping reflector is opposite to that of the seafloor reflector
269 (Fig. 5c), which indicates that the seismic velocity below the negative-polarity reflector
270 is lower than that of the overlying sequence. DSV and ROV observations of the small
271 peak immediately west of the hydrothermal field (e.g., NT07-11 cruise; Fig. 2) indicate
272 that most of the seafloor there is paved with potential sheet carbonate crust containing
273 numerous mussel shells and pieces of pumiceous breccia (Figs. 5b and 8). Based on the
274 pumiceous volcanic structures exposed at the seafloor and the layered structures below
275 and parallel to the negative-polarity reflector (gray circles in Fig. 5b), we interpret the

negative-polarity reflector and underlying low-velocity sequence to represent porous volcanic breccia.

Because the permeability of the layered sequence underlying the negative-polarity horizon is likely higher than those above and below it, large-scale permeability anisotropy would be expected within the layered sequence; low-permeability sediments above the layered sequence would provide a cap structure to prevent vertical upward migration of fluids. Furthermore, as already pointed out, the shallow eastern extent of the negative-polarity reflector approaches the hydrothermal field (Fig. 6). Thus, we suggest that the layered sequence immediately below the negative-polarity reflector provides a major conduit that gathers hydrothermal fluids and channels them to their point of discharge at the Iheya North Knoll hydrothermal field.

The large number of dead mussel shells on the seafloor above the shallowest point of the negative-polarity reflector (Figs. 2, 5b, and 8) suggest that past methanotrophic mussel populations was sustained by considerable amounts of hydrothermally derived methane seepage. The seafloor reflector above the negative-polarity reflector is stronger than that of the surrounding seafloor (Fig. 5e), suggesting that a consolidated carbonate crust is exposed at the seafloor there. Although it is still unclear how old the mussel shells and carbonate crust are formed, the volatile components of the hydrothermal

fluids such as CO₂ and CH₄ would escape from the hydrothermal reservoir by diffusion through the cap structure above the negative-polarity reflector, which may have been a less-effective seal in the past.

As shown by both the E-W and N-S seismic profiles, the negative-polarity event and underlying layered sequence extend into the thick trough-fill sediments that are widespread on the western and northern sides of the Iheya North Knoll (Figs. 5b and 5e). Considering the 3D structure of the negative-polarity horizon (Fig. 6), the large reservoir of pore water in the thick trough-fill sediments may provide the source of hydrothermal fluids, which then flow updip within the continuous permeable layered sequence before discharging at Iheya North Knoll hydrothermal field. Both the sedimentary sequence and the interbedded volcanic units were uplifted by the volcanic intrusion that formed the knoll. The resultant structure and the presence of a heat source beneath the knoll facilitate lateral migration of recharged fluids along the pumiceous layers toward the hydrothermal field. This conceptual hydrothermal system is consistent with the extraordinarily high concentrations of CH₄ and NH₄ in the hydrothermal fluids of the Iheya North Knoll hydrothermal field (Kawagucci et al., 2011). Rather, it is consistent with the prediction that the abundant CH₄ and NH₄ in the hydrothermal fluid should be derived from their reservoir in the more organics-rich, trough-filling

sediments accumulated by subseafloor microbial methanogenesis and ammonification in certain geological time scales (Kawagucci et al., 2011).

We identified many normal faults within the trough-fill sediments around the Iheya North Knoll (Fig. 4). The major faults strike parallel to the trough axis (ENE–WSW). Although the lithology is the same on both sides of a major normal fault on line YK-0703-18, the reflection amplitude east of the fault is weaker than that west of the fault (Fig. 7b). If the normal fault creates a hydrogeological boundary to lateral flow, the higher amplitudes west of the fault might represent active hydrothermal fluid flow and/or gas trapped within sedimentary sequence near the knoll. On the other hand, these major normal faults extend from the seafloor to deep crustal basement (Figs. 3 and 4; Lee et al., 1980) and, if fluid flow is driven by thermal advection from the deep heat source beneath the Iheya North Knoll, it is also possible that these deep crustal faults provide vertical hydrothermal fluid migration paths (Fig. 8).

6. Discussion and conclusions

In the region of mid-ocean ridges, the primary conduits for hydrothermal fluids are highly permeable zones of upper oceanic crust and abyssal-hill faults (Fisher, 1998; Fisher and Becker, 2000; Fisher et al., 2008; Hutnak et al., 2008; Fisher et al., 2011).

The thick sedimentary sequences that overlie basaltic crust on the flanks of mid-ocean ridges are generally considered to be impermeable to hydrothermal fluid flow. However, unlike the high-temperature fluids in sediment-hosted hydrothermal vent fields, the hydrothermal fluids there are not highly enriched with CH_4 and NH_4 (Wheat et al., 2003, 2004). A plausible explanation for this difference is that the primary fluid source on the ridge flank is nutrient-poor oxygenated seawater recharged directly through volcanic outcrops on the seafloor (Fisher et al., 2003) such that the chemistry of the hydrothermal fluids is little affected by geochemical and microbiological alteration during migration.

The Iheya North Knoll hydrothermal system is part of a continental backarc basin where subseafloor silicic arc volcanic deposits provide complex migration networks for recharge and discharge of sediment pore fluids. Taking into consideration the extraordinary CH_4 and NH_4 enrichment of the fluids of the Iheya North Knoll hydrothermal field, there are four important components in the model we propose here for the hydrothermal fluid flow system.

1. The primary source of hydrothermal fluids is the nutrient-rich, reductive pore fluids of the trough-fill sediments around the Iheya North Knoll.
2. The primary fluid recharge and migration paths are through permeable trough-fill

volcaniclastic sediments and normal faults that extend from the seafloor to crustal basement.

3. Geochemical and microbiological alteration of hydrothermal fluids passing through the complex network of flow paths enriches them in CH_4 and NH_4 .

4. The structural culmination around the knoll of the highly permeable volcaniclastic sequences that integrate multiple flow paths focuses fluid flow in the region of the hydrothermal vents.

Acknowledgements

We thank CDEX/JAMSTEC and Nippon Marine Enterprises (NME) for the acquisition of seismic reflection data and detailed bathymetric data. We thank S. Shimizu (NME) and Y. Kido and S. Uraki (CDEX/JAMSTEC) for their cooperation and assistance in seismic processing, and two reviewers for helpful comments. This research was supported by Grant-in-Aid for Scientific Research on Innovative Areas TAIGA “Structure of oceanic crust and its properties revealed by seismic reflection method: Insight into subseafloor hydrothermal fluid circulation” (21109502) and the Excellent Young Researcher Overseas Visit Program of the Japan Society for the Promotion of Science (21-5338).

References

- Expedition 331 Scientists, 2010. Deep hot biosphere. IODP Prel. Rept. 331.
doi:10.2204/iodp.pr.331.2010.
- Fisher, A. T., 1998. Permeability within basaltic oceanic crust, *Rev. Geophys.* 36,
143–182.
- Fisher, A.T., Becker, K., 2000. Reconciling heat flow and permeability data with a
model of channelized flow in oceanic crust, *Nature* 403, 71–74.
- Fisher, A.T., Becker, K., Davis, E.E., 1997. The permeability of young oceanic crust
east of Juan de Fuca Ridge determined using borehole thermal measurements.
Geophys. Res. Lett. 24, 1311–1314.
- Fisher, A.T., Davis, E.E., Hutnak, M., Spiess, V., Zuhlsdorff, L., Cherkaoui, A.,
Christiansen, L., Edwards, K., Macdonald, R., Villinger, H., Mottl, M.J., Wheat,
C.G., Becker, K., 2003. Hydrothermal recharge and discharge across 50 km guided
by seamounts on a young ridge flank, *Nature*, 421, 618–621,
doi:10.1038/nature01352.
- Fisher, A.T., Davis, E.E., Becker, K., 2008. Borehole-to-borehole hydrologic response

- 384 across 2.4 km in the upper oceanic crust: Implications for crustal-scale properties,
385 J. Geophys. Res., 113, B07106, doi:10.1029/2007JB005447.
- 386 Fisher, A.T., Tsuji, T., Petronotis, K., and the Expedition 327 Scientists, 2011. Proc.
387 IODP, 327: Tokyo (Integrated Ocean Drilling Program Management International,
388 Inc.). doi:10.2204/iodp.proc.327.2011.
- 389 French, W.S., 1974. Two-dimensional and three-dimensional migration of
390 model-experiment reflection profiles, Geophysics, 39, 265-277.
- 391 Gamo, T., 1995. Wide variation of chemical characteristics of submarine hydrothermal
392 fluids due to secondary modification processes after high temperature water-rock
393 interaction: a review. p. 425-451. *In* H. Sakai, and Y. Nozaki (eds.),
394 Biogeochemical processes and ocean flux in the western Pacific, Terra Scientific
395 Publisher, Tokyo.
- 396 Gamo, T., Ishibashi, J., Tsunogai, U., Okamura, K., Chiba, H., 2006. Unique
397 geochemistry of submarine hydrothermal fluids from arc-backarc settings of the
398 western Pacific. In "Back-Arc Spreading Systems: Geological, Biological, Chemical,
399 and Physical Interactions" (D. M. Christie, C. R. Fisher, S.-M. Lee and S. Givens
400 ed.) AGU Monograph series, 166, 147-161.
- 401 Hsu, S.-K., Sibuet, J.-C. Shyu C.-T., 2001. Magnetic inversion in the East China Sea

- 402 and Okinawa Trough: tectonic implications, *Tectonophysics*, 333, 111-122.
- 403 Huh, C.A., Su, C.C., 1999. Sedimentation dynamics in the East China Sea elucidated
404 from ^{210}Pb , ^{137}Cs and $^{239,240}\text{Pu}$, *Mar. Geol.*, 160, 183-196.
- 405 Hutnak, M., Fisher, A.T., Harris, R., Stein, C., Wang, K., Spinelli, G., Schindler, M.,
406 Villnger, H., Silver, E., 2008. Large heat and fluid fluxes driven through mid-plate
407 outcrops on ocean crust, *Nature Geoscience*, 1, 611-614.
- 408 Ishibashi, J., Urabe, T., 1995. Hydrothermal activity related to arc-backarc magmatism
409 in the western Pacific, *In Backarc Basins: Tectonics and Magmatism*, Ed. B Taylor,
410 Plenum Press, New York, 451-495.
- 411 Johnson, H.P., Hutnak, M., Dziak, R.P., Fox, C.G., Urcuyo, I., Cowen, J.P., Nabelek, J.,
412 Fisher, C., 2000. Earthquake-induced changes in a hydrothermal system on the Juan
413 de Fuca mid-ocean ridge, *Nature*, 407, 174-177.
- 414 Kawagucci, S., Chiba, H., Ishibashi, J., Yamanaka, T., Toki, T., Muramatsu, Y., Ueno,
415 Y., Makebe, A., Inoue, K., Yoshida, N., Nakagawa, S., Nunoura, T., Takai, K.,
416 Takahata, N., Sano, Y., Narita, T., Teranishi, G., Obata, H., and Gamo, T., 2011.
417 Hydrothermal fluid geochemistry at the Iheya North field in the mid-Okinawa
418 Trough: Implication for origin of methane in subseafloor fluid circulation systems,
419 *Geochem. J.*, 45, 109-124.

- 420 Konno, U., Tsunogai, U., Nakagawa, F., Nakaseama, M., Ishibashi, J., Nunoura, T.,
421 Nakamura, K., 2006. Liquid CO₂ venting on the seafloor: Yonaguni Knoll IV
422 hydrothermal system, Okinawa Trough, *Geophys. Res. Lett.*, 33, L16607,
423 doi:10.1029/2006GL026115.
- 424 Kumagai, H., Tsukioka, S., Yamamoto, H., Tsuji, T., Shitashima, K., Asada, M.,
425 Yamamoto, F., and Kinoshita, M., 2010. Hydrothermal plumes imaged by high -
426 resolution side - scan sonar on a cruising AUV, Urashima, *Geochem. Geophys.*
427 *Geosyst.*, 11, Q12013, doi:10.1029/2010GC003337.
- 428 Lee, C., Shor, G.G., Bibee, L.D., Lu, R.S., Hilde, T.W.C., 1980. Okinawa Trough,
429 origin of a back-arc basin, *Mar. Geol.*, 35, 219–241.
- 430 Letouzey, J., Kimura, M., 1986. Okinawa Trough: genesis of a back-arc basin
431 developing along a continental margin, *Tectonophysics*, 125, 209-230.
- 432 Lilley, M.D., Butterfield, D.A., Olson, E.J., Lupton, J.E., Macko, S.A., Mcduff, R.E.,
433 1993. Anomalous CH₄ and NH₄⁺ concentrations at an unsedimented
434 mid-ocean-ridge hydrothermal system, *Nature*, 364, 45-47.
- 435 Masaki, Y., Kinoshita, M., Inagaki, F., Nakagawa, S., Takai, K., 2011. Possible
436 kilometer-scale hydrothermal circulation within the Iheya-North field, mid-Okinawa
437 Trough, as inferred from heat flow data, *JAMSTEC Rep. Res. Dev.*, 12, 1-12

- 438 Momma, H., Iwase, R., Mitsuzawa, K., Kaiho, Y., Fujiwara, Y., Amitani, Y., Aoki, M.,
439 1996. Deep tow survey in Nanseisyoto region (K95-07-NSS), JAMSTEC J
440 Deep-sea Res., 12, 195-210 (in Japanese with English abstract).
- 441 Nakagawa, S., Takai, K., Inagaki, F., Chiba, H., Ishibashi, J., Kataoka, S., Hirayama, H.,
442 Nunoura, T., Horikoshi, K., Sako, Y., 2005. Variability in microbial community and
443 venting chemistry in a sediment-hosted backarc hydrothermal system: impacts of
444 subseafloor phase-separation, FEMS Microbiol. Ecol., 54, 141-155.
- 445 Nakamura, M., Otakia, K. and Takeuchi, S., 2008. Permeability and pore-connectivity
446 variation of pumices from a single pyroclastic flow eruption: Implications for partial
447 fragmentation, Journal of Volcanology and Geothermal Research, 176, 302-314.
- 448 Narita, H., Harada, K., Tsunogai, S., 1990. Lateral transport of sediment in particles in
449 the Okinawa Trough determined by natural radionuclides, Geochem J., 24, 207-216.
- 450 Oiwane, H., Kumagai, H., Masaki, Y., Nakamura, Y., Tokuyama, H., 2008.
451 Characteristics of Sediment in Iheya North Knoll and 'the acoustic blanking layer,
452 JPGU meeting, J164-P013 (English abstract).
- 453 Pearson, A., Seewald, J.S., Eglinton, T.I., 2005. Bacterial incorporation of relict carbon
454 in the hydrothermal environment of Guaymas Basin, Geochim. Cosmochim. Acta,
455 69, 5477-5486.

- 456 Shanks W.C., Bohlke, J.K., Seal, R.R., 1995. Stable isotopes in Mid-Ocean Ridge
457 hydrothermal systems: interactions between fluids, minerals and organisms, p.
458 194-221. *In* S.E. Humphris, R.A. Zierenberg, L.S. Mullineaux, and R.E. Thompson
459 (eds.), Geophysical Monograph 91 Seafloor Hydrothermal System, American
460 Geographical Union, Washington DC.
- 461 Sheriff, R.E., 2002. Encyclopedic Dictionary of Exploration Geophysics, Geophys. Ref.
462 Ser., vol. 13, 4th ed., Soc. of Explor. Geophys., Tulsa, Okla.
- 463 Sibuet, J.-C., Hsu, S.-K., Shyu, C.-T., Liu, C.-S., 1995. Structural and kinematic
464 evolution of the Okinawa trough backarc basin. In: Taylor, B. (Ed.), Backarc
465 Basins: Tectonics and Magmatism. Plenum, New York, pp. 343– 378.
- 466 Stein, C., Stein, S., 1992. A model for the global variation in oceanic depth and heat
467 flow with lithospheric age, *Nature*, 359, 123–129.
- 468 Takai, K., and Nakamura, K., 2010. Compositional, Physiological and Metabolic
469 Variability in Microbial Communities Associated with Geochemically Diverse,
470 Deep-Sea Hydrothermal Vent Fluids, L.L. Barton et al. (eds.), *Geomicrobiology:
471 Molecular and Environmental Perspective*, Springer Netherlands.
- 472 Takai, K., Nakagawa, S., Reysenbach, A.-L., Hoek, J., 2006. Microbial ecology of
473 Mid-Ocean Ridges and Back-Arc Basins. *In* "Back-Arc Spreading Systems:

- 474 Geological, Biological, Chemical, and Physical Interactions" (D. M. Christie, C. R.
475 Fisher, S.-M. Lee and S. Givens ed.) AGU Monograph series, 166, 185-213.
- 476 Taner, M. T., Koehler, F., Sheriff, R.E., 1979. Complex seismic traces analysis,
477 Geophysics, 44, 1041– 1063.
- 478 Tsugaru, R., Tamaki, Y., Sato, M., Kimura, M., Shimamura, K., 1991. Report on DELP
479 1988 cruises in the Okinawa Trough: part4, Preliminary report on the sediment
480 samples, Bull Earthq. Res. Inst., 66, 71-89.
- 481 Tsuji, T., Matsuoka, T., Yamada, Y., Nakamura, Y., Ashi, J., Tokuyama, H., Kuramoto,
482 S., Bangs, N., 2005. Initiation of plate boundary slip in the Nankai Trough off the
483 Muroto peninsula, southwest Japan, Geophys. Res. Lett., 32, L12306,
484 doi:10.1029/2004GL021861.
- 485 Tsuji, T., Nakamura, Y., Tokuyama, H., Coffin, M.F., Koda, K., 2007. Oceanic crust and
486 Moho of the Pacific Plate in the eastern Ogasawara Plateau region, Island Arc, 16,
487 361-373.
- 488 Waldhauser, F., and Tolstoy, M., 2011. Seismogenic structure and processes associated
489 with magma inflation and hydrothermal circulation beneath the East Pacific Rise at
490 9°50' N, Geochem. Geophys. Geosyst., 12, Q08T10, doi:10.1029/2011GC003568.
- 491 Welhan, J. A., Lupton, J.E., 1987. Light hydrocarbon gases in Guaymas Basin

hydrothermal fluids: Thermogenic versus abiogenic origin, Amer. Assoc. Petrol.

Geol. Bull., 71, 215-223.

Wheat, C.G., Jannasch, H.W., Kastner, M., Plant, J.N., DeCarlo, E.H., 2003. Seawater

transport and reaction in upper oceanic basaltic basement: chemical data from

continuous monitoring of sealed boreholes in a mid-ocean ridge flank environment,

Earth Planet. Sci. Lett., 216, 549–564.

Wheat, C.G., Jannasch, H.W., Kastner, M., Plant, J.N., DeCarlo, E.H., Lebon, G., 2004.

Venting formation fluids from deep sea boreholes in a ridge flank setting: ODP Sites

1025 and 1026, Geochem. Geophys. Geosyst., 5, Q08007,

doi:10.1029/2004GC000710.

Yilmaz, O., 2001. Seismic Data Processing: Processing, Inversion, and Interpretation of

Seismic Data, Invest. Geophys. Ser., vol. 10, 2nd ed., edited by S. M. Doherty, Soc.

of Explor. Geophys., Tulsa, Okla.

Table 1. Acquisition parameters for seismic surveys

Cruise	Vessel	Survey lines	Streamer	Source
KY02-11	<i>R/V Kaiyo</i>	2 lines	24-channel, 600 m streamer	~2.5 L (150 inch ³) GI airgun
YK06-09	<i>R/V Yokosuka</i>	26 lines	1-channel Streamer	~2.5 L (150 inch ³) GI airgun
KY07-03	<i>R/V Kaiyo</i>	19 lines	48-channel, 1200 m streamer	~5.8 L (355 inch ³) GI airgun
KR10-02	<i>R/V Kairei</i>	7 lines	360-channel 4500 m streamer	~54 L (3300 inch ³) Tuned airgun

Figure Captions




Figure 1. (a) Location map and bathymetry of the Okinawa Trough. (b) Bathymetric map of the Iheya North Knoll area showing seismic survey lines used in this study. The yellow star shows the location of the hydrothermal field. The seismic profiles displayed in Figures 3, 4, and 7 are shown as thick red lines. (c) Enlarged bathymetric map of the Iheya North Knoll area with seismic survey lines shown. The seismic profiles displayed in Figures 5 and 7 are shown as thick red lines.






Figure 2. Bathymetric map of the central valley within the Iheya North Knoll acquired by AUV *Urashima* during cruise YK07-07. White arrows indicate hydrothermal mounds. The volcanic ridge in the central valley and the peak on the western side of the hydrothermal field are also shown.

Hydrothermal fluid paths in continental margin



540 **Figure 3.** Prestack depth migrated seismic profiles across the Iheya North Knoll
541 hydrothermal field with and without geological interpretation. (a) E–W line
542 KR1002-18 and (b) N–S line KR1002-E2. Locations of the profiles are shown in
543 Figure 1b. The two profiles intersect at the Iheya North Knoll hydrothermal field
544 (red arrow). Gray shaded areas indicate volcanic bodies or deposits with high
545 permeability. Gray lines indicate geological boundaries, some of which are
546 pumiceous deposits. The normal faults indicated by blue lines.

Hydrothermal fluid paths in continental margin



548 **Figure 4.** N–S prestack depth migrated seismic profiles (with and without
549 geological interpretation) west of the Iheya North Knoll, showing intense normal
550 faulting. (a) Line KR1002-A02 and (b) line KR1002-A01. Locations of the profiles
551 are shown in Figure 1b. Normal faults (blue lines) may provide fluid recharge
552 paths to the deep crust. Red arrows indicate the location of a possible magma
553 chamber.

554

555




Figure 5. Interpreted time-domain seismic reflection profiles around the Iheya North Knoll hydrothermal field. Locations of the profiles are shown in Figure 1c. No amplitude gain control (AGC) was applied during processing of these sections. (a) E–W profile KY0703-18 and (d) N–S profile KY0703-E. Vertical white dashed lines indicate the intersection of the two profiles. Panels (b) and (e) show enlargements from profiles KY0703-18 and KY0703-E, respectively. Black circles indicate negative-polarity reflections. Gray circles indicate reflections

564 parallel to the negative-polarity reflections. Black diamonds indicate the upper
565 surface of pyroclastic flow deposits. Gray diamonds indicate the base of
566 pyroclastic flow deposits. (c) Enlargement of panel (b) showing a seafloor
567 reflection wavelet and a negative-polarity reflection. In panels (a) and (b), blue
568 lines represent faults.

569

570

571

572




Figure 6. Fence diagram showing a 3D view of the seismic profiles and the geometry of the negative-polarity reflector and faults within trough-fill sediments. The shallowest part of the subseafloor negative-polarity reflection (top of low-velocity sequence) corresponds to the location of the hydrothermal field (white arrow). The faults displayed here were interpreted from intersecting orthogonal 2D profiles.




Figure 7. (a) N–S seismic profile (line YK0609-G2) crossing the central valley

east of the hydrothermal field showing volcanic pumice deposits (upper), and
amplitude envelope profile (lower) used to identify volcanic deposits within the
sedimentary sequence. Locations of the seismic profiles are shown in Figure 1c.
(b) Amplitude envelope profile of line KY0703-18. This profile is an enlargement
from Figure 5a showing interpreted volcanic deposits (white dashed lines).
Volcanic rocks within the sedimentary sequence are represented by
high-amplitude, low-frequency reflections. (c) N–S seismic profile west of the
Iheya North Knoll and amplitude envelope profile with interpreted volcanic
deposits (white dashed lines).

Hydrothermal fluid paths in continental margin




Figure 8. Model of the hydrothermal fluid system around the Iheya North Knoll (based on seismic profile KR1002-18; Fig. 3a). Fluids flow laterally and updip along permeable pumiceous layers within the sedimentary sequence and laterally along basement faults within the crust. Near the knoll, fluids migrate upward due to heating and are then trapped below the hydrogeologic barrier (red line) and flow beneath it to vents in the hydrothermal field.