

S1

A Short, Truncated, and Partial History of Chaos

Philip Holmes¹

¹*Program in Applied and Computational Mathematics, Department of Mechanical and Aerospace Engineering, and Neuroscience Institute, Princeton University, Princeton, NJ 08544-1000, USA, pholmes@math.princeton.edu*

Abstract

Starting with the story of Henri Poincaré's correction of a serious mistake in his prize-winning paper on the three-body problem of classical mechanics [1] and his subsequent discovery of deterministic chaos in the form of homoclinic tangles (cf. [2, Volume 3]), I shall sketch some key ideas and developments in dynamical systems theory. I shall emphasize problems in periodically-forced nonlinear oscillators, including the work of van der Pol and van den Mark [3], G.D. Birkhoff [4, 5, 6], M.L. Cartright and J.E. Littlewood [7], and N. Levinson [8], culminating in S. Smale's construction of the "horseshoe" in 1959-60 [9, 10]. I will also explain how V.K. Melnikov and V.I. Arnold [11, 12] provided rather general perturbative methods that provide proofs of the existence of homoclinic tangles such as those recognized by Poincaré and Smale.

It is perhaps no coincidence that I truncate this history shortly after the independent discoveries of chaos and strange attractors in 1961 by Y. Ueda [13] and E.N. Lorenz [14, 15]: discoveries that did not immediately attract the attention of mathematicians, although in [16], Ueda and his colleagues provided elegant illustrations of homoclinic tangles. Moreover, stopping in the 1960's allows me to escape submersion in the tsunami of activity, papers and books that followed this period, and gives me time to speculate on the rôles that chance, conservatism, publication delays, and general ignorance of work in other fields played in this history, and (along with fashionable fads) the rôles that they still play on the development of science. If time permits, I will also make some remarks on the influence of "chaos theory" on the study of turbulence in fluids.

Extended versions of parts of this story and comments on more recent developments can be found in [17, 18]. For a more detailed account of Poincaré's work on dynamics, see [19], and for reflections on the sociological and cultural context of nonlinear dynamics and chaos, see [20].

Acknowledgments

The past 40 years of work that led to this presentation were indulged and partially supported by the Universities of Southampton (UK), Cornell and Princeton, and by many federal agencies including the late Science and Engineering Research Council of the UK, and the National Science Foundation, the Office of Naval Research, the Department of Energy, the Air Force Office of Scientific Research, and the National Institutes of Health of the USA. I thank these institutions and agencies, and the many colleagues and students who have taught me more than I now remember.

References

- [1] H.J. Poincaré, Sur le problème des trois corps et les équations de la dynamique, *Acta Mathematica*, vol. 13, pp. 1–270, 1890.

- [2] H.J. Poincaré, *Les méthodes nouvelles de la mécanique celeste, Vols 1-3*, Gauthiers - Villars, Paris, 1892, 1893, 1899. [English translation edited by D. Goroff, published by the American Institute of Physics, New York, 1993.]
- [3] B. van der Pol and B. van der Mark, Frequency demultiplication, *Nature*, vol. 120, pp. 363–364, 1927.
- [4] G.D. Birkhoff, *Dynamical Systems*, American Mathematical Society, Providence, RI, 1927. [reprinted with an introduction by J. Moser and a preface by M. Morse, 1966]
- [5] G.D. Birkhoff, Sur quelques courbes fermées remarquables, *Bull. de la Soc Mathém de France*, vol. 60, pp 1-26, 1932.
- [6] G.D. Birkhoff, Sur l'existence de région d'instabilité en Dynamique, *Annales de l'institut Henri Poincaré*, vol 2, pp.369-386, 1932.
- [7] M.L. Cartwright and J.E. Littlewood, On nonlinear differential equations of the second order, I: the equation $\ddot{y} + k(1 - y^2)\dot{y} + y = b\lambda k \cos(\lambda t + a)$, k large, *J. London Math. Soc.*, vol. 20, pp. 180–189, 1945.
- [8] N. Levinson, A second-order differential equation with singular solutions, *Annals of Mathematics*, vol. 50, pp. 127–153, 1949.
- [9] S. Smale, Diffeomorphisms with many periodic points. In *Differential and Combinatorial Topology: A Symposium in Honor of Marston Morse*, pp. 63-70, S.S. Cairns(ed), Princeton University Press, Princeton, NJ, 1965.
- [10] S. Smale, Differentiable Dynamical Systems, *Bull. Amer. Math. Soc.*, vol. 73, pp. 747–817, 1967.
- [11] V.K. Melnikov, On the stability of the center for time-periodic perturbations. *Trans. Moscow Math. Soc.*, vol. 12, pp. 1–57, 1963.
- [12] V.I. Arnold, Instability of dynamical systems with several degrees of freedom, *Soviet Mathematics Doklady*, vol. 5, pp. 342–355, 1964.
- [13] Y. Ueda, *The Road to Chaos-II*, Aerial Press, Inc., 2001.
- [14] E.N. Lorenz, Deterministic non-periodic flow, *J. Atmospheric Sci.*, vol. 20, pp. 130–141, 1963.
- [15] E.N. Lorenz, *The Essence of Chaos*, University of Washington Press, Seattle, WA, 1993.
- [16] C. Hayashi, Y. Ueda, N. Akamatsu, and H. Itakura, On the behavior of self-oscillatory systems with external force, *Trans. IECE*, vol.53-A, no. 3, pp. 150-158, 1970. [Reprinted in [13].]
- [17] F. Diacu and P. Holmes, *Celestial Encounters: The Origins of Chaos and Stability*, Princeton University Press, Princeton, NJ, 1996.
- [18] P. Holmes, Ninety plus thirty years of nonlinear dynamics: More is different and less is more, *Int. J. Bifurcation and Chaos*, vol. 15, pp. 2703–2716, 2005.
- [19] J. Barrow-Green, *Poincaré and the Three Body Problem*, American Mathematical Society, Providence, RI, 1997.
- [20] D. Aubin and A. Dahan Dalmedico. Writing the history of dynamical systems and chaos: Longue durée and revolution, disciplines and cultures. *Historia Mathematica*, 29, 273–339, 2002.