

A method for estimating colony size using queen fecundity in termites under field conditions

Authors: Mamoru Takata^{1*}, Kiyotaka Yabe¹, Takuya Noro¹, Shun Mizote¹, Takao Konishi¹, Eisuke Tasaki^{1,2}, Kenji Matsuura¹

Author affiliation

¹Laboratory of Insect Ecology, Graduate School of Agriculture, Kyoto University
Kitashirakawa-Oiwake-cho, Sakyo-ku, Kyoto, 606-8502, Japan

²Present address: Department of Biology, Faculty of Science, Niigata University, 8050 Ikarashi 2-no-cho, Nishi-ku, Niigata 950-2181, Japan.

*Corresponding author: Mamoru Takata (E-mail: takata.mamoru.7z@kyoto-u.ac.jp)

16-digit ORCID: 0000-0002-8181-9987

Abstract

Colony size in social insects is one of the most important factors in shaping their self-organized system. It affects a wide variety of traits such as foraging and defense strategies, social immune responses, the degree of polymorphism, and reproductive output. However, colony size estimation of subterranean termites in the field has been challenging, due to their extremely cryptic biology and multiple site-nesting behavior. Since natural selection favors workers that maximize the number of their siblings, the amount of egg production may reflect the number of workers in the colony. Here, we report a method for inferring colony size in the field using total egg production in each colony from a subterranean termite, *Reticulitermes speratus*. Our investigation of field colonies revealed that the body weight of queens reaches a peak and had the largest variance in June and July and accurately predicts the number of eggs laid by the queen per 24 h. Using laboratory-reared colonies, we found that the total egg production in each colony is proportional to the number of workers. We also estimated the colony size of 198 field colonies and found that the median and maximum colony size was 24,500 and 451,800 workers per colony. The method for inferring colony size presented here may also be applicable to termite species with a clear seasonality in egg production. The colony size estimate will contribute to understanding the life history strategies and social systems of termites.

Keywords: colony size, population estimate, termites, social insects

33 **Introduction**

34 Division of labor between reproductive and non-reproductive castes is fundamental to the ecological
35 success of social insects (Wilson 1971; Oster and Wilson 1978; Beshers and Fewell 2001; Hölldobler
36 and Wilson 2009). Societies of social insects can be considered a superorganism, with the number of
37 non-reproductive individuals in a colony (i.e., colony size) in eusocial species corresponding to the
38 body size in solitary organisms (Bourke 1999; Dornhaus et al. 2012; Kennedy et al. 2017; Boomsma
39 and Gawne 2018). The colony size is an important social characteristic shaping defense and foraging
40 strategies, task specialization, the initiation time of the production of fertile dispersers (i.e., alates),
41 and the lifelong reproduction of alates (Wilson 1971; Thorne 1997; Su 2003; Hölldobler and Wilson
42 2009). To comprehensively understand the regulatory mechanisms and dynamics of life history traits
43 in social insects, it is essential to conduct researches that take into account colony size under natural
44 conditions.

45 In termites, especially subterranean species, investigation of the colony size in the field can
46 be challenging due to multiple-site nesting behavior and a cryptic biology. Several methods for
47 investigating the colony size have been proposed, including destructive sampling by collecting
48 termite nests and habitats (King and Spink 1969; Howard et al. 1982; Darlington 1990) or trapping
49 (Haverty et al. 1975; La Fage et al. 1983), and non-destructive methods such as using radioisotopes
50 (Spragg and Paton 1980) and mark–release–recapture protocols (Su and Scheffrahn 1988; Su et al.
51 1993; Su 1994; Kenneth Grace et al. 1995; Tsunoda et al. 1999; Su and Lee 2008). The mark–
52 recapture methods have been widely used for the estimation of population size in field studies;
53 however, since subterranean termites have large colony sizes and extensive foraging systems, it is
54 difficult to accurately estimate the whole population size by these methods (Thorne et al. 1996; Evans
55 et al. 1999). Although some successful colony size estimation methods have been proposed for
56 laboratory-kept colonies (Su 2013; Patel et al. 2020), to date, no method is suitable for estimating the
57 whole population size in the field termites.

58 In this study, we developed a method for estimating colony size under field conditions in a
59 subterranean termite *Reticulitermes speratus*. They live mostly in pine and Japanese cedar forests in
60 Japan (Takematsu 1999; Park et al. 2006). A single colony uses multiple logs which are connected by
61 underground tunnels (Abe 1987; Shellman-Reeve 1997). Mature colonies are typically headed by
62 one primary king (PK) and multiple secondary queens (SQs), which are produced by parthenogenesis
63 and eventually replace the primary queen (Matsuura et al. 2009, 2018). Egg production is a seasonal
64 event and is limited from June to August (Matsuura et al. 2007; Nozaki and Matsuura 2021; Takata

et al. 2023a). The resources required for egg production are produced by workers and predominantly consumed by queens (Konishi et al. 2023). We initially validated the mark–recapture method in *R. speratus*; however, the method was not applicable for estimating the whole colony size. We then developed a new method for estimating colony size based on the fecundity of the queens. A previous study reported that the body weight of a queen strongly correlates with the number of workers captured in and around the nests in *Macrotermes subhyalinus* (Darlington 1990). Although workers who were away from the nest were not taken into account in the study, there could be a correlation between the queen's fecundity and the total number of individuals in the colony. Thus, the objective of the present study was to determine whether the fecundity of queens could be used to estimate the number of workers in the field colonies (Fig. 1). First, we investigated the seasons with the greatest differences in fecundity of the queens among colonies. Second, we measured the relationship between the body weight of the queen and her fecundity during those seasons. Third, we evaluated the relationship between daily egg production and colony size under laboratory conditions also during those seasons. Finally, the number of workers was estimated in the field colonies.

Materials and methods

Termite sampling

Decayed logs containing a primary king and secondary queens of *R. speratus* were collected in pine or Japanese cedar forests in Kyoto, Osaka, Nara, Shiga, Hyogo, and Fukui, Japan, from 2017 to 2022 (see Dataset 1 for details). Six colonies were used for a mark–release–recapture program. Another 342 colonies were used to investigate the seasonal body weight of queens. To investigate the relationship between the body and ovaries weight of the queen, 25 out of the 342 colonies were used. To investigate the relationship between the body weight of the queen and her daily egg production, 18 out of the 25 colonies were used. Fifteen colonies, including 13 out of the 342 colonies used to investigate the queen weight and an additional 2 colonies, were used to investigate the presence or absence of satellite nests. An additional 15 colonies were used to evaluate the relationship between daily egg production and colony size. To estimate the number of workers in field colonies, 188 out of the 342 colonies were used. In every experiment, each colony was individually processed.

Evaluation of mark–recapture methods

To investigate whether the assumption of the mark–recapture methods (i.e., the workers distribute equally in the colony) are met in *R. speratus*, the proportion of marked workers was compared

between different colony areas under laboratory conditions. Parts of the log containing and not containing royals were collected for each of the six colonies. The colonies were individually transferred into a container ($70 \times 50 \times 40$ cm: $W \times D \times H$) lined with 1 cm of moistened mountain soil (Takagi Kenzai Co., Ltd., Japan), and maintained at 25°C under dark conditions. Six months later, 500 workers were extracted from the edge of each log, and marked by sprays (Tamiya Color TS-15 and TS-35). Different colors were applied to workers from each log (royal chamber/foraging area: green/blue or blue/green). Then, the workers were returned to their original logs. The logs were returned to the original position in the container and maintained at 25°C under dark conditions. Two weeks later, all termites were extracted from each log, and the number of marked and unmarked workers was recorded in three colonies. Four weeks later, the remaining three colonies were processed in the same manner.

Relationship between queen weight and daily egg production in field colonies

To determine the month with the largest variance in the body weight of queens, 342 colonies with kings and queens were collected during the breeding season (from April to September). Within 3 days of collection, all termites were extracted from the log, and the fresh body weight of queens was recorded to the nearest 0.1 mg.

Since the body weight of queens had the largest variance in June and July, 25 colonies were collected in June and July to evaluate the relationships between the queen's body weight and her fecundity in the field colonies. Within 24 h of collection, all termites were extracted and the fresh body weight of queens was recorded to the nearest 0.1 mg. To evaluate the relationships between the body weight of queens and ovary size, one SQ from each of the colonies was individually dissected in PBS buffer (Wako Pure Chemical Industry Co., Ltd., Japan) within 24 h of collection. Their ovaries were removed and transferred into a centrifuge tube to avoid desiccation after the removal of water on its surface. The fresh weight of their ovaries was recorded to the nearest 0.01 mg. Twenty-five SQs were used in total.

To evaluate the relationships between the body weight of queens and egg production, another two SQs were randomly selected from each of 18 out of the 25 colonies, individually transferred into dishes (ca. 30 mm) lined with a moist unwoven cloth and 10 non-nestmate workers (including both sexes), and maintained at 25°C under dark conditions. Thirty-six SQs were used in total. After 24 h, the number of eggs laid by SQs was recorded.

Relationship between queen weight and colony size in laboratory conditions

To evaluate the relationship between daily egg production and colony size, decayed logs of Japanese cedar containing kings and secondary queens were collected in April ($n = 15$ colonies). The colonies were individually transferred into a container ($626 \times 426 \times 365$ mm: $W \times D \times H$) keeping them separate from one another. The individual containers were then lined with 4 cm of moistened mountain soil. If the logs were too large to fit in a single container, they were split into multiple containers. Then an opening was made on the lower side of the containers connected by 15-cm tubes (6 mm inner diameter), so the workers could move between the containers. A moistened Japanese cedar log (ca. diameter 10 cm, length 35 cm) was added to each container as an additional food source. Then, the logs were maintained at ambient temperature. In June and early July, all termites were extracted from the container. The number of individuals who died during the extraction process was recorded by counting the number of individuals (or more precisely, the number of heads) directly for each caste. To calculate the ratio of the surviving individuals in each caste, all surviving termites were separated from their nest material and the fresh body weight of king and queens was recorded to the nearest 0.1 mg. The total weight of the remaining colony members was recorded to the nearest 1 mg. Then, a 500-mg subset of termite colony members was randomly selected and the numbers of each individual caste was recorded (worker, soldier, nymph, or larva). The number of surviving caste members was determined by the ratio of individuals in each caste from the randomly selected subset. The total number of members in each caste was determined by adding the calculated number of the surviving individuals to the counted number of the dead individuals.

Colony size estimation in the field colonies

Colony size estimation using queen fecundity requires the collection of all queens in the colony. Therefore, before estimating colony size in the field colonies, we confirmed the absence of satellite nests, which are apart from the location where the king is present and consist only of the queen and non-reproductive members. First, we searched for a log with one or more queens in the field. Once the queens were found, we brought the entire log back to the laboratory for further examination ($n = 15$ colonies). Within eight days of collection, we examined whether there was a king in the logs.

To estimate the number of workers in field colonies, 188 colonies with kings and queens were collected in June and July. Within 3 days of collection, all secondary queens were extracted, and their fresh body weight was recorded to the nearest 0.1 mg. The daily egg production in each colony was calculated using a regression formula between the queen's body weight and her daily egg production.

Then, the number of workers in the colony was estimated using a regression formula between daily egg production and the number of workers in the colony.

Statistical analysis

To investigate whether the workers distribute randomly in the colony as assumed in mark–recapture–methods, the proportion of marked workers was compared using a generalized linear mixed model (GLMM) with a binomial error distribution and logit link function. The response variable was the proportion of marked workers, and the explanatory variables were the log (whether the marked workers were introduced in the log or not), test period, and their interactions. Colony was included as a random factor. The proportion of marked workers was compared in each test period using the same model without the interaction term. Two-tailed paired *F*-tests were used to compare the variance in the mean body weight of the queens in each colony among different months. A two-sided Bonferroni-corrected significance level of $P < 0.0033$ was set. We analyzed the relationship between the body weight of queens and the size of their sexual organs using a generalized linear model (GLM) with a gamma error distribution and log link function. The response variable was the weight of the sexual organ, and the explanatory variable was the body weight of the queens. We also analyzed the relationship between the body weight of the queens and egg production using a GLM with a Poisson error distribution and log link function. The response variable was the number of eggs laid, and explanatory variable was the log-transformed the body weight of the queens. We analyzed the correlation between egg production and colony size using a GLM with a Poisson error distribution and log link function. The response variable was the number of workers, and explanatory variable was the log-transformed the daily egg production in each colony. We also analyzed the effect of the body weight of primary king, mean weight of secondary queen, number of the queens, or total weight of the queens on the colony size using a GLM with a Poisson error distribution and log link function. The response variable was the number of workers, and explanatory variable was the log-transformed the body weight of primary king, mean weight of secondary queen, number of the queens, or total weight of the queens in each colony.

All analyses were performed by the software R v4.2.3 (R Core Team 2022), with the lme4 and car packages. For the GLMMs and GLMs, likelihood ratio tests (LRTs) were used to determine the statistical significance of each explanatory variable. A significance value of $p < 0.05$ was considered to indicate statistical significance.

193 **Results**

194 **Distribution of marked workers**

195 The marked workers were not distributed equally in the colony but were found at a higher frequency
 196 from the logs in which they were introduced (GLMM, LRT: $\chi^2 = 13.537$, $df = 1$, $p < 0.001$, Fig. 2).
 197 The interaction between the log (whether the marked workers were introduced in the log or not) and
 198 the test period was not significant (GLMM, LRT: $\chi^2 = 0.044$, $df = 1$, $p = 0.834$). There were
 199 statistically significant differences in the proportion of marked workers between the logs in the 2-
 200 week (GLMM, LRT: $\chi^2 = 9.571$, $df = 1$, $p = 0.002$, Fig. 2a) and 4-week test period group (GLMM,
 201 LRT: $\chi^2 = 4.010$, $df = 1$, $p = 0.045$, Fig.2b).

203 **Relationship between queen weight and daily egg production in field colonies**

204 In total, 15,637 SQs from 342 colonies were collected from the field from April to September. We
 205 found that the body weight of the SQs begins to increase in May, peaks in June, and then decreases
 206 through September (Fig. 3a). The variance in mean body weight of the queens among the colonies
 207 was the largest in June and July (F -test with Bonferroni correction). In June and July, the body weight
 208 of queens correlated with their ovaries weight (GLM, LRT: $\chi^2 = 610.89$, $df = 1$, $p < 0.001$, $R^2 = 0.951$,
 209 Fig. 3b) and the number of eggs produced in 24 hours (GLM, LRT: $\chi^2 = 805.36$, $df = 1$, $p <$
 210 0.001 , $R^2 = 0.914$, Fig. 3c). The regression formula between body weight of queens and their daily
 211 egg production is as follows:

$$213 \quad y_i = e^{(-1.052582 + 1.855425 * \log(x_i))} \quad (1)$$

214
 215 where y_i is the number of eggs produced in 24 h by the queen i and x_i is the body weight of the queen
 216 i .

218 **Relationship between queen weight and colony size in laboratory conditions**

219 Using laboratory-reared colonies, we found that colony size is predictable from the body weight of
 220 SQs in the colony. The total egg production in each colony was calculated from the regression formula
 221 (1), and was best explained the variation in the number of workers in a colony (GLM, LRT: $\chi^2 =$
 222 $1,571,616$, $df = 1$, $p < 0.001$, $AIC = 36,630$, $R^2 = 0.975$; null model: $AIC = 1,608,244$, Fig. 4). The
 223 number of workers was weakly correlated with the weight of primary king (GLM, LRT: $\chi^2 = 887,905$,
 224 $df = 1$, $p < 0.001$, $AIC = 720,341$, $R^2 = 0.645$), the mean weight of the secondary queen (GLM,

LRT: $\chi^2 = 909,198$, $df = 1$, $p < 0.001$, $AIC = 699,048$, $R^2 = 0.310$), the number of queens (GLM, LRT: $\chi^2 = 847,66$, $df = 1$, $p < 0.001$, $AIC = 1,523,480$, $R^2 = 0.034$), and the total weight of the queens (GLM, LRT: $\chi^2 = 1,265,010$, $df = 1$, $p < 0.001$, $AIC = 343,236$, $R^2 = 0.820$). The regression formula between the total daily egg production and the number of workers is as follows:

229

$$w = e^{(0.8946205 + 1.411185 * \log(\sum y_i))} \quad (2)$$

231

232 where w is the number of workers in the colony and $\sum y_i$ is the total number of eggs produced in
233 the colony in 24 h.

234

235 **Estimated colony size in the field colonies**

236 We confirmed that satellite nests are either absent or extremely rare. In all 15 logs where queens were
237 found, kings were also collected in each of them (see Dataset for details). The colony sizes were
238 calculated using the regression formulas (1) and (2) in 188 field colonies collected in June and July.
239 The first quartile, median, third quartile, and maximum of the estimated number of workers in a
240 colony were 10,700, 24,500, 58,600, and 451,800, respectively (Fig. 5).

241

242 **Discussion**

243 We present a reliable method for estimating colony size in field colonies of *R. speratus*. We initially
244 evaluated a mark–recapture method commonly used in estimating colony size for subterranean
245 termite species (Grace et al. 1989; Su et al. 1993; Tsunoda et al. 1999). However, we found that the
246 assumption of the mark–recapture method is not met in *R. speratus*, similar to other species of
247 subterranean termites (Thorne et al. 1996; Evans et al. 1999). Our results showed that marked workers
248 are not equally distributed throughout the colony, and the proportion of marked workers varies even
249 between neighboring logs (Fig. 2). The mark–recapture methods requires an assumption of
250 equilibrium (Su 2013), which is not met in *R. speratus* as presented here (see also Dataset) and
251 *Coptotermes formosanus* (Su et al. 2017), as the different instars are distributed unequally over the
252 distance from the central part of the nest. The actual degree of the relative lack of equilibrium may
253 provide the data necessary to estimate population size more accurately. Although we do not
254 completely reject the potential use of mark–recapture methods in *R. speratus*, we decided to consider
255 an alternative approach. As a result, we found that the number of workers in the colony can be
256 estimated from the body weight of the queens. The weight of the queen reflects the number of eggs

laid per 24 h (Fig. 3), and the total number of eggs produced per unit of time in the colony is strongly correlated with the total number of workers (Fig. 4). The queens of this species are distributed in a single log and does not exist across multiple sites. Therefore, by collecting all the queens from a single log we can estimate the total egg production and the number of workers in the whole colony.

Our colony size estimation is based on the valid assumption that termite colonies are adapted to maximize egg production based on colony size within the constraints of a limited growing season. While termites are typically found in tropical regions, some species, such as *R. speratus*, have successfully adapted to and colonized temperate zones (Emerson 1955; Eggleton et al. 1994; Takematsu 1999; Eggleton 2000; Park et al. 2006; Evans et al. 2013). *R. speratus* exhibits a clear seasonality in egg production, which is limited to the period from June to August (Nozaki and Matsuura 2021). The queens are specialized for egg production, and their fecundity is reflected in their body weight (Fig. 3). The queens rely on a food supply from the workers for the nutrients required for egg production (Tasaki et al. 2023; Konishi et al. 2023), and unlike the workers, they have no symbiotic microorganisms which are essential for wood digestion (Shimada et al. 2013; Inagaki and Matsuura 2016). Therefore, it is reasonable to infer that the body weight of queens can serve as an indicator of worker population size within a colony, as it reflects the number of workers available for food provisions. The present study demonstrated that the number of workers affects egg production. The annual production of new members influences age demography and is fundamental for the maintenance and health of termite societies (Chouvenc et al. 2022). Future comparative studies that examine the impact of seasonal egg production patterns on overall colony functionality will be valuable in revealing how termite colonies adapt to cyclical changes in environmental conditions that occur, for example, in temperate regions where temperature fluctuates more dramatically than tropical regions.

Our method, which requires the dissection of the royal chamber to collect all the queens, is not suitable for monitoring and follow-up studies, but it allows for snapshot studies. The organization of colonial organisms is analogous to that of multicellular organisms, and colony size in eusocial species corresponds to body size in solitary species (Bourke 1999; Dornhaus et al. 2012; Kennedy et al. 2017; Boomsma and Gawne 2018). Colony size is one of the most important parameters that shape the collective organization of colonial organisms, as it influences defense and foraging strategies, task specialization, and the production of dispersers (Wilson 1971; Thorne 1997; Su 2003; Hölldobler and Wilson 2009). Furthermore, colony size may also play a crucial role in breeding systems and caste development, as seen with the replacement of primary queens with secondary queens, and changes

in caste development occurring during the colony life cycle in *R. speratus* (Matsuura et al. 2009, 2018; Takata et al. 2023b). Our method has broad applicability in studying the role and contribution of colony size in these life history events and traits in the field and opens new avenues for understanding the life history strategies and social systems of termites.

The data must be interpreted carefully because the relationship between queen weight and the number of workers may vary slightly in the field, and our data may underestimate the population size of field colonies. In this study, termite nesting logs were placed adjacent to each other, but in the field, *R. speratus* foraging territories can extend up to 56.6 m² per colony, with foraging occurring from logs as far as 10 m linear distance away (Tsunoda et al. 1999). Hence, additional studies are needed to investigate the potential influence of energy expenditure associated with nest-to-nest movement on egg production, as this would allow a more accurate estimation of population size.

Data availability

All data generated or analyzed during this study are included in this published article.

Code availability

The codes for graphing and statistical analyses used for this paper are available at GitHub (https://github.com/MamoruTakata/Colony_size_estimate).

References

- Abe T (1987) Evolution of life types in termites. In: Kawano S, Connell JH, Hidaka T (eds) Evolution and coadaptation in biotic communities. University of Tokyo Press, Tokyo, Japan, pp 125–148
- Beshers SN, Fewell JH (2001) Models of division of labor in social insects. *Annu Rev Entomol* 46:413–440. <https://doi.org/10.1146/annurev.ento.46.1.413>
- Boomsma JJ, Gawne R (2018) Superorganismality and caste differentiation as points of no return: how the major evolutionary transitions were lost in translation. *Biol Rev* 93:28–54. <https://doi.org/10.1111/brev.12330>
- Bourke AFG (1999) Colony size, social complexity and reproductive conflict in social insects. *J Evol Biol* 12:245–257. <https://doi.org/10.1046/j.1420-9101.1999.00028.x>
- Chouvenc T, Ban PM, Su N-Y (2022) Life and death of termite colonies, a decades-long age demography perspective. *Front Ecol Evol* 10:1–21.

- 321 <https://doi.org/10.3389/fevo.2022.911042>
- 322 Darlington JPEC (1990) Populations in nests of the termite *Macrotermes subhyalinus* in Kenya.
- 323 *Insectes Soc* 37:158–168. <https://doi.org/10.1007/BF02224028>
- 324 Dornhaus A, Powell S, Bengston S (2012) Group size and its effects on collective organization.
- 325 *Annu Rev Entomol* 57:123–141. <https://doi.org/10.1146/annurev-ento-120710-100604>
- 326 Eggleton P (2000) Global patterns of termite diversity. In: Abe T, Bignell DE, Higashi M (eds)
- 327 *Termites, Evolution, Sociality, Symbioses, Ecology*. Springer Netherlands, Dordrecht, pp
- 328 25–51
- 329 Eggleton P, Williams PH, Gaston KJ (1994) Explaining global termite diversity: productivity or
- 330 history? *Biodivers Conserv* 3:318–330. <https://doi.org/10.1007/BF00056505>
- 331 Emerson AE (1955) Geographic origins and dispersion of termite genera. *Fieldiana Zool* 37:465–
- 332 521
- 333 Evans TA, Forschler BT, Grace JK (2013) Biology of invasive termites: a worldwide review.
- 334 *Annu Rev Entomol* 58:455–474. <https://doi.org/10.1146/annurev-ento-120811-153554>
- 335 Evans TA, Lenz M, Gleeson P V. (1999) Estimating population size and forager movement in a
- 336 tropical subterranean termite (Isoptera: Rhinotermitidae). *Environ Entomol* 28:823–830.
- 337 <https://doi.org/10.1093/ee/28.5.823>
- 338 Grace JK, Abdallay A, Farr KR (1989) Eastern subterranean termite (Isoptera: Rhinotermitidae)
- 339 foraging territories and populations in Toronto. *Can Entomol* 121:551–556.
- 340 <https://doi.org/10.4039/Ent121551-7>
- 341 Haverty MI, Nutting WL, Lafage JP (1975) Density of colonies and spatial distribution of
- 342 foraging territories of the desert subterranean termite, *Heterotermes aureus* (Snyder).
- 343 *Environ Entomol* 4:105–109. <https://doi.org/10.1093/ee/4.1.105>
- 344 Hölldobler B, Wilson EO (2009) The superorganism: the beauty, elegance, and strangeness of
- 345 insect societies. W. W. Norton, New York
- 346 Howard RW, Jones SC, Mauldin JK, Beal RH (1982) Abundance, distribution, and colony size
- 347 estimates for *Reticulitermes* spp. (Isoptera: Rhinotermitidae) in southern Mississippi.
- 348 *Environ Entomol* 11:1290–1293. <https://doi.org/10.1093/ee/11.6.1290>
- 349 Inagaki T, Matsuura K (2016) Colony-dependent sex differences in protozoan communities of
- 350 the lower termite *Reticulitermes speratus* (Isoptera: Rhinotermitidae). *Ecol Res* 31:749–
- 351 755. <https://doi.org/10.1007/s11284-016-1387-2>
- 352 Kennedy P, Baron G, Qiu B, et al (2017) Deconstructing superorganisms and societies to address

- big questions in biology. *Trends Ecol Evol* 32:861–872.
<https://doi.org/10.1016/j.tree.2017.08.004>
- Kenneth Grace J, Yamamoto RT, Tamashiro M (1995) Relationship of individual worker mass and population decline in a Formosan subterranean termite colony (Isoptera: Rhinotermitidae). *Environ Entomol* 24:1258–1262. <https://doi.org/10.1093/ee/24.5.1258>
- King EG, Spink WT (1969) Foraging galleries of the Formosan subterranean termite, *Coptotermes formosanus*, in Louisiana. *Ann Entomol Soc Am* 62:536–542.
<https://doi.org/10.1093/aesa/62.3.536>
- Konishi T, Tasaki E, Takata M, Matsuura K (2023) King- and queen-specific degradation of uric acid contributes to reproduction in termites. *Proc R Soc B Biol Sci* 290:20221942.
<https://doi.org/10.1098/rspb.2022.1942>
- La Fage JP, Su N-Y, Jones MJ, Esenther GR (1983) A rapid method for collecting large numbers of subterranean termites from wood. *Sociobiology* 7:305–309
- Matsuura K, Kobayashi N, Yashiro T (2007) Seasonal patterns of egg production in field colonies of the termite *Reticulitermes speratus* (Isoptera: Rhinotermitidae). *Popul Ecol* 49:179–183. <https://doi.org/10.1007/s10144-006-0030-4>
- Matsuura K, Mizumoto N, Kobayashi K, et al (2018) A genomic imprinting model of termite caste determination: not genetic but epigenetic inheritance influences offspring caste fate. *Am Nat* 191:677–690. <https://doi.org/10.1086/697238>
- Matsuura K, Vargo EL, Kawatsu K, et al (2009) Queen succession through asexual reproduction in termites. *Science* 323:1687–1687. <https://doi.org/10.1126/science.1169702>
- Nozaki T, Matsuura K (2021) Oocyte resorption in termite queens: seasonal dynamics and controlling factors. *J Insect Physiol* 131:104242.
<https://doi.org/10.1016/j.jinsphys.2021.104242>
- Oster GF, Wilson EO (1978) *Caste and ecology in the social insects*. Princeton University Press, Princeton, NJ
- Park YC, Kitade O, Schwarz M, et al (2006) Intraspecific molecular phylogeny, genetic variation and phylogeography of *Reticulitermes speratus* (Isoptera: Rhinotermitidae). *Mol Cells* 21:89–103
- Patel JS, Lee S-B, Chouvenc T, Su N-Y (2020) Inferring termite colony size using wood consumption in subterranean termites (Blattodea: Rhinotermitidae) in laboratory-rearing conditions. *J Econ Entomol* 113:905–910. <https://doi.org/10.1093/jee/toz353>

- 385 R Core Team (2022) R: a language and environment for statistical computing. R Foundation for
386 Statistical Computing
- 387 Shellman-Reeve JS (1997) The spectrum of eusociality in termites. In: Choe JC, Crespi BJ (eds)
388 Evolution of social behavior in insects and arachnids. Cambridge University Press,
389 Cambridge, UK, pp 52–93
- 390 Shimada K, Lo N, Kitade O, et al (2013) Cellulolytic protist numbers rise and fall dramatically in
391 termite queens and kings during colony foundatio. *Eukaryot Cell* 12:545–550.
392 <https://doi.org/10.1128/EC.00286-12>
- 393 Spragg WT, Paton R (1980) Tracing, trophallaxis and population measurement of colonies of
394 subterranean termites (Isoptera) using a radioactive tracer. *Ann Entomol Soc Am* 73:708–
395 714. <https://doi.org/10.1093/aesa/73.6.708>
- 396 Su N-Y (2003) Behavioral ecology of foraging in subterranean termites. In: Bignell DE, Roisin
397 Y, Lo N (eds) *Biology of Termites: a Modern Synthesis*. Springer, Dordrecht, Netherlands,
398 pp 165–191
- 399 Su N-Y (1994) Field evaluation of a hexaflumuron bait for population suppression of
400 subterranean termites (Isoptera: Rhinotermitidae). *J Econ Entomol* 87:389–397.
401 <https://doi.org/10.1093/jee/87.2.389>
- 402 Su N-Y (2013) Estimating population size of large laboratory colonies of the Formosan
403 subterranean termite using the capture probability equilibrium. *J Econ Entomol* 106:2442–
404 2447. <https://doi.org/10.1603/EC13258>
- 405 Su N-Y, Ban PM, Scheffrahn RH (1993) Foraging populations and territories of the eastern
406 subterranean termite (Isoptera: Rhinotermitidae) in southeastern Florida. *Environ Entomol*
407 22:1113–1117. <https://doi.org/10.1093/ee/22.5.1113>
- 408 Su N-Y, Lee S-H (2008) Estimating the population size and colony boundary of subterranean
409 termites by using the density functions of directionally averaged capture probability. *J Econ*
410 *Entomol* 101:592–604. <https://doi.org/10.1093/jee/101.2.592>
- 411 Su N-Y, Osbrink W, Kakkar G, et al (2017) Foraging distance and population size of juvenile
412 colonies of the Formosan subterranean termite (Isoptera: Rhinotermitidae) in laboratory
413 extended arenas. *J Econ Entomol* 110:1728–1735. <https://doi.org/10.1093/jee/tox153>
- 414 Su N-Y, Scheffrahn RH (1988) Foraging population and territory of the Formosan subterranean
415 termite (Isoptera: Rhinotermitidae) in an urban environment. *Sociobiology* 14:353–359
- 416 Takata M, Konishi T, Nagai S, et al (2023a) Discovery of an underground chamber to protect

- 417 kings and queens during winter in temperate termites. *Sci Rep* 13:8809.
 418 <https://doi.org/10.1038/s41598-023-36035-1>
- 419 Takata M, Nagai S, Inagaki T, et al (2023b) Heritable effects on caste determination and colony-
 420 level sex allocation in termites under field conditions. *iScience* 26:106207.
 421 <https://doi.org/10.1016/j.isci.2023.106207>
- 422 Takematsu Y (1999) The genus *Reticulitermes* (Isoptera: Rhinotermitidae) in Japan, with
 423 description of a new species. *Entomol Sci* 2:231–243
- 424 Tasaki E, Mitaka Y, Takahashi Y, et al (2023) The royal food of termites shows king and queen
 425 specificity. *PNAS Nexus* Accepted: <https://doi.org/10.1093/pnasnexus/pgad222>
- 426 Thorne BL (1997) Evolution of eusociality in termites. *Annu Rev Ecol Syst* 28:27–54.
 427 <https://doi.org/10.1146/annurev.ecolsys.28.1.27>
- 428 Thorne BL, Russek-Cohen E, Forschler BT, et al (1996) Evaluation of mark–release–recapture
 429 methods for estimating forager population size of subterranean termite (Isoptera:
 430 Rhinotermitidae) colonies. *Environ Entomol* 25:938–951.
 431 <https://doi.org/10.1093/ee/25.5.938>
- 432 Tsunoda K, Matsuoka H, Yoshimura T, Tokoro M (1999) Foraging populations and territories of
 433 *Reticulitermes speratus* (Isoptera: Rhinotermitidae). *J Econ Entomol* 92:604–609.
 434 <https://doi.org/10.1093/jee/92.3.604>

435 Wilson EO (1971) *The insect societies*. Belknap Press, Cambridge, Massachusetts
 436

437 **Acknowledgements**

438 We thank Matthew Tatsuo Kamiyama for the fruitful discussion, and Wu Yao, Chihiro Tamaki,
 439 Takehiro Morimoto, Hiroki Noda, Shuya Nagai for assistance in collecting termites and helpful
 440 discussion.

441

442 **Funding**

443 This work was supported by JSPS KAKENHI Grant Numbers JP18H05268, JP20K20380, and
 444 JP23H00332 to K.M., JP21K14863 to M.T., and JP20J20278 to T.K.

445

446 **Author information**

447 **Authors and Affiliations**

448 Laboratory of Insect Ecology, Graduate School of Agriculture, Kyoto University, Kitashirakawa-

449 Oiwake-cho, Sakyo-ku, Kyoto, 606-8502, Japan

450 Mamoru Takata, Kiyotaka Yabe, Takuya Noro, Shun Mizote, Takao Konishi, Eisuke

451 Tasaki & Kenji Matsuura

452 Present address: Department of Biology, Faculty of Science, Niigata University, 8050 Ikarashi 2-

453 no-cho, Nishi-ku, Niigata, 950-2181, Japan

454 Eisuke Tasaki

455

456 **Contributions**

457 Conceptualization, M.T.; methodology, M.T. and K.Y.; validation, M.T.; formal analysis, M.T.;

458 investigation, M.T., K.Y., T.N. S.M., T.K., and E.T.; resources, M.T., K.Y., T.N. S.M., T.K., and

459 E.T.; writing—original draft: M.T.; writing—review and editing: M.T., K.Y., T.N. S.M., T.K.,

460 E.T., and K.M.; visualization, M.T. and T.K.; supervision, M.T. and K.M.; project

461 administration, M.T.; funding acquisition, M.T., T.K., and K.M.

462

463 **Corresponding author**

464 Correspondence to Mamoru Takata.

465

466 **Ethics declarations**

467 **Ethics approval**

468 Not applicable.

469

470 **Consent to participate**

471 Not applicable.

472

473 **Consent for publication**

474 Not applicable.

475

476 **Competing interests**

477 The authors declare no competing interests.

478

479 **Additional information**

480 Communicated by: Sean O'Donnell

481

482 **Supplementary information**


483 **ESM 1**

484 (XLSX 348 kb)

485

486 **Figures**

487




488

489 **Figure 1** Schematic representation of the nutritional contribution of workers on egg production in
 490 small (**a**) and large colonies (**b**). A single colony uses multiple logs which are connected by tunnels.
 491 Even if the workers forage from multiple logs, the nutrients necessary for egg production are
 492 concentrated in the queens, so the number of eggs produced is likely to reflect the number of workers
 493 in the colony. The size of the queens varies remarkably among colonies.

494

495



496


497

498

499


500

Figure 2 Comparison of the proportion of marked workers between the logs in which they were introduced and the other log in the two-week (a) and 4-week test period group (b). Asterisks indicate significant differences (likelihood ratio test, $*p < 0.05$, $**p < 0.01$).



502 **Figure 3** Body weight and fecundity of secondary queens in *Reticulitermes speratus*. **a** Seasonal
 503 changes in the variance in mean body weight of queens among colonies. Plots indicate the mean body
 504 weight of queens in each colony ($n = 342$ colonies). Different letters indicate significant differences
 505 in variance (F -test with Bonferroni correction, $p < 0.05$). **b** Correlation between queen weight and
 506 ovaries weight in field colonies. The fresh weights of the ovaries are plotted as a function of the body
 507 weight of queens ($n = 25$ secondary queens from 25 colonies). The solid line represents the estimated
 508 regression line, $y_i = e^{(-3.071019 + 1.754776 * \log(x_i))}$. The y_i and x_i are the ovaries and the body
 509 weight of the queen i , respectively. **c** Correlation between queen weight and daily egg production in
 510 field colonies. The number of eggs laid by queens is plotted as a function of her body weight ($n = 36$
 511 secondary queens from 18 colonies). The solid line represents the estimated regression
 512 line, $y_i = e^{(-1.052582 + 1.855425 * \log(x_i))}$. The y_i is the number of eggs produced in 24 h by the
 513 queen i , and x_i is the body weight of the queen i .
 514


515



516

517 **Figure 4** Correlation between the amount of daily egg production and the number of workers under
518 laboratory conditions. The number of workers in each colony is plotted as a function of the total
519 number of daily eggs laid in the colony ($n = 15$ colonies). The solid line represents the estimated
520 regression line, $w = e ^ { (0.8946205 + 1.411185 * \log(\sum y_i))}$. The w is the number of workers in the
521 colony and $\sum y_i$ is the total number of eggs produced in the colony in 24 h.
522

523



524

525 **Figure 5** Histogram of colony size in *Reticulitermes speratus* in the field. The x-axis represents the
526 estimated number of workers in each colony, and the y-axis represents the number of colonies. The
527 colony sizes were estimated for 188 field colonies. The estimated number of workers in a colony
528 ranged from 10,700 (first quartile) to 24,500 (median) to 58,600 (third quartile) to 451,800
529 (maximum).