

On the Arakawa lifting

Atsushi Murase

Faculty of Science, Kyoto Sangyo University

Hiro-aki Narita

Faculty of Science and Engineering, Waseda University

1 Introduction

For a reductive dual pair (H, G) , we have the theta lifting

$$\mathcal{L}: \{\text{automorphic forms on } H\} \rightarrow \{\text{automorphic forms on } G\}$$

defined by

$$\mathcal{L}(f)(g) = \int_{H_{\mathbb{Q}} \backslash H_{\mathbb{A}}} \theta(h, g) f(h) dh \quad (g \in G_{\mathbb{A}})$$

for an automorphic form f on H , where $\theta(h, g)$ is the theta kernel on $H_{\mathbb{Q}} \backslash H_{\mathbb{A}} \times G_{\mathbb{Q}} \backslash G_{\mathbb{A}}$. One of the fundametal problems is to characterize $\text{Ker}(\mathcal{L})$ and $\text{Im}(\mathcal{L})$ in terms of, say, Fourier expansions or L-values.

The kernel of $\mathcal{L}$ has been studied in many cases. For example, Böcherer and Schulze-Pillot ([BS]) and Hsieh and Namikawa ([HN1],[HN2]) proved the injectivity of the Yoshida lifting. On the other hand, the image of $\mathcal{L}$ is characterized in terms of automorphic L -functions in several cases. As a typical example, we recall the following results on the Saito-Kurokawa lifting due to Evdokimov and Oda.

Theorem 1.1 ([E],[O]). *Let F be a Hecke eigenform on $\text{Sp}_2(\mathbb{Z})$ of even weight. Denote by*

$$L^*(s, F; \text{spin}) = (2\pi)^{-s} \Gamma\left(s + k - \frac{3}{2}\right) \Gamma\left(s + \frac{1}{2}\right) \cdot L(s, F; \text{spin})$$

the completed spinor L -function of F with functional equation

$$L^*(s, F; \text{spin}) = L^*(1 - s, F; \text{spin}).$$

Then the following three conditions are equivalent.

- (1) F is in the Maass space (the image of the Saito-Kurokawa lifting).
- (2) $L^*(s, F; \text{spin})$ has a simple pole at $s = \frac{3}{2}$.
- (3) $L(s, F; \text{spin})$ does not vanish at $s = -\frac{1}{2}$.

In this note, we give a partial analogue of Theorem 1.1 for the *Arakawa lifting*, the theta lifting for the reductive dual pair $(O^*(2, 2), \text{Sp}(1, 1))$.

2 The Arakawa lifting

Let B be a definite quaternion algebra over $\mathbb{Q}$ of discriminant d_B . For $x \in B$, let $\text{tr}(x) = x + \bar{x}$ and $N(x) = x\bar{x}$ be the reduced trace and the reduced norm of x respectively, where $x \mapsto \bar{x}$ denotes the main involution of B . We fix an identification $B \otimes_{\mathbb{Q}} \mathbb{R} = \mathbb{H}$, where $\mathbb{H}$ is the Hamilton quaternion algebra.

We define two quaternion unitary groups by

$$H = \left\{ h \in \text{GL}_2(B) : \bar{t}hJh = J \right\}, \quad J = \begin{pmatrix} 0 & 1 \\ -1 & 0 \end{pmatrix},$$

$$G = \left\{ g \in \text{GL}_2(B) : \bar{t}gQg = Q \right\}, \quad Q = \begin{pmatrix} 0 & 1 \\ 1 & 0 \end{pmatrix}.$$

Then (H, G) forms a dual reductive pair. Note that $H = O^*(2, 2)$ is isogeneous to $\text{PGL}_2 \times (B^\times / \mathbb{Q}^\times)$ and that $G = \text{Sp}(1, 1)$ is an inner form of Sp_2 .

We henceforth fix a maximal order $\mathcal{O}$ of B . For a finite place p of $\mathbb{Q}$, set

$$L_p = \begin{cases} (\mathcal{O}_p, \mathcal{O}_p) & (p \nmid d_B) \\ (\mathcal{O}_p, \mathfrak{P}_p^{-1}) & (p \mid d_B) \end{cases},$$

$$L'_p = \begin{pmatrix} \mathcal{O}_p \\ \mathcal{O}_p \end{pmatrix},$$

where $\mathfrak{P}_p$ is the maximal ideal of $\mathcal{O}_p = \mathcal{O} \otimes_{\mathbb{Z}} \mathbb{Z}_p$. Let

$$K_f = \prod_{p < \infty} K_p, \quad K_p = \{k \in G_p \mid L_p k = L_p\},$$

$$U_f = \prod_{p < \infty} U_p, \quad U_p = \{u \in H_p \mid u L'_p = L'_p\}.$$

We fix an embedding $\mathbb{H}$ into $M_2(\mathbb{C})$. Let

$$K_\infty = \left\{ \begin{pmatrix} a & b \\ b & a \end{pmatrix} \mid a, b \in \mathbb{H}, a \pm b \in \mathbb{H}^1 \right\} \subset G_\infty,$$

$$U_\infty = \mathbb{H}^1 \cdot \mathrm{SO}(2, \mathbb{R}) \subset H_\infty,$$

where $\mathbb{H}^1 = \{x \in \mathbb{H} \mid \bar{x}x = 1\}$.

We henceforth suppose that $\kappa > 4$. Let σ_κ be the symmetric tensor representation of $M_2(\mathbb{C})$ on V_κ , the space of homogeneous polynomials in X and Y of total degree κ .

Let S_κ^G be the space of smooth functions $F: G_{\mathbb{Q}} \backslash G_{\mathbb{A}} \rightarrow V_\kappa$ satisfying

- (1) $F(gk_fk_\infty) = \sigma_\kappa(a+b)^{-1}F(g) \quad (g \in G_{\mathbb{A}}, k_f \in K_f, k_\infty = \begin{pmatrix} a & b \\ b & a \end{pmatrix} \in K_\infty).$
- (2) $F(g) = \frac{\kappa(\kappa-1)}{8\pi^2} \int_{G_\infty} \Omega(x_\infty) F(gx_\infty^{-1}) dx_\infty$, where $\Omega: G_\infty \rightarrow \mathrm{End}(V_\kappa)$ is the spherical function defined by $\Omega(g) = \sigma_\kappa(\Delta_g)^{-1} N(\Delta_g)^{-1}$ with $\Delta_g = \frac{1}{2}(a+b+c+d) \in \mathbb{H}^\times$ for $g = \begin{pmatrix} a & b \\ c & d \end{pmatrix} \in G_\infty$.
- (3) F is bounded.

Next let S_κ^H be the space of smooth functions $f: H_{\mathbb{Q}} \backslash H_{\mathbb{A}} \rightarrow V_\kappa$ satisfying

- (1) $f(hu_fu_\infty) = \sigma_\kappa(c + \sqrt{-1}d)^{-1}f(h) \quad (h \in H_{\mathbb{A}}, u_f \in U_f, u_\infty = \begin{pmatrix} a & b \\ c & d \end{pmatrix} \in U_\infty).$
- (2) $f(h) = \frac{\kappa-1}{2\pi} \int_{H_\infty} \omega(x_\infty) f(hx_\infty^{-1}) dx_\infty$, where $\omega: H_\infty \rightarrow \mathrm{End}(V_\kappa)$ is the spherical function defined by $\omega(h) = \sigma_\kappa(\delta_h)^{-1}$ with $\delta_h = \frac{1}{2}(a+d + \sqrt{-1}(-b+c))$ for $h = \begin{pmatrix} a & b \\ c & d \end{pmatrix} \in H_\infty$.

(3) f is bounded.

To define the theta kernel, we define the Weil representation r of $H_{\mathbb{A}} \times G_{\mathbb{A}}$ on $\mathcal{S}(B_{\mathbb{A}}^2) \otimes_{\mathbb{C}} \text{End}(V_{\kappa})$ by

$$\begin{aligned} r(h, 1)\varphi(X) &= \varphi({}^t\bar{h}X) \quad (h \in H_{\mathbb{A}}), \\ r\left(1, \begin{pmatrix} \alpha & 0 \\ 0 & {}^t\bar{\alpha}^{-1} \end{pmatrix}\right)\varphi(X) &= |N(\alpha)|^2 \varphi(X\alpha) \quad (\alpha \in B_{\mathbb{A}}^{\times}), \\ r\left(1, \begin{pmatrix} 1 & \beta \\ 0 & 1 \end{pmatrix}\right)\varphi(X) &= \psi(-\text{tr}(\beta {}^t\bar{X}JX)) \varphi(X) \quad (\beta \in B_{\mathbb{A}}, \beta + \bar{\beta} = 0), \end{aligned}$$

where ψ is the nontrivial additive character of $\mathbb{A}/\mathbb{Q}$ with $\psi(x_{\infty}) = \mathbf{e}(x_{\infty}) := e^{2\pi i x_{\infty}}$ for $x_{\infty} \in \mathbb{R}$, and $\mathcal{S}(B_{\mathbb{A}}^2)$ is the space of Schwartz-Bruhat functions on $B_{\mathbb{A}}^2$. The theta kernel $\theta: H_{\mathbb{Q}} \backslash H_{\mathbb{A}} \times G_{\mathbb{Q}} \backslash G_{\mathbb{A}} \rightarrow \text{End}(V_{\kappa})$ is defined by

$$\theta(h, g) = \sum_{X \in B^2} r(h, g)\varphi_0(X),$$

where

$$\varphi_0(X_f X_{\infty}) = \text{char}_{L_f}(X_f) \exp(-2\pi {}^t\bar{X}_{\infty} X_{\infty}) \sigma_{\kappa}((1, -\sqrt{-1})\bar{X}_{\infty})$$

for $X_f \in B_{\mathbb{A}_f}^2$ and $X_{\infty} \in \mathbb{H}^2$ and $L_f = \prod_{p < \infty} L_p$.

For $f \in \mathcal{S}_{\kappa}^H$, set

$$\mathcal{L}(f)(g) = \int_{H_{\mathbb{Q}} \backslash H_{\mathbb{A}}} \theta(h, g) f(h) dh \quad (g \in G_{\mathbb{A}}).$$

In [Ar], Arakawa proved the following result, which was later generalized to the case of $\text{Sp}(1, q)$ ($q \geq 1$) by the second named author ([Na]).

Theorem 2.1. *We have $\mathcal{L}(f) \in \mathcal{S}_{\kappa}^G$.*

We call $\mathcal{L}: \mathcal{S}_{\kappa}^H \rightarrow \mathcal{S}_{\kappa}^G$ the *Arakawa lifting*. The authors of this note studied arithmetic properties of $\mathcal{L}$ in [MN1], [MN2] and [MN3].

3 The main results

Let $\mathcal{L}^*: \mathcal{S}_\kappa^G \rightarrow \mathcal{S}_\kappa^H$ be the adjoint of $\mathcal{L}$ with respect to the Petersson inner products on $\mathcal{S}_\kappa^H$ and $\mathcal{S}_\kappa^G$. The main results of this note are given as follows.

Theorem 3.1 (Murase-Narita). *Suppose that $F \in \mathcal{S}_\kappa^G$ is a Hecke eigenform. Then $f = \mathcal{L}^*(F) \in \mathcal{S}_\kappa^H$ is a Hecke eigenform and we have*

$$L(F, s; \text{std}) = \zeta(s) L(f, s; \text{std}).$$

Here $L(F, s; \text{std})$ is the standard L -function of F of degree 5 and $L(f, s; \text{std})$ is the standard L -function of f of degree 4, and $\zeta(s)$ is the Riemann zeta function.

Theorem 3.2 (Murase-Narita). *Suppose that d_B is a prime number and let $F \in \mathcal{S}_\kappa^G$ be a Hecke eigenform.*

(1) *We have*

$$\mathcal{L}(\mathcal{L}^*F)(g) = c L(F, 0; \text{std}) F(g) \quad (g \in G_\mathbb{A}), \quad (3.1)$$

where c is an elementary constant.

(2) *We have an inner product formula:*

$$||\mathcal{L}^*(F)||^2 = c L(F, 0; \text{std}) \cdot ||F||^2,$$

and hence $\mathcal{L}^(F) = 0 \iff L(F, 0; \text{std}) = 0$.*

(3) *If $L(F, 0; \text{std}) \neq 0$, then $f := \mathcal{L}^*(F)$ does not vanish and F is the Arakawa lift of $(c \cdot L(F, 0; \text{std}))^{-1} f$. In particular $F \in \text{Im}(\mathcal{L})$.*

Remarks:

1. The proof of Theorem 3.2 is based on

- the doubling method,
- the Siegel-Weil formula for quaternion unitary groups due to Yamana ([Ya]),

- a trick using the spherical function Ω .
- 2. The assumption on d_B comes from this Siegel-Weil formula. When d_B is a composite, we need a modification of the assertion of Theorem 3.2 (we omit the detail in this note).
- 3. We have a formula for $\mathcal{L}^*(\mathcal{L}(f))$ similar to (3.1) for a Hecke eigenform f on H .
- 4. Let $\mathcal{L}': \{\text{Jacobi forms}\} \rightarrow \{\text{automorphic forms on } O(2, m)\}$ be a generalization of Maass lift studied independently by Gritsenko ([G]) and Sugano ([Su]). In [Su], Sugano showed that the lift is related to the theta lift for the reductice dual pair $(\text{SL}_2, O(2, m))$ studied by Oda and Rallis-Schiffmann, and proved a formula for $(\mathcal{L}')^* \circ \mathcal{L}'$ similar to (3.1) by a method different from ours.

References

- [Ar] T. Arakawa, unpublished notes.
- [BS] S. Böcherer and R. Schulze-Pillot: Siegel modular forms and theta series attached to quaternion algebras II, Nagoya Math. J. 147 (1997), 71–106, With errata to: Siegel modular forms and theta series attached to quaternion algebras, Nagoya Math. J. 121 (1991), 35–96.
- [E] S. A. Evdokimov, Characterization of the Maass space of Siegel modular cusp forms of genus 2, Mat. Sbor. 112 (1980), 133–142, 144.
- [G] V. A. Gritsenko, Fourier-Jacobi functions in n variables (Russian) Zap. Nauchn. Sem. Leningrad. Otdel. Mat. Inst. Steklov. (LOMI) 168, Anal. Teor. Chisel i Teor. Funktsii. 9(1988), 32–44, 187–188; translation in J. Soviet Math. 53 (1991), no.3, 243–252.
- [HN1] M-L. Hsieh and K. Namikawa, Bessel periods and the non-vanishing of Yoshida lifts modulo a prime, Math. Z. 285 (2017), no.3-4, 851–878.

- [HN2] M-L. Hsieh and K. Namikawa, Inner product formula for Yoshida lifts, *Ann. Math. Qué.* 42 (2018), no.2, 215–253.
- [MN1] A. Murase and H. Narita, Commutation relations of Hecke operators for Arakawa lifting, *Tohoku Math. J.* 60 (2008), 227–251.
- [MN2] A. Murase and H. Narita, Fourier expansion of Arakawa lifting I: An explicit formula and examples of non-vanishing lifts, *Israel J. Math.* 187 (2012), 317–369.
- [MN3] A. Murase and H. Narita, Fourier expansion of Arakawa lifting II: Relation with central L-values, *Internat. J. Math.* 27 (2016), 1650001.
- [Na] H. Narita, Theta lifting from elliptic cusp forms to automorphic forms on $\mathrm{Sp}(1, q)$, *Math. Z.* 259 (2008), 591–615.
- [O] T. Oda, On the poles of Andrianov L-functions, *Math. Ann.* 256 (1981), no.3, 323–340.
- [Su] T. Sugano, Jacobi forms and the theta lifting, *Comment. Math. Univ. St. Paul.* 44 (1995), no.1, 1–58.
- [Ya] S. Yamana, On the Siegel-Weil formula for quaternionic unitary groups, *Amer. J. Math.* 135 (2013), 1383–1432.

Atsushi Murase
 Department of Mathematical Science
 Faculty of Science
 Kyoto Sangyo University
 Motoyama, Kamigamo, Kita-ku
 Kyoto 603-8555
 JAPAN
 E-mail address: murase@cc.kyoto-su.ac.jp

Hiro-aki Narita
 Department of Mathematics
 Faculty of Science and Engineering
 Waseda University
 3-4-1 Okubo, Shinjuku-ku
 Tokyo 169-8555
 JAPAN
 E-mail address: hnarita@waseda.jp