

Reverse of Bebiano-Lemos-Providência inequality and Complementary Furuta inequality

大阪府立能勢高校 松本 明美 (Akemi Matsumoto)
Nose Senior Highschool
茨城大学工学部 中本 律男 (Rituo Nakamoto)
Department of Engineering, Ibaraki University
大阪教育大学 藤井 正俊 (Masatoshi Fujii)
Osaka Kyoiku University

We give a simultaneous extension of Bebiano-Lemos-Providência inequality and Araki-Cordes one: Let A and B be positive operators. Then for each $r \geq 0$

$$\|A^{\frac{r+t}{2}} B^t A^{\frac{r+t}{2}}\| \leq \|A^{\frac{r}{2}} (A^{\frac{s}{2}} B^s A^{\frac{s}{2}})^{\frac{t}{s}} A^{\frac{r}{2}}\| \quad \text{for } s \geq t \geq 0.$$

In succession, we prove a reverse inequality: Let A and B be positive operators. Then for each $r \geq 0$

$$\|A^{\frac{r+t}{2}} B^t A^{\frac{r+t}{2}}\| \geq \|A^{\frac{r}{2}} (A^{\frac{s}{2}} B^s A^{\frac{s}{2}})^{\frac{t}{s}} A^{\frac{r}{2}}\| \quad \text{for } t \geq s \geq r \geq 0.$$

Furthermore, we discuss reverse of generalized BLP inequality in a general setting, in which we point out that the restriction $t \geq s \geq r$ in the above is quite reasonable.

1 Introduction

Let A be a bounded linear operator acting on a Hilbert space H . Then A is positive, denoted by $A \geq 0$, if $(Ax, x) \geq 0$ for all $x \in H$. In particular, $A > 0$ means that A is invertible and positive.

Recently, Bebiano-Lemos-Providência showed an interesting norm inequality (BLP): Let A and B be positive operators. Then

$$(1) \quad \|A^{\frac{1+t}{2}} B^t A^{\frac{1+t}{2}}\| \leq \|A^{\frac{1}{2}} (A^{\frac{s}{2}} B^s A^{\frac{s}{2}})^{\frac{t}{s}} A^{\frac{1}{2}}\| \quad \text{for } s \geq t \geq 0.$$

If we delete $A^{\frac{1}{2}}$ on both sides in (1), we have the Araki-Cordes inequality (AC)

$$(2) \quad \|A^p B^p A^p\| \leq \|ABA\|^p \quad \text{for } 0 \leq p \leq 1.$$

In this sense, BLP inequality (1) is regarded as an extension of AC inequality (2).

In this note, we first give a simultaneous extension of (1) and (2) as follows generalized BLP inequality (GBLP): Let A and B be positive operators. Then for each $r \geq 0$

$$(3) \quad \|A^{\frac{r+t}{2}} B^t A^{\frac{r+t}{2}}\| \leq \|A^{\frac{r}{2}} (A^{\frac{s}{2}} B^s A^{\frac{s}{2}})^{\frac{t}{s}} A^{\frac{r}{2}}\| \quad \text{for } s \geq t \geq 0.$$

Next we discuss a reverse inequality of (3) (R-GBLP): Let A and B be positive operators. Then for each $r \geq 0$,

$$(4) \quad \|A^{\frac{r+t}{2}} B^t A^{\frac{r+t}{2}}\| \geq \|A^{\frac{r}{2}} (A^{\frac{s}{2}} B^s A^{\frac{s}{2}})^{\frac{t}{s}} A^{\frac{r}{2}}\| \quad \text{for } t \geq s \geq r \text{ and } s > 0.$$

As a corollary, the case $r = 1$ in (4) corresponds to the reverse of BLP(R-BLP) inequality and the case $r = 0$ in (4) corresponds to the reverse of AC inequality (2) (R-AC).

2 A generalization of BLP inequality

In this section, we generalize BLP inequality (1). For this, we cite Furuta inequality: Let $A \geq B$ for A and B be positive operators. Then for each $r \geq 0$.

$$(A^{\frac{r}{2}} B^p A^{\frac{r}{2}})^{\frac{1}{q}} \leq (A^{\frac{r}{2}} A^p A^{\frac{r}{2}})^{\frac{1}{q}} \quad \text{for } p \geq 0, \quad q \geq 1 \quad \text{with } (1+r)q \geq p+r.$$

We prove the following theorem including BLP inequality and AC one.

Theorem. 1. *Let A and B be positive operators. Then for each $r \geq 0$*

$$\| A^{\frac{r+t}{2}} B^t A^{\frac{r+t}{2}} \| \leq \| A^{\frac{r}{2}} (A^{\frac{t}{2}} B^s A^{\frac{t}{2}})^{\frac{1}{s}} A^{\frac{r}{2}} \| \quad \text{for } s \geq t \geq 0.$$

Proof. Since this inequality is AC inequality when $r = 0$, we may assume $r > 0$. It suffices to prove that

$$A^{\frac{r}{2}} (A^{\frac{t}{2}} B^s A^{\frac{t}{2}})^{\frac{1}{s}} A^{\frac{r}{2}} \leq 1 \implies A^{\frac{r+t}{2}} B^t A^{\frac{r+t}{2}} \leq 1 \quad \text{for } s \geq t \geq 0.$$

If $A^{\frac{r}{2}} (A^{\frac{t}{2}} B^s A^{\frac{t}{2}})^{\frac{1}{s}} A^{\frac{r}{2}} \leq 1$, then $(A^{\frac{t}{2}} B^s A^{\frac{t}{2}})^{\frac{1}{s}} \leq A^{-r}$. We put

$$A_1 = A^{-r}, \quad B_1 = (A^{\frac{t}{2}} B^s A^{\frac{t}{2}})^{\frac{1}{s}} \quad r_1 = \frac{s}{r} \quad \text{and} \quad p = q = \frac{s}{t}.$$

Then

$$A_1 \geq B_1 \geq 0, \quad r_1 \geq 0, \quad p = q \geq 1 \quad \text{and} \quad (1+r_1)q \geq p+r_1,$$

so that Furuta inequality implies

$$(A_1^{\frac{r_1}{2}} B_1^p A_1^{\frac{r_1}{2}})^{\frac{1}{q}} \leq (A_1^{\frac{r_1}{2}} A_1^p A_1^{\frac{r_1}{2}})^{\frac{1}{q}},$$

that is,

$$(A_1^{\frac{r_1}{2}} B_1^{\frac{t}{s}} A_1^{\frac{r_1}{2}})^{\frac{1}{s}} \leq (A_1^{\frac{r_1}{2}} A_1^{\frac{t}{s}} A_1^{\frac{r_1}{2}})^{\frac{1}{s}}.$$

Since we have

$$A = A_1^{-\frac{1}{r}}, \quad B = (A^{-\frac{t}{2}} B_1^{\frac{t}{s}} A^{-\frac{t}{2}})^{\frac{1}{s}} = (A_1^{\frac{r_1}{2}} B_1^{\frac{t}{s}} A_1^{\frac{r_1}{2}})^{\frac{1}{s}},$$

it follows that

$$\begin{aligned} A^{\frac{r+t}{2}} B^t A^{\frac{r+t}{2}} &= (A_1^{-\frac{1}{r}})^{\frac{r+t}{2}} (A_1^{\frac{r_1}{2}} B_1^{\frac{t}{s}} A_1^{\frac{r_1}{2}})^{\frac{1}{s}} (A_1^{-\frac{1}{r}})^{\frac{r+t}{2}} \\ &\leq A_1^{-\frac{r+t}{2r}} (A_1^{-\frac{t}{2}} A_1^{\frac{t}{s}} A_1^{-\frac{t}{2}})^{\frac{1}{s}} A_1^{-\frac{r+t}{2r}} = A_1^{-\frac{r+t}{2r}} A_1^{\frac{r+t}{r}} A_1^{-\frac{r+t}{2r}} = I. \end{aligned}$$

□

Remark. It is obvious that the case $r = 1$ in Theorem 1 is just the BLP inequality and the case $r = 0$ is AC one.

3 Reverse inequalities

In this section, we show a reverse inequality of Theorem 1, in which we use the well-known formula (Lemma of Furuta),

$$(5) \quad (A^{\frac{1}{2}}BA^{\frac{1}{2}})^p = A^{\frac{1}{2}}B^{\frac{1}{2}}(B^{\frac{1}{2}}AB^{\frac{1}{2}})^{p-1}B^{\frac{1}{2}}A^{\frac{1}{2}} \quad \text{for } p \geq 1$$

and Löwner-Heinz inequality (LH)

$$(6) \quad A \geq B \geq 0 \quad \text{implies} \quad A^p \geq B^p \quad \text{for all } 0 \leq p \leq 1.$$

Theorem. 2. *Let A and B be positive operators. Then for each $r \geq 0$*

$$(7) \quad \|A^{\frac{r+t}{2}}B^tA^{\frac{r+t}{2}}\| \geq \|A^{\frac{r}{2}}(A^{\frac{1}{2}}B^sA^{\frac{1}{2}})^{\frac{t}{s}}A^{\frac{r}{2}}\| \quad \text{for } t \geq s \geq r \quad \text{and } s > 0.$$

Proof. It suffices to prove that

$$A^{\frac{r+t}{2}}B^tA^{\frac{r+t}{2}} \leq 1 \implies A^{\frac{r}{2}}(A^{\frac{1}{2}}B^sA^{\frac{1}{2}})^{\frac{t}{s}}A^{\frac{r}{2}} \leq 1 \quad \text{for } t \geq s \geq r \quad \text{and } s > 0.$$

Suppose that $\lfloor \frac{t}{s} \rfloor = 2l - 1$ or $2l$ for some natural number l . Then following equations are obtained by (5).

$$\begin{aligned} & A^{\frac{r}{2}}(A^{\frac{1}{2}}B^sA^{\frac{1}{2}})^{\frac{t}{s}}A^{\frac{r}{2}} \\ &= A^{\frac{r+t}{2}}B^{\frac{t}{s}}(B^{\frac{1}{2}}A^sB^{\frac{1}{2}})^{\frac{t}{s}-1}B^{\frac{1}{2}}A^{\frac{r+t}{2}} \\ &= A^{\frac{r+t}{2}}B^sA^{\frac{r}{2}}(A^{\frac{1}{2}}B^sA^{\frac{1}{2}})^{\frac{t}{s}-2}A^{\frac{1}{2}}B^sA^{\frac{r+t}{2}} \\ &= A^{\frac{r+t}{2}}(B^sA^s)B^{\frac{t}{s}}(B^{\frac{1}{2}}A^sB^{\frac{1}{2}})^{\frac{t}{s}-3}B^{\frac{1}{2}}(A^sB^s)A^{\frac{r+t}{2}} \\ &= A^{\frac{r+t}{2}}(B^sA^s)B^sA^{\frac{r}{2}}(A^{\frac{1}{2}}B^sA^{\frac{1}{2}})^{\frac{t}{s}-4}A^{\frac{1}{2}}B^s(A^sB^s)A^{\frac{r+t}{2}} \\ &= A^{\frac{r+t}{2}}(B^sA^s)(B^sA^s)B^{\frac{t}{s}}(B^{\frac{1}{2}}A^sB^{\frac{1}{2}})^{\frac{t}{s}-5}B^{\frac{1}{2}}(A^sB^s)(A^sB^s)A^{\frac{r+t}{2}} \\ &\vdots \\ &= A^{\frac{r+t}{2}} \overbrace{(B^sA^s) \cdots (B^sA^s)}^{(l-1)\text{-times}} B^{\frac{t}{s}} (B^{\frac{1}{2}}A^sB^{\frac{1}{2}})^{\frac{t}{s}-(2l-1)} B^{\frac{1}{2}} \overbrace{(A^sB^s) \cdots (A^sB^s)}^{(l-1)\text{-times}} A^{\frac{r+t}{2}} \end{aligned}$$

Since $t \geq s \geq r \geq 0$, we have

$$0 \leq \frac{s}{r+t} \leq 1, \quad 0 \leq \frac{s}{t} \leq 1, \quad 0 \leq \frac{t(r+s) - 2(l-h)rs}{t(r+t)} \leq 1 \quad \text{for } h = 1, 2, \dots, l$$

and

$$0 \leq \frac{t(s-r) + 2(l-k)rs}{t(r+t)} \leq 1 \quad \text{for } k = 0, 1, \dots, l-1.$$

We suppose that $A^{\frac{r+t}{2}}B^tA^{\frac{r+t}{2}} \leq 1$, that is,

$$B^t \leq A^{-(r+t)} \quad \text{and} \quad A^{(r+t)} \leq B^{-t}.$$

So the Löwner-Heinz inequality (6) implies that

$$A^s \leq B^{-\frac{st}{r+t}}, \quad B^s \leq A^{-\frac{s(r+t)}{t}}, \quad B^{\frac{t(r+s)-2(l-h)rs}{(r+t)}} \leq A^{-\frac{t(r+s)-2(l-h)rs}{t}} \quad \text{for } h = 1, 2, \dots, l$$

and

$$A^{\frac{t(s-r)+2(l-k)rs}{t}} \leq B^{-\frac{t(s-r)+2(l-k)rs}{(r+t)}} \quad \text{for } k = 0, 1, \dots, l-1.$$

Now we assume that $\lfloor \frac{t}{s} \rfloor = 2l - 1$ for some natural number l . Since $0 \leq \frac{t}{s} - (2l - 1) \leq 1$, we have

$$\begin{aligned} & A^{\frac{r}{2}} (A^{\frac{s}{2}} B^s A^{\frac{s}{2}})^{\frac{t}{s}} A^{\frac{r}{2}} \\ &= A^{\frac{r}{2}} \overbrace{(B^s A^s) \cdots (B^s A^s)(B^s A^s)}^{(l-1)\text{-times}} B^{\frac{s}{2}} (B^{\frac{s}{2}} A^s B^{\frac{s}{2}})^{\frac{t}{s} - (2l-1)} B^{\frac{s}{2}} \overbrace{(A^s B^s)(A^s B^s) \cdots (A^s B^s)}^{(l-1)\text{-times}} A^{\frac{r}{2}} \\ &\leq A^{\frac{r}{2}} (B^s A^s) \cdots (B^s A^s)(B^s A^s) B^{\frac{s}{2}} (B^{\frac{s}{2}} B^{-\frac{st}{r+t}} B^{\frac{s}{2}})^{\frac{t}{s} - (2l-1)} B^{\frac{s}{2}} (A^s B^s)(A^s B^s) \cdots (A^s B^s) A^{\frac{r}{2}} \\ &= A^{\frac{r}{2}} (B^s A^s) \cdots (B^s A^s)(B^s A^s) B^{\frac{t(s+r)-2(l-1)rs}{r+t}} (A^s B^s)(A^s B^s) \cdots (A^s B^s) A^{\frac{r}{2}} \\ &= A^{\frac{r}{2}} \overbrace{(B^s A^s) \cdots (B^s A^s)}^{(l-2)\text{-times}} B^s A^s B^{\frac{t(s+r)-2(l-1)rs}{r+t}} A^s B^s \overbrace{(A^s B^s) \cdots (A^s B^s)}^{(l-2)\text{-times}} A^{\frac{r}{2}} \\ &\leq A^{\frac{r}{2}} (B^s A^s) \cdots (B^s A^s) B^s A^{\frac{t(s-r)+2(l-1)rs}{t}} B^s (A^s B^s) \cdots (A^s B^s) A^{\frac{r}{2}} \\ &\leq A^{\frac{r}{2}} (B^s A^s) \cdots (B^s A^s) B^{\frac{t(s+r)-2(l-2)rs}{r+t}} (A^s B^s) \cdots (A^s B^s) A^{\frac{r}{2}} \\ &\vdots \\ &\leq A^{\frac{r}{2}} B^s A^{\frac{t(s-r)+2(l-(l-1))rs}{t}} B^s A^{\frac{r}{2}} \\ &\leq A^{\frac{r}{2}} B^{\frac{ts+rt}{r+t}} A^{\frac{r}{2}} \\ &\leq A^{\frac{r}{2}} A^{-(r+s)} A^{\frac{r}{2}} \\ &= I. \end{aligned}$$

On the other hand, we assume that $\lfloor \frac{t}{s} \rfloor = 2l$ for some natural number l . Since $0 \leq \frac{t}{s} - 2l \leq 1$, similarly we have the following, in which the first equality is ensured in the first paragraph.

$$\begin{aligned} & A^{\frac{r}{2}} (A^{\frac{s}{2}} B^s A^{\frac{s}{2}})^{\frac{t}{s}} A^{\frac{r}{2}} \\ &= A^{\frac{r}{2}} \overbrace{(B^s A^s) \cdots (B^s A^s)(B^s A^s)}^{(l-1)\text{-times}} B^{\frac{s}{2}} (B^{\frac{s}{2}} A^s B^{\frac{s}{2}})^{\frac{t}{s} - (2l-1)} B^{\frac{s}{2}} \overbrace{(A^s B^s)(A^s B^s) \cdots (A^s B^s)}^{(l-1)\text{-times}} A^{\frac{r}{2}} \\ &= A^{\frac{r}{2}} \overbrace{(B^s A^s) \cdots (B^s A^s)(B^s A^s)}^{(l-1)\text{-times}} B^s A^{\frac{s}{2}} (A^{\frac{s}{2}} B^s A^{\frac{s}{2}})^{\frac{t}{s} - 2l} A^{\frac{s}{2}} B^s \overbrace{(A^s B^s)(A^s B^s) \cdots (A^s B^s)}^{(l-1)\text{-times}} A^{\frac{r}{2}} \\ &\leq A^{\frac{r}{2}} (B^s A^s) \cdots (B^s A^s)(B^s A^s) B^s A^{\frac{s}{2}} (A^{\frac{s}{2}} A^{-\frac{s(r+t)}{t}} A^{\frac{s}{2}})^{\frac{t}{s} - 2l} A^{\frac{s}{2}} B^s (A^s B^s)(A^s B^s) \cdots (A^s B^s) A^{\frac{r}{2}} \\ &= A^{\frac{r}{2}} (B^s A^s) \cdots (B^s A^s)(B^s A^s) B^s A^{\frac{t(s-r)+2ls}{t}} B^s (A^s B^s)(A^s B^s) \cdots (A^s B^s) A^{\frac{r}{2}} \\ &\leq A^{\frac{r}{2}} (B^s A^s) \cdots (B^s A^s)(B^s A^s) B^s B^{-\frac{t(s-r)+2lrs}{r+s}} B^s (A^s B^s)(A^s B^s) \cdots (A^s B^s) A^{\frac{r}{2}} \\ &= A^{\frac{r}{2}} \overbrace{(B^s A^s) \cdots (B^s A^s)}^{(l-2)\text{-times}} B^s A^s B^{\frac{t(r+s)-2(l-1)rs}{r+t}} A^s B^s \overbrace{(A^s B^s) \cdots (A^s B^s)}^{(l-2)\text{-times}} A^{\frac{r}{2}} \\ &\vdots \\ &\leq I. \end{aligned}$$

Hence the proof is complete. $\square$

4 Complementary Furuta inequality

In this section, we consider **R-GBLP**, in which Kamei's theorem(**Theorem K**) on complements of Furuta inequality corresponds to our result. So now recall it due to Kamei.

Theorem K. If $A \geq B > 0$, then for $0 < p \leq \frac{1}{2}$

$$(8) \quad A^t \natural_{\frac{2p-t}{p-t}} B^p \leq A^{2p} \quad \text{for } 0 \leq t \leq p$$

and for $\frac{1}{2} \leq p \leq 1$

$$(9) \quad A^t \natural_{\frac{1-t}{p-t}} B^p \leq A \quad \text{for } 0 \leq t \leq p.$$

Here $\natural_q$ for $q \notin [0, 1]$ has been used as

$$A \natural_q B := A^{\frac{1}{2}} (A^{-\frac{1}{2}} B A^{-\frac{1}{2}})^q A^{\frac{1}{2}} \quad \text{for } A, B > 0.$$

First we prove the following Theorem.

Theorem. 3. Let $A, B \geq 0$ and $0 < p \leq 1$. Then

$$(10) \quad \| A^{\frac{1+s}{2}} B^{1+s} A^{\frac{1+s}{2}} \|_{\frac{p+s}{p(1+s)}} \geq \| A^{\frac{1}{2}} (A^{\frac{s}{2}} B^{p+s} A^{\frac{s}{2}})^{\frac{1}{p}} A^{\frac{1}{2}} \|$$

for all $s \geq 0$ with $s \geq 1 - 2p$.

Proof. It suffices to show that

$$(11) \quad B^{1+s} \leq A^{-(1+s)} \Rightarrow A^{\frac{1}{2}} (A^{\frac{s}{2}} B^{p+s} A^{\frac{s}{2}})^{\frac{1}{p}} A^{\frac{1}{2}} \leq 1$$

for $0 < p \leq 1$ and $s \geq 0$ with $s \geq 1 - 2p$. So we put

$$A_1 = A^{-(1+s)}, \quad B_1 = B^{1+s}.$$

Then (11) is rephrased as

$$A_1 \geq B_1 > 0 \Rightarrow A_1^{\frac{1}{1+s}} \natural_{\frac{1}{p}} B_1^{\frac{p+s}{1+s}} \leq A_1.$$

for $0 < p \leq 1$ and $s \geq 0$ with $s \geq 1 - 2p$. Moreover if we replace

$$t_1 = \frac{s}{1+s}, \quad p_1 = \frac{p+s}{1+s},$$

then we have $\frac{1-t_1}{p_1-t_1} = \frac{1}{p}$, and $\frac{1}{2} \leq p_1 (\leq 1)$ if and only if $1 - 2p \leq s$, so that (11) has the following equivalent expression:

$$A_1 \geq B_1 > 0 \Rightarrow A_1^{t_1} \natural_{\frac{1-t_1}{p_1-t_1}} B_1^p \leq A_1 \quad \text{for } 0 \leq t_1 < p_1.$$

Fortunately, since $\frac{1}{2} \leq p_1 \leq 1$, it has been ensured by Theorem A due to Kamei. $\square$

Next we show a reverse inequality of BLP inequality (**R-BLP**) is obtained as colloary of Theorem 3.

Proof of R-BLP We put $p = \frac{s}{t}$ for $t \geq s \geq 0$. Then we have $1 - 2p \leq s$ if and only if $\frac{t}{t+2} \leq s$. Since $s \geq 1$ is assumed, $\frac{t}{t+2} \leq s$ holds for arbitrary $t > 0$, so that Theorem 3 is applicable.

Now we take $B_1 = B^{\frac{1+t}{t}}$, i.e., $B = B_1^{\frac{t}{1+t}}$. Then Araki-Cordes inequality and Theorem 3 imply that

$$\begin{aligned} \| A^{\frac{1+t}{2}} B_1^t A^{\frac{1+t}{2}} \| &\geq \| A^{\frac{1+s}{2}} B_1^{\frac{t(1+s)}{1+t}} A^{\frac{1+s}{2}} \| \frac{1+t}{1+s} = \| A^{\frac{1+s}{2}} B^{1+s} A^{\frac{1+s}{2}} \| \frac{2+s}{2(1+s)} \\ &\geq \| A^{\frac{1}{2}} (A^{\frac{s}{2}} B^{p+s} A^{\frac{s}{2}})^{\frac{1}{p}} A^{\frac{1}{2}} \| = \| A^{\frac{1}{2}} (A^{\frac{s}{2}} B_1^s A^{\frac{s}{2}})^{\frac{1}{p}} A^{\frac{1}{2}} \|, \end{aligned}$$

as desired.

R-BLP is generalized a bit as follows:

Corollary. For $A, B > 0$ and $r \geq 0$

$$(12) \quad \| A^{\frac{r+t}{2}} B^t A^{\frac{r+t}{2}} \| \geq \| A^{\frac{r}{2}} (A^{\frac{s}{2}} B^s A^{\frac{s}{2}})^{\frac{1}{p}} A^{\frac{r}{2}} \|$$

holds for all $t \geq s \geq r$.

Proof. It is proved by applying **R-BLP** to $A_1 = A^r$, $B_1 = B^r$ and $t_1 = \frac{t}{r}$, $s_1 = \frac{s}{r}$. $\square$

Finally we consider a reverse inequality of generalized BLP inequality which corresponds to another Kamei's complement (7): If $A \geq B > 0$, then for $0 < p \leq \frac{1}{2}$

$$A^t \natural_{\frac{2p-t}{p-t}} B^p \leq A^{2p} \quad \text{for } 0 \leq t < p.$$

Theorem. 4. Let $A, B \geq 0$ and $0 < p \leq \frac{1}{2}$. Then

$$(13) \quad \| A^{\frac{1+s}{2}} B^{1+s} A^{\frac{1+s}{2}} \| \frac{(2p+s)(p+s)}{p(1+s)} \geq \| A^{p+\frac{s}{2}} (A^{\frac{s}{2}} B^{p+s} A^{\frac{s}{2}})^{\frac{2p+s}{p}} A^{p+\frac{s}{2}} \|$$

for all $0 \leq s \leq 1 - 2p$.

Proof. A proof is quite similar to that of Theorem 3. We put

$$A_1 = A^{-(1+s)}, \quad B_1 = B^{1+s}; \quad t_1 = \frac{s}{1+s}, \quad p_1 = \frac{p+s}{1+s}.$$

Then (7) gives us that

$$A_1 \geq B_1 > 0 \Rightarrow A_1^{t_1} \natural_{\frac{2p_1-t_1}{p_1-t_1}} B_1^{p_1} \leq A_1^{2p_1},$$

for $0 \leq t_1 < p_1 \leq \frac{1}{2}$, so that

$$A^{-(1+s)} \geq B^{1+s} \Rightarrow A^{-s} \natural_{\frac{2p+s}{p}} B^{p+s} \leq A^{-2(p+s)}$$

for $0 \leq s \leq 1 - 2p$. Obviously it implies the desired norm inequality (12). $\square$


5 Remarks

We first consider a relation between Furuta inequality and Theorem K. Furuta inequality has the following representation by α -geometric mean $\sharp_\alpha$: For $A \geq B \geq 0$


$$A^t \sharp_{\frac{1-t}{p-t}} B^p \leq A \quad \text{for } p \geq 1, t \leq 0$$

where $A \sharp_\alpha B$ for $0 \leq \alpha \leq 1$ is defined by

$$A \sharp_\alpha B := A^{\frac{1}{2}} (A^{-\frac{1}{2}} B A^{-\frac{1}{2}})^\alpha A^{\frac{1}{2}} \quad \text{for } A, B > 0.$$



As a consequence we have relations among the inequalities discussed above.



References

- [1] H. Araki, *On an inequality of Lieb and Thirring*, Lett. Math. Phys., **19**(1990), 167–170.
- [2] N. Bebiano, R. Lemos and J. Providência, *Inequalities for quantum relative entropy*, Linear Algebra Appl. **401**(2005), 159–172.
- [3] H.O. Cordes, *Spectral Theory of Linear Differential Operators and Comparison Algebras*, Cambridge University Press, 1987.
- [4] M. Fujii, *Furuta's inequality and its mean theoretic approach*, J. Operator theory, **23**(1990), 67–72.
- [5] M. Fujii, R. Nakamoto and M. Tominaga, *Generalized Bebiano-Lemos-Providência inequalities and their reverses*, Linear Algebra Appl., to appear.
- [6] T. Furuta, $A \geq B \geq 0$ assures $(B^r A^p B^r)^{1/q} \geq B^{(p+2r)/q}$ for $r \geq 0, p \geq 0, q \geq 1$ with $(1+2r)q \geq p+2r$, Proc. Amer. Math. Soc., **101**(1987), 85–88.
- [7] T. Furuta, *Elementary proof of an order preserving inequality*, Proc. Japan Acad., **65**(1989), 126.
- [8] T. Furuta, J. Mićić, J.E. Pečarić and Y. Seo, *Mond-Pečarić Method in Operator Inequalities*, Monographs in Inequalities 1, Element, Zagreb, 2005.
- [9] E. Kamei, *A satellite to Furuta's inequality*, Math. Japon., **33**(1988), 883–886.
- [10] E. Kamei, *Complements to Furuta inequality, II*, Math. Japon., **45**(1997), 15–23.
- [11] A. Matsumoto and R. Nakamoto, *Reverse of Bebiano-Lemos-Providência inequality*, preprint.
- [12] A. Matsumoto and M. Fujii, *Generalizations of reverse of Bebiano-Lemos-Providência inequality*, preprint.
- [13] G.K. Pedersen, *Some operator monotone functions*, Proc. Amer. Math. Soc., **36**(1972), 309–310.
- [14] K. Tanahashi, *Best possibility of the Furuta inequality*, Proc. Amer. Math. Soc., **124**(1996), 141–146.